

# Ambavirumab/romlusevimab for COVID-19: real-time meta analysis of 3 studies

@CovidAnalysis, June 2024, Version 1  
<https://c19early.org/ammeta.html>

## Abstract

Statistically significant lower risk is seen for viral clearance. 2 studies from 2 independent teams (both from the same country) show significant improvements.



Meta analysis using the most serious outcome reported shows 11% [-154-69%] lower risk, without reaching statistical significance. Results are better for Randomized Controlled Trials and higher quality studies. Early treatment shows efficacy while late treatment does not, consistent with expectations for an antiviral treatment.

Currently there is limited data, with only 29 control events for the most serious outcome in trials to date. Studies to date are from only 3 different groups.

**Efficacy is variant dependent.** mAb use may create new variants that spread globally<sup>1,2</sup>, and may be associated with prolonged viral loads, clinical deterioration, and immune escape<sup>2-4</sup>.

No treatment or intervention is 100% effective. All practical, effective, and safe means should be used based on risk/benefit analysis. Multiple treatments are typically used in combination, and other treatments are significantly more effective.

All data to reproduce this paper and sources are in the appendix.




## AMBAVIRUMAB/ROMLUSEVIMAB FOR COVID-19 – HIGHLIGHTS

Ambavirumab/romlusevimab reduces risk with low confidence for viral clearance. **Efficacy is variant dependent.**


Outcome specific analyses and combined evidence from all studies, incorporating treatment delay, a primary confounding factor.

Real-time updates and corrections, transparent analysis with all results in the same format, consistent protocol for 72 treatments.

### 3 ambavirumab COVID-19 studies



### Timeline of COVID-19 ambavirumab studies (pooled effects)



**Figure 1. A. Random effects meta-analysis.** This plot shows pooled effects, see the specific outcome analyses for individual outcomes. Analysis validating pooled outcomes for COVID-19 can be found below. Effect extraction is pre-specified, using the most serious outcome reported. For details see the appendix. **B. Timeline of results in ambavirumab studies.**


## Introduction

**Immediate treatment recommended.** SARS-CoV-2 infection primarily begins in the upper respiratory tract and may progress to the lower respiratory tract, other tissues, and the nervous and cardiovascular systems, which may lead to cytokine storm, pneumonia, ARDS, neurological injury<sup>5-10</sup> and cognitive deficits<sup>7</sup>, cardiovascular complications<sup>11</sup>, organ failure, and death. Minimizing replication as early as possible is recommended.

**Many treatments are expected to modulate infection.** SARS-CoV-2 infection and replication involves the complex interplay of 50+ host and viral proteins and other factors<sup>A,12-16</sup>, providing many therapeutic targets for which many existing compounds have known activity. Scientists have predicted that over 7,000 compounds may reduce COVID-19 risk<sup>17</sup>, either by directly minimizing infection or replication, by supporting immune system function, or by minimizing secondary complications.

**Analysis.** We analyze all significant controlled studies of ambavirumab for COVID-19. Search methods, inclusion criteria, effect extraction criteria (more serious outcomes have priority), all individual study data, PRISMA answers, and statistical methods are detailed in Appendix 1. We present random effects meta-analysis results for all studies, studies within each treatment stage, individual outcomes, Randomized Controlled Trials (RCTs), and higher quality studies.

**Treatment timing.** Figure 2 shows stages of possible treatment for COVID-19. Prophylaxis refers to regularly taking medication before becoming sick, in order to prevent or minimize infection. Early Treatment refers to treatment immediately or soon after symptoms appear, while Late Treatment refers to more delayed treatment.



**Figure 2.** Treatment stages.

## Variant Dependence

Extensive mutations in SARS-CoV-2 have resulted in variants that evade neutralizing antibodies from monoclonal antibody treatments<sup>18</sup>, resulting in efficacy that is highly variant dependent. Table 1 shows efficacy by variant for several monoclonal antibodies. This table covers earlier SARS-CoV-2 variants and has not been updated for more recent variants and more recent monoclonal antibodies.

	<i>Bamlanivimab/etesevimab</i>	<i>Casirivimab/imdevimab</i>	<i>Sotrovimab</i>	<i>Bebtelovimab</i>	<i>Tixagevimab/cilgavimab</i>
Alpha B.1.1.7	likely effective	likely effective	likely effective	likely effective	likely effective
Beta/Gamma BA1.351/P.1	likely ineffective	likely effective	likely effective	likely effective	likely effective
Delta B.1.617.2	likely effective	likely effective	likely effective	likely effective	likely effective
Omicron BA.1/BA.1.1	likely ineffective	likely ineffective	likely effective	likely effective	unknown
Omicron BA.2	likely ineffective	likely ineffective	likely ineffective	likely effective	likely effective
Omicron BA.5	likely ineffective	likely ineffective	likely ineffective	likely effective	likely effective
Omicron BA.4.6	likely ineffective	likely ineffective	likely ineffective	likely effective	likely ineffective
Omicron BQ.1.1	likely ineffective	likely ineffective	likely ineffective	likely ineffective	likely ineffective

**Table 1.** Predicted efficacy by variant from *Davis et al.* (not updated for more recent variants). ■: likely effective ■: likely ineffective ■: unknown. Submit updates.

## Results


Table 2 summarizes the results for all stages combined, for Randomized Controlled Trials, after exclusions, and for specific outcomes. Table 3 shows results by treatment stage. Figure 3 plots individual results by treatment stage. Figure 4, 5, 6, 7, 8, and 9 show forest plots for random effects meta-analysis of all studies with pooled effects, mortality results, ICU admission, hospitalization, recovery, and viral clearance.

	Improvement	Studies	Patients	Authors
All studies	11% [-154-69%]	3	1,531	91
After exclusions	56% [-319-95%]	2	1,191	85
Randomized Controlled Trials	56% [-319-95%]	2	1,191	85
Mortality	11% [-154-69%]	3	1,531	91
Hospitalization	48% [-74-85%]	2	1,177	24
RCT mortality	56% [-319-95%]	2	1,191	85

**Table 2.** Random effects meta-analysis for all stages combined, for Randomized Controlled Trials, after exclusions, and for specific outcomes. Results show the percentage improvement with treatment and the 95% confidence interval. \*  $p < 0.05$   
\*\*  $p < 0.01$  \*\*\*\*  $p < 0.0001$ .

	Early treatment	Late treatment
All studies	89% [12-99%] *	-34% [-135-24%]
After exclusions	89% [12-99%] *	-15% [-141-46%]
Randomized Controlled Trials	89% [12-99%] *	-15% [-141-46%]
Mortality	89% [12-99%] *	-34% [-135-24%]
Hospitalization	73% [50-86%] ****	8% [3-13%] **
RCT mortality	89% [12-99%] *	-15% [-141-46%]


**Table 3.** Random effects meta-analysis results by treatment stage. Results show the percentage improvement with treatment, the 95% confidence interval, and the number of studies for the stage. \*  $p < 0.05$   
\*\*  $p < 0.01$  \*\*\*\*  $p < 0.0001$ .



**Figure 3.** Scatter plot showing the most serious outcome in all studies, and for studies within each stage. Diamonds shows the results of random effects meta-analysis.

### 3 ambavirumab COVID-19 studies


c19early.org  
June 2024



**Figure 4.** Random effects meta-analysis for all studies. This plot shows pooled effects, see the specific outcome analyses for individual outcomes. Analysis validating pooled outcomes for COVID-19 can be found below. Effect extraction is pre-specified, using the most serious outcome reported. For details see the appendix.

### 3 ambavirumab COVID-19 mortality results


c19early.org  
June 2024



**Figure 5.** Random effects meta-analysis for mortality results.

### 1 ambavirumab COVID-19 ICU result

c19early.org  
June 2024



**Figure 6.** Random effects meta-analysis for ICU admission.

## 2 ambavirumab COVID-19 hospitalization results

c19early.org  
June 2024




Figure 7. Random effects meta-analysis for hospitalization.

## 1 ambavirumab COVID-19 recovery result

c19early.org  
June 2024




Figure 8. Random effects meta-analysis for recovery.

## 1 ambavirumab COVID-19 viral clearance result

c19early.org  
June 2024





Figure 9. Random effects meta-analysis for viral clearance.

## Randomized Controlled Trials (RCTs)

Figure 10 shows a comparison of results for RCTs and non-RCT studies. Figure 11 shows a forest plot for random effects meta-analysis of all Randomized Controlled Trials. RCT results are included in Table 2 and Table 3.



**Figure 10.** Results for RCTs and non-RCT studies.

## 2 ambavirumab COVID-19 Randomized Controlled Trials



**Figure 11.** Random effects meta-analysis for all Randomized Controlled Trials. This plot shows pooled effects, see the specific outcome analyses for individual outcomes. Analysis validating pooled outcomes for COVID-19 can be found below. Effect extraction is pre-specified, using the most serious outcome reported. For details see the appendix.

RCTs have many potential biases. RCTs help to make study groups more similar and can provide a higher level of evidence, however they are subject to many biases<sup>20</sup>, and analysis of double-blind RCTs has identified extreme levels of bias<sup>21</sup>. For COVID-19, the overhead may delay treatment, dramatically compromising efficacy; they may encourage monotherapy for simplicity at the cost of efficacy which may rely on combined or synergistic effects; the participants that sign up may not reflect real world usage or the population that benefits most in terms of age, comorbidities, severity of illness, or other factors; standard of care may be compromised and unable to evolve quickly based on emerging research for new diseases; errors may be made in randomization and medication delivery; and investigators may have hidden agendas or vested interests influencing design, operation, analysis, reporting, and the potential for fraud. All of these biases have been observed with COVID-19 RCTs. There is no guarantee that a specific RCT provides a higher level of evidence.

**Conflicts of interest for COVID-19 RCTs.** RCTs are expensive and many RCTs are funded by pharmaceutical companies or interests closely aligned with pharmaceutical companies. For COVID-19, this creates an incentive to show efficacy for patented commercial products, and an incentive to show a lack of efficacy for inexpensive treatments. The bias is expected to be significant, for example *Als-Nielsen et al.* analyzed 370 RCTs from Cochrane reviews, showing that trials funded by for-profit organizations were 5 times more likely to recommend the experimental drug compared with those funded by nonprofit organizations. For COVID-19, some major philanthropic organizations are largely funded by investments with extreme conflicts of interest for and against specific COVID-19 interventions.

**RCTs for novel acute diseases requiring rapid treatment.** High quality RCTs for novel acute diseases are more challenging, with increased ethical issues due to the urgency of treatment, increased risk due to enrollment delays, and more difficult design with a rapidly evolving evidence base. For COVID-19, the most common site of initial infection is the upper respiratory tract. Immediate treatment is likely to be most successful and may prevent or slow progression to other parts of the body. For a non-prophylaxis RCT, it makes sense to provide treatment in advance and instruct patients to use it immediately on symptoms, just as some governments have done by providing medication

kits in advance. Unfortunately, no RCTs have been done in this way. Every treatment RCT to date involves delayed treatment. Among the 72 treatments we have analyzed, 64% of RCTs involve very late treatment 5+ days after onset. No non-prophylaxis COVID-19 RCTs match the potential real-world use of early treatments. They may more accurately represent results for treatments that require visiting a medical facility, e.g., those requiring intravenous administration.

**Non-RCT studies have been shown to be reliable.** Evidence shows that non-RCT studies can also provide reliable results. *Concato et al.* found that well-designed observational studies do not systematically overestimate the magnitude of the effects of treatment compared to RCTs. *Anglemyer et al.* summarized reviews comparing RCTs to observational studies and found little evidence for significant differences in effect estimates. *Lee et al.* showed that only 14% of the guidelines of the Infectious Diseases Society of America were based on RCTs. Evaluation of studies relies on an understanding of the study and potential biases. Limitations in an RCT can outweigh the benefits, for example excessive dosages, excessive treatment delays, or Internet survey bias may have a greater effect on results. Ethical issues may also prevent running RCTs for known effective treatments. For more on issues with RCTs see<sup>26,27</sup>.

**Using all studies identifies efficacy 7+ months faster (8+ months for low-cost treatments).** Currently, 45 of the treatments we analyze show statistically significant efficacy or harm, defined as  $\geq 10\%$  decreased risk or  $>0\%$  increased risk from  $\geq 3$  studies. Of these, 29 have been confirmed in RCTs, with a mean delay of 7.1 months. When considering only low cost treatments, 24 have been confirmed with a delay of 8.5 months. For the 16 unconfirmed treatments, 3 have zero RCTs to date. The point estimates for the remaining 13 are all consistent with the overall results (benefit or harm), with 10 showing  $>20\%$ . The only treatments showing  $>10\%$  efficacy for all studies, but  $<10\%$  for RCTs are sotrovimab and aspirin.

**Summary.** We need to evaluate each trial on its own merits. RCTs for a given medication and disease may be more reliable, however they may also be less reliable. For off-patent medications, very high conflict of interest trials may be more likely to be RCTs, and more likely to be large trials that dominate meta analyses.

## Exclusions


To avoid bias in the selection of studies, we analyze all non-retracted studies. Here we show the results after excluding studies with major issues likely to alter results, non-standard studies, and studies where very minimal detail is currently available. Our bias evaluation is based on analysis of each study and identifying when there is a significant chance that limitations will substantially change the outcome of the study. We believe this can be more valuable than checklist-based approaches such as Cochrane GRADE, which can be easily influenced by potential bias, may ignore or underemphasize serious issues not captured in the checklists, and may overemphasize issues unlikely to alter outcomes in specific cases (for example certain specifics of randomization with a very large effect size and well-matched baseline characteristics).

The studies excluded are as below. Figure 12 shows a forest plot for random effects meta-analysis of all studies after exclusions.

*Yalan*, unadjusted differences between groups.



## 2 ambavirumab COVID-19 studies after exclusions



**Figure 12.** Random effects meta-analysis for all studies after exclusions. This plot shows pooled effects, see the specific outcome analyses for individual outcomes. Analysis validating pooled outcomes for COVID-19 can be found below. Effect extraction is pre-specified, using the most serious outcome reported. For details see the appendix.

## Heterogeneity


Heterogeneity in COVID-19 studies arises from many factors including:

**Treatment delay.** The time between infection or the onset of symptoms and treatment may critically affect how well a treatment works. For example an antiviral may be very effective when used early but may not be effective in late stage disease, and may even be harmful. Oseltamivir, for example, is generally only considered effective for influenza when used within 0-36 or 0-48 hours<sup>29,30</sup>. Baloxavir studies for influenza also show that treatment delay is critical — *Ikematsu et al.* report an 86% reduction in cases for post-exposure prophylaxis, *Hayden et al.* show a 33 hour reduction in the time to alleviation of symptoms for treatment within 24 hours and a reduction of 13 hours for treatment within 24-48 hours, and *Kumar et al.* report only 2.5 hours improvement for inpatient treatment.


Treatment delay	Result
Post-exposure prophylaxis	86% fewer cases <sup>31</sup>
<24 hours	-33 hours symptoms <sup>32</sup>
24-48 hours	-13 hours symptoms <sup>32</sup>
Inpatients	-2.5 hours to improvement <sup>33</sup>

**Table 4.** Studies of baloxavir for influenza show that early treatment is more effective.

Figure 13 shows a mixed-effects meta-regression of efficacy as a function of treatment delay in COVID-19 ambavirumab studies, with group estimates for different stages when a specific value is not provided. For comparison, Figure 14 shows a meta-regression for all studies providing specific values across 72 treatments. Efficacy declines rapidly with treatment delay. Early treatment is critical for COVID-19.



**Figure 14.** Early treatment is more effective. Meta-regression showing efficacy as a function of treatment delay in COVID-19 ambavirumab studies.



**Figure 14.** Early treatment is more effective. Meta-regression showing efficacy as a function of treatment delay in COVID-19 studies from 72 treatments.

**Patient demographics.** Details of the patient population including age and comorbidities may critically affect how well a treatment works. For example, many COVID-19 studies with relatively young low-comorbidity patients show all patients recovering quickly with or without treatment. In such cases, there is little room for an effective treatment to improve results, for example as in *López-Medina et al.*

**Variants.** Efficacy may depend critically on the distribution of SARS-CoV-2 variants encountered by patients. Risk varies significantly across variants<sup>35</sup>, for example the Gamma variant shows significantly different characteristics<sup>36-39</sup>. Different mechanisms of action may be more or less effective depending on variants, for example the degree to which TMPRSS2 contributes to viral entry can differ across variants<sup>40,41</sup>.

**Regimen.** Effectiveness may depend strongly on the dosage and treatment regimen.

**Other treatments.** The use of other treatments may significantly affect outcomes, including supplements, other medications, or other interventions such as prone positioning. Treatments may be synergistic<sup>42-52</sup>, therefore efficacy may depend strongly on combined treatments.

**Medication quality.** The quality of medications may vary significantly between manufacturers and production batches, which may significantly affect efficacy and safety. *Williams et al.* analyze ivermectin from 11 different sources, showing highly variable antiparasitic efficacy across different manufacturers. *Xu et al.* analyze a treatment from two different manufacturers, showing 9 different impurities, with significantly different concentrations for each manufacturer.

**Effect measured.** Across all studies there is a strong association between different outcomes, for example improved recovery is strongly associated with lower mortality. However, efficacy may differ depending on the effect measured, for example a treatment may be more effective against secondary complications and have minimal effect on viral clearance.

**Meta analysis.** The distribution of studies will alter the outcome of a meta analysis. Consider a simplified example where everything is equal except for the treatment delay, and effectiveness decreases to zero or below with increasing delay. If there are many studies using very late treatment, the outcome may be negative, even though early treatment is very effective. All meta analyses combine heterogeneous studies, varying in population, variants, and potentially all factors above, and therefore may obscure efficacy by including studies where treatment is less effective. Generally, we expect the estimated effect size from meta analysis to be less than that for the optimal case. Looking at all studies is valuable for providing an overview of all research, important to avoid cherry-picking, and informative when a positive result is found despite combining less-optimal situations. However, the resulting estimate does not apply to specific cases such as early treatment in high-risk populations. While we present results for all studies, we also present treatment time and individual outcome analyses, which may be more informative for specific use cases.

## Pooled Effects

**Combining studies is required.** For COVID-19, delay in clinical results translates into additional death and morbidity, as well as additional economic and societal damage. Combining the results of studies reporting different outcomes is required. There may be no mortality in a trial with low-risk patients, however a reduction in severity or improved viral clearance may translate into lower mortality in a high-risk population. Different studies may report lower severity, improved recovery, and lower mortality, and the significance may be very high when combining the results. "*The studies reported different outcomes*" is not a good reason for disregarding results.


**Specific outcome and pooled analyses.** We present both specific outcome and pooled analyses. In order to combine the results of studies reporting different outcomes we use the most serious outcome reported in each study, based on the thesis that improvement in the most serious outcome provides comparable measures of efficacy for a treatment. A critical advantage of this approach is simplicity and transparency. There are many other ways to combine evidence for different outcomes, along with additional evidence such as dose-response relationships, however these increase complexity.

**Using more information.** Another way to view pooled analysis is that we are using more of the available information. Logically we should, and do, use additional information. For example dose-response and treatment delay-response relationships provide significant additional evidence of efficacy that is considered when reviewing the evidence for a treatment.


**Ethical and practical issues limit high-risk trials.** Trials with high-risk patients may be restricted due to ethics for treatments that are known or expected to be effective, and they increase difficulty for recruiting. Using less severe outcomes as a proxy for more serious outcomes allows faster collection of evidence.

**Improvement across outcomes.** For many COVID-19 treatments, a reduction in mortality logically follows from a reduction in hospitalization, which follows from a reduction in symptomatic cases, which follows from a reduction in PCR positivity. We can directly test this for COVID-19.


Validating pooled outcome analysis for COVID-19. Analysis of the the association between different outcomes across studies from all 72 treatments we cover confirms the validity of pooled outcome analysis for COVID-19. Figure 15 shows that lower hospitalization is very strongly associated with lower mortality ( $p < 0.00000000001$ ). Similarly, Figure 16 shows that improved recovery is very strongly associated with lower mortality ( $p < 0.00000000001$ ). Considering the extremes, *Singh et al.* show an association between viral clearance and hospitalization or death, with  $p = 0.003$  after excluding one large outlier from a mutagenic treatment, and based on 44 RCTs including 52,384 patients. Figure 17 shows that improved viral clearance is strongly associated with fewer serious outcomes. The association is very similar to *Singh et al.*, with higher confidence due to the larger number of studies. As with *Singh et al.*, the confidence increases when excluding the outlier treatment, from  $p = 0.0000014$  to  $p = 0.000000005$ .



**Figure 15.** Lower hospitalization is associated with lower mortality, supporting pooled outcome analysis.



**Figure 16.** Improved recovery is associated with lower mortality, supporting pooled outcome analysis.



**Figure 15.** Improved viral clearance is associated with fewer serious outcomes, supporting pooled outcome analysis.

Pooled outcomes identify efficacy 5 months faster (6 months for RCTs). Currently, 45 of the treatments we analyze show statistically significant efficacy or harm, defined as  $\geq 10\%$  decreased risk or  $>0\%$  increased risk from  $\geq 3$  studies. 90% of these have been confirmed with one or more specific outcomes, with a mean delay of 5.0 months. When restricting to RCTs only, 56% of treatments showing statistically significant efficacy/harm with pooled effects have been confirmed with one or more specific outcomes, with a mean delay of 6.4 months. Figure 18 shows when treatments were found effective during the pandemic. Pooled outcomes often resulted in earlier detection of efficacy.

## Time when COVID-19 studies showed efficacy

c19early.org  
June 2024



**Figure 18.** The time when studies showed that treatments were effective, defined as statistically significant improvement of  $\geq 10\%$  from  $\geq 3$  studies. Pooled results typically show efficacy earlier than specific outcome results. Results from all studies often shows efficacy much earlier than when restricting to RCTs. Results reflect conditions as used in trials to date, these depend on the population treated, treatment delay, and treatment regimen.

**Limitations.** Pooled analysis could hide efficacy, for example a treatment that is beneficial for late stage patients but has no effect on viral clearance may show no efficacy if most studies only examine viral clearance. In practice, it is rare for a non-antiviral treatment to report viral clearance and to not report clinical outcomes; and in practice other sources of heterogeneity such as difference in treatment delay is more likely to hide efficacy.

**Summary.** Analysis validates the use of pooled effects and shows significantly faster detection of efficacy on average. However, as with all meta analyses, it is important to review the different studies included. We also present individual outcome analyses, which may be more informative for specific use cases.

## Discussion

**Publication bias.** Publishing is often biased towards positive results. Trials with patented drugs may have a financial conflict of interest that results in positive studies being more likely to be published, or bias towards more positive results. For example with molnupiravir, trials with negative results remain unpublished to date (CTRI/2021/05/033864 and CTRI/2021/08/0354242). For ambavirumab, there is currently not enough data to evaluate publication bias with high confidence.

**Limitations.** Summary statistics from meta analysis necessarily lose information. As with all meta analyses, studies are heterogeneous, with differences in treatment delay, treatment regimen, patient demographics, variants, conflicts of interest, standard of care, and other factors. We provide analyses for specific outcomes and by treatment delay, and we aim to identify key characteristics in the forest plots and summaries. Results should be viewed in the context of study characteristics.

Some analyses classify treatment based on early or late administration, as done here, while others distinguish between mild, moderate, and severe cases. Viral load does not indicate degree of symptoms — for example patients may have a high viral load while being asymptomatic. With regard to treatments that have antiviral properties, timing of treatment is critical — late administration may be less helpful regardless of severity.

Details of treatment delay per patient is often not available. For example, a study may treat 90% of patients relatively early, but the events driving the outcome may come from 10% of patients treated very late. Our 5 day cutoff for early treatment may be too conservative, 5 days may be too late in many cases.

Comparison across treatments is confounded by differences in the studies performed, for example dose, variants, and conflicts of interest. Trials with conflicts of interest may use designs better suited to the preferred outcome.

In some cases, the most serious outcome has very few events, resulting in lower confidence results being used in pooled analysis, however the method is simpler and more transparent. This is less critical as the number of studies increases. Restriction to outcomes with sufficient power may be beneficial in pooled analysis and improve accuracy when there are few studies, however we maintain our pre-specified method to avoid any retrospective changes.

Studies show that combinations of treatments can be highly synergistic and may result in many times greater efficacy than individual treatments alone<sup>42-52</sup>. Therefore standard of care may be critical and benefits may diminish or disappear if standard of care does not include certain treatments.

This real-time analysis is constantly updated based on submissions. Accuracy benefits from widespread review and submission of updates and corrections from reviewers. Less popular treatments may receive fewer reviews.

No treatment or intervention is 100% available and effective for all current and future variants. Efficacy may vary significantly with different variants and within different populations. All treatments have potential side effects. Propensity to experience side effects may be predicted in advance by qualified physicians. We do not provide medical advice. Before taking any medication, consult a qualified physician who can compare all options, provide personalized advice, and provide details of risks and benefits based on individual medical history and situations.

## Perspective

**Results compared with other treatments.** SARS-CoV-2 infection and replication involves a complex interplay of 50+ host and viral proteins and other factors<sup>12-16</sup>, providing many therapeutic targets. Over 7,000 compounds have been predicted to reduce COVID-19 risk<sup>17</sup>, either by directly minimizing infection or replication, by supporting immune system function, or by minimizing secondary complications. Figure 19 shows an overview of the results for ambavirumab in the context of multiple COVID-19 treatments, and Figure 20 shows a plot of efficacy vs. cost for COVID-19 treatments.

# Efficacy in COVID-19 studies (pooled effects)

c19early.org

June 2024




Figure 19. Scatter plot showing results within the context of multiple COVID-19 treatments. Diamonds shows the results of random effects meta-analysis. 0.6% of 7,000+ proposed treatments show efficacy<sup>56</sup>.

# Efficacy vs. cost for COVID-19 treatments

c19early.org

June 2024




Figure 20. Efficacy vs. cost for COVID-19 treatments.



## Conclusion


Statistically significant lower risk is seen for viral clearance. 2 studies from 2 independent teams (both from the same country) show significant improvements. Meta analysis using the most serious outcome reported shows 11% [-154-69%] lower risk, without reaching statistical significance. Results are better for Randomized Controlled Trials and higher quality studies. Early treatment shows efficacy while late treatment does not, consistent with expectations for an antiviral treatment.

Currently there is limited data, with only 29 control events for the most serious outcome in trials to date. Studies to date are from only 3 different groups.

**Efficacy is variant dependent.** mAb use may create new variants that spread globally<sup>1,2</sup>, and may be associated with prolonged viral loads, clinical deterioration, and immune escape<sup>2-4</sup>.


## Study Notes

### Evering




*Evering*: RCT 837 non-hospitalized high-risk COVID-19 patients showing 78% reduction in hospitalization and death with BR11-196/BR11-198 (monoclonal antibodies). Study was terminated early. BR11-196/BR11-198 had fewer grade 3+ adverse events (3.8% vs 13.4%).

### Self



Self: RCT with 182 sotrovimab patients, 176 BR11-196+BR11-198 patients, and 178 control patients, median 8 days from symptom onset, showing no significant differences and terminated early due to futility.

## Yalan



Yalan: Retrospective 340 COVID-19 patients in China showing shorter length of hospital stay and faster viral clearance with BR11-196 plus BR11-198 monoclonal antibody treatment, especially when given early. The treatment did not show efficacy for improving clinical outcomes among severe or critical cases.

## Appendix 1. Methods and Data

We perform ongoing searches of PubMed, medRxiv, Europe PMC, ClinicalTrials.gov, The Cochrane Library, Google Scholar, Research Square, ScienceDirect, Oxford University Press, the reference lists of other studies and meta-analyses, and submissions to the site c19early.org. Search terms are ambavirumab and COVID-19 or SARS-CoV-2. Automated searches are performed twice daily, with all matches reviewed for inclusion. All studies regarding the use of ambavirumab for COVID-19 that report a comparison with a control group are included in the main analysis. Sensitivity analysis is performed, excluding studies with major issues, epidemiological studies, and studies with minimal available information. This is a living analysis and is updated regularly.

We extracted effect sizes and associated data from all studies. If studies report multiple kinds of effects then the most serious outcome is used in pooled analysis, while other outcomes are included in the outcome specific analyses. For example, if effects for mortality and cases are both reported, the effect for mortality is used, this may be different to the effect that a study focused on. If symptomatic results are reported at multiple times, we used the latest time, for example if mortality results are provided at 14 days and 28 days, the results at 28 days have preference. Mortality alone is preferred over combined outcomes. Outcomes with zero events in both arms are not used, the next most serious outcome with one or more events is used. For example, in low-risk populations with no mortality, a reduction in mortality with treatment is not possible, however a reduction in hospitalization, for example, is still valuable. Clinical outcomes are considered more important than viral test status. When basically all patients recover in both treatment and control groups, preference for viral clearance and recovery is given to results mid-recovery where available. After most or all patients have recovered there is little or no room for an effective treatment to do better, however faster recovery is valuable. If only individual symptom data is available, the most serious symptom has priority, for example difficulty breathing or low SpO<sub>2</sub> is more important than cough. When results provide an odds ratio, we compute the relative risk when possible, or convert to a relative risk according to<sup>59</sup>. Reported confidence intervals and  $p$ -values were used when available, using adjusted values when provided. If multiple types of adjustments are reported propensity score matching and multivariable regression has preference over propensity score matching or weighting, which has preference over multivariable regression. Adjusted results have preference over unadjusted results for a more serious outcome when the adjustments significantly alter results. When needed, conversion between reported  $p$ -values and confidence intervals followed *Altman, Altman (B)*, and Fisher's exact test was used to calculate  $p$ -values for event data. If continuity correction for zero values is required, we use the reciprocal of the opposite arm with the

sum of the correction factors equal to 1<sup>62</sup>. Results are expressed with RR < 1.0 favoring treatment, and using the risk of a negative outcome when applicable (for example, the risk of death rather than the risk of survival). If studies only report relative continuous values such as relative times, the ratio of the time for the treatment group versus the time for the control group is used. Calculations are done in Python (3.12.3) with scipy (1.13.1), pythonmeta (1.26), numpy (1.26.4), statsmodels (0.14.2), and plotly (5.22.0).

Forest plots are computed using PythonMeta<sup>63</sup> with the DerSimonian and Laird random effects model (the fixed effect assumption is not plausible in this case) and inverse variance weighting. Results are presented with 95% confidence intervals. Heterogeneity among studies was assessed using the I<sup>2</sup> statistic. Mixed-effects meta-regression results are computed with R (4.4.0) using the metafor (4.6-0) and rms (6.8-0) packages, and using the most serious sufficiently powered outcome. For all statistical tests, a *p*-value less than 0.05 was considered statistically significant. Grobid 0.8.0 is used to parse PDF documents.

We have classified studies as early treatment if most patients are not already at a severe stage at the time of treatment (for example based on oxygen status or lung involvement), and treatment started within 5 days of the onset of symptoms. If studies contain a mix of early treatment and late treatment patients, we consider the treatment time of patients contributing most to the events (for example, consider a study where most patients are treated early but late treatment patients are included, and all mortality events were observed with late treatment patients). We note that a shorter time may be preferable. Antivirals are typically only considered effective when used within a shorter timeframe, for example 0-36 or 0-48 hours for oseltamivir, with longer delays not being effective<sup>29,30</sup>.

We received no funding, this research is done in our spare time. We have no affiliations with any pharmaceutical companies or political parties.

A summary of study results is below. Please submit updates and corrections at <https://c19early.org/ammeta.html>.

## Early treatment

Effect extraction follows pre-specified rules as detailed above and gives priority to more serious outcomes. For pooled analyses, the first (most serious) outcome is used, which may differ from the effect a paper focuses on. Other outcomes are used in outcome specific analyses.

<i>Evering</i> , 11/1/2021, Double Blind Randomized Controlled Trial, placebo-controlled, multiple countries, peer-reviewed, median age 49.0, 18 authors, study period January 2021 - July 2021, trial NCT03383419 (history).	risk of death, 88.9% lower, RR 0.11, <i>p</i> = 0.02, treatment 1 of 418 (0.2%), control 9 of 419 (2.1%), NNT 52.
	risk of hospitalization, 73.3% lower, RR 0.27, <i>p</i> < 0.001, treatment 12 of 418 (2.9%), control 45 of 419 (10.7%), NNT 13.
	risk of death/hospitalization, 78.7% lower, RR 0.21, <i>p</i> < 0.001, treatment 10 of 418 (2.4%), control 47 of 419 (11.2%), NNT 11.

## Late treatment

Effect extraction follows pre-specified rules as detailed above and gives priority to more serious outcomes. For pooled analyses, the first (most serious) outcome is used, which may differ from the effect a paper focuses on. Other outcomes are used in outcome specific analyses.

<i>Self</i> , 12/23/2021, Double Blind Randomized Controlled Trial, multiple countries, peer-reviewed, 67 authors, study period 16 December, 2020 - 1 March, 2021, average treatment delay 8.0 days, trial NCT04501978 (history) (TICO).	risk of death, 15.0% higher, RR 1.15, <i>p</i> = 0.72, treatment 15 of 176 (8.5%), control 13 of 178 (7.3%), adjusted per study, day 90.
	risk of no recovery, 7.4% lower, RR 0.93, <i>p</i> = 0.48, treatment 21 of 176 (11.9%), control 27 of 178 (15.2%), adjusted per study, inverted to make RR<1 favor treatment, day 90, primary

	outcome.
	risk of no recovery, no change, RR 1.00, $p = 0.99$ , treatment 173, control 178, adjusted per study, inverted to make $RR < 1$ favor treatment, pulmonary-plus ordinal outcome @day 5, day 5.
Yalan, 4/19/2024, retrospective, China, peer-reviewed, median age 72.0, 6 authors, study period October 2022 - November 2022, excluded in exclusion analyses: unadjusted differences between groups.	risk of death, 71.4% higher, RR 1.71, $p = 0.35$ , treatment 12 of 170 (7.1%), control 7 of 170 (4.1%).
	risk of ICU admission, 7.7% higher, RR 1.08, $p = 0.80$ , treatment 42 of 170 (24.7%), control 39 of 170 (22.9%).
	hospitalization time, 7.7% lower, relative time 0.92, $p = 0.004$ , treatment 170, control 170.
	time to viral-, 6.7% lower, relative time 0.93, $p = 0.004$ , treatment 170, control 170.

## Supplementary Data

Supplementary Data

## Footnotes

- a. Viral infection and replication involves attachment, entry, uncoating and release, genome replication and transcription, translation and protein processing, assembly and budding, and release. Each step can be disrupted by therapeutics.

## References

1. **Focosi** et al., *Analysis of SARS-CoV-2 mutations associated with resistance to therapeutic monoclonal antibodies that emerge after treatment*, Drug Resistance Updates, doi:10.1016/j.drug.2023.100991.
2. **Leducq** et al., *Spike protein genetic evolution in patients at high-risk of severe COVID-19 treated by monoclonal antibodies*, The Journal of Infectious Diseases, doi:10.1093/infdis/jiad523.
3. **Choudhary** et al., *Emergence of SARS-CoV-2 Resistance with Monoclonal Antibody Therapy*, medRxiv, doi:10.1101/2021.09.03.21263105.
4. **Günther** et al., *Variant-specific humoral immune response to SARS-CoV-2 escape mutants arising in clinically severe, prolonged infection*, medRxiv, doi:10.1101/2024.01.06.24300890.
5. **Yang** et al., *SARS-CoV-2 infection causes dopaminergic neuron senescence*, Cell Stem Cell, doi:10.1016/j.stem.2023.12.012.
6. **Scardua-Silva** et al., *Microstructural brain abnormalities, fatigue, and cognitive dysfunction after mild COVID-19*, Scientific Reports, doi:10.1038/s41598-024-52005-7.
7. **Hampshire** et al., *Cognition and Memory after Covid-19 in a Large Community Sample*, New England Journal of Medicine, doi:10.1056/NEJMoa2311330.
8. **Duloquin** et al., *Is COVID-19 Infection a Multiorgan Disease? Focus on Extrapulmonary Involvement of SARS-CoV-2*, Journal of Clinical Medicine, doi:10.3390/jcm13051397.

9. **Sodagar** et al., *Pathological Features and Neuroinflammatory Mechanisms of SARS-CoV-2 in the Brain and Potential Therapeutic Approaches*, *Biomolecules*, doi:10.3390/biom12070971.
10. **Sagar** et al., *COVID-19-associated cerebral microbleeds in the general population*, *Brain Communications*, doi:10.1093/braincomms/fcae127.
11. **Eberhardt** et al., *SARS-CoV-2 infection triggers pro-atherogenic inflammatory responses in human coronary vessels*, *Nature Cardiovascular Research*, doi:10.1038/s44161-023-00336-5.
12. **Malone** et al., *Structures and functions of coronavirus replication–transcription complexes and their relevance for SARS-CoV-2 drug design*, *Nature Reviews Molecular Cell Biology*, doi:10.1038/s41580-021-00432-z.
13. **Murigneux** et al., *Proteomic analysis of SARS-CoV-2 particles unveils a key role of G3BP proteins in viral assembly*, *Nature Communications*, doi:10.1038/s41467-024-44958-0.
14. **Lv** et al., *Host proviral and antiviral factors for SARS-CoV-2*, *Virus Genes*, doi:10.1007/s11262-021-01869-2.
15. **Lui** et al., *Nsp1 facilitates SARS-CoV-2 replication through calcineurin-NFAT signaling*, *Virology*, doi:10.1128/mbio.00392-24.
16. **Niarakis** et al., *Drug-target identification in COVID-19 disease mechanisms using computational systems biology approaches*, *Frontiers in Immunology*, doi:10.3389/fimmu.2023.1282859.
17. **c19early.org**, [c19early.org/treatments.html](http://c19early.org/treatments.html).
18. **Hattab** et al., *SARS-CoV-2 journey: from alpha variant to omicron and its sub-variants*, *Infection*, doi:10.1007/s15010-024-02223-y.
19. **Davis** et al., *The Promise and Peril of Anti-SARS-CoV-2 Monoclonal Antibodies*, *Clinical Infectious Diseases*, doi:10.1093/cid/ciac902.
20. **Jadad** et al., *Randomized Controlled Trials: Questions, Answers, and Musings, Second Edition*, doi:10.1002/9780470691922.
21. **Göttsche**, P., *Bias in double-blind trials*, Doctoral Thesis, University of Copenhagen, [www.scientificfreedom.dk/2023/05/16/bias-in-double-blind-trials-doctoral-thesis/](http://www.scientificfreedom.dk/2023/05/16/bias-in-double-blind-trials-doctoral-thesis/).
22. **Als-Nielsen** et al., *Association of Funding and Conclusions in Randomized Drug Trials*, *JAMA*, doi:10.1001/jama.290.7.921.
23. **Concato** et al., *NEJM*, 342:1887-1892, doi:10.1056/NEJM200006223422507.
24. **Anglemyer** et al., *Healthcare outcomes assessed with observational study designs compared with those assessed in randomized trials*, *Cochrane Database of Systematic Reviews* 2014, Issue 4, doi:10.1002/14651858.MR000034.pub2.
25. **Lee** et al., *Analysis of Overall Level of Evidence Behind Infectious Diseases Society of America Practice Guidelines*, *Arch Intern Med.*, 2011, 171:1, 18-22, doi:10.1001/archinternmed.2010.482.
26. **Deaton** et al., *Understanding and misunderstanding randomized controlled trials*, *Social Science & Medicine*, 210, doi:10.1016/j.socscimed.2017.12.005.
27. **Nichol** et al., *Challenging issues in randomised controlled trials*, *Injury*, 2010, doi: 10.1016/j.injury.2010.03.033, [www.injuryjournal.com/article/S0020-1383\(10\)00233-0/fulltext](http://www.injuryjournal.com/article/S0020-1383(10)00233-0/fulltext).
28. **Yalan** et al., *Treatment for Covid-19 with SARS-CoV-2 neutralizing antibody BR11-196(Ambavirumab) plus BR11-198(Lomisivir): a retrospective cohort study*, *BMC Pharmacology and Toxicology*, doi:10.1186/s40360-024-00753-7.
29. **Treanor** et al., *Efficacy and Safety of the Oral Neuraminidase Inhibitor Oseltamivir in Treating Acute Influenza: A Randomized Controlled Trial*, *JAMA*, 2000, 283:8, 1016-1024, doi:10.1001/jama.283.8.1016.
30. **McLean** et al., *Impact of Late Oseltamivir Treatment on Influenza Symptoms in the Outpatient Setting: Results of a Randomized Trial*, *Open Forum Infect. Dis.* September 2015, 2:3, doi:10.1093/ofid/ofv100.
31. **Ikematsu** et al., *Baloxavir Marboxil for Prophylaxis against Influenza in Household Contacts*, *New England Journal of Medicine*, doi:10.1056/NEJMoa1915341.
32. **Hayden** et al., *Baloxavir Marboxil for Uncomplicated Influenza in Adults and Adolescents*, *New England Journal of Medicine*, doi:10.1056/NEJMoa1716197.

33. **Kumar** et al., *Combining baloxavir marboxil with standard-of-care neuraminidase inhibitor in patients hospitalised with severe influenza (FLAGSTONE): a randomised, parallel-group, double-blind, placebo-controlled, superiority trial*, *The Lancet Infectious Diseases*, doi:10.1016/S1473-3099(21)00469-2.
34. **López-Medina** et al., *Effect of Ivermectin on Time to Resolution of Symptoms Among Adults With Mild COVID-19: A Randomized Clinical Trial*, *JAMA*, doi:10.1001/jama.2021.3071.
35. **Korves** et al., *SARS-CoV-2 Genetic Variants and Patient Factors Associated with Hospitalization Risk*, medRxiv, doi:10.1101/2024.03.08.24303818.
36. **Faria** et al., *Genomics and epidemiology of the P.1 SARS-CoV-2 lineage in Manaus, Brazil*, *Science*, doi:10.1126/science.abh2644.
37. **Nonaka** et al., *SARS-CoV-2 variant of concern P.1 (Gamma) infection in young and middle-aged patients admitted to the intensive care units of a single hospital in Salvador, Northeast Brazil, February 2021*, *International Journal of Infectious Diseases*, doi:10.1016/j.ijid.2021.08.003.
38. **Karita** et al., *Trajectory of viral load in a prospective population-based cohort with incident SARS-CoV-2 G614 infection*, medRxiv, doi:10.1101/2021.08.27.21262754.
39. **Zavascki** et al., *Advanced ventilatory support and mortality in hospitalized patients with COVID-19 caused by Gamma (P.1) variant of concern compared to other lineages: cohort study at a reference center in Brazil*, *Research Square*, doi:10.21203/rs.3.rs-910467/v1.
40. **Willett** et al., *The hyper-transmissible SARS-CoV-2 Omicron variant exhibits significant antigenic change, vaccine escape and a switch in cell entry mechanism*, medRxiv, doi:10.1101/2022.01.03.21268111.
41. **Peacock** et al., *The SARS-CoV-2 variant, Omicron, shows rapid replication in human primary nasal epithelial cultures and efficiently uses the endosomal route of entry*, bioRxiv, doi:10.1101/2021.12.31.474653.
42. **Jitobaom** et al., *Favipiravir and Ivermectin Showed in Vitro Synergistic Antiviral Activity against SARS-CoV-2*, *Research Square*, doi:10.21203/rs.3.rs-941811/v1.
43. **Jitobaom (B)** et al., *Synergistic anti-SARS-CoV-2 activity of repurposed anti-parasitic drug combinations*, *BMC Pharmacology and Toxicology*, doi:10.1186/s40360-022-00580-8.
44. **Jeffreys** et al., *Remdesivir-ivermectin combination displays synergistic interaction with improved in vitro activity against SARS-CoV-2*, *International Journal of Antimicrobial Agents*, doi:10.1016/j.ijantimicag.2022.106542.
45. **Ostrov** et al., *Highly Specific Sigma Receptor Ligands Exhibit Anti-Viral Properties in SARS-CoV-2 Infected Cells*, *Pathogens*, doi:10.3390/pathogens10111514.
46. **Alsaïdi** et al., *Griffithsin and Carrageenan Combination Results in Antiviral Synergy against SARS-CoV-1 and 2 in a Pseudoviral Model*, *Marine Drugs*, doi:10.3390/md19080418.
47. **Andreani** et al., *In vitro testing of combined hydroxychloroquine and azithromycin on SARS-CoV-2 shows synergistic effect*, *Microbial Pathogenesis*, doi:10.1016/j.micpath.2020.104228.
48. **De Forni** et al., *Synergistic drug combinations designed to fully suppress SARS-CoV-2 in the lung of COVID-19 patients*, *PLoS ONE*, doi:10.1371/journal.pone.0276751.
49. **Wan** et al., *Synergistic inhibition effects of andrographolide and baicalin on coronavirus mechanisms by downregulation of ACE2 protein level*, *Scientific Reports*, doi:10.1038/s41598-024-54722-5.
50. **Said** et al., *The effect of Nigella sativa and vitamin D3 supplementation on the clinical outcome in COVID-19 patients: A randomized controlled clinical trial*, *Frontiers in Pharmacology*, doi:10.3389/fphar.2022.1011522.
51. **Fiaschi** et al., *In Vitro Combinatorial Activity of Direct Acting Antivirals and Monoclonal Antibodies against the Ancestral B.1 and BQ.1.1 SARS-CoV-2 Viral Variants*, *Viruses*, doi:10.3390/v16020168.
52. **Thairu** et al., *A Comparison of Ivermectin and Non Ivermectin Based Regimen for COVID-19 in Abuja: Effects on Virus Clearance, Days-to-discharge and Mortality*, *Journal of Pharmaceutical Research International*, doi:10.9734/jpri/2022/v34i44A36328.



53. **Williams, T.**, *Not All Ivermectin Is Created Equal: Comparing The Quality of 11 Different Ivermectin Sources*, Do Your Own Research, [doyourownresearch.substack.com/p/not-all-ivermectin-is-created-equal](https://doyourownresearch.substack.com/p/not-all-ivermectin-is-created-equal).
54. **Xu et al.**, *A study of impurities in the repurposed COVID-19 drug hydroxychloroquine sulfate by UHPLC-Q/TOF-MS and LC-SPE-NMR*, Rapid Communications in Mass Spectrometry, doi:10.1002/rcm.9358.
55. **Singh et al.**, *The relationship between viral clearance rates and disease progression in early symptomatic COVID-19: a systematic review and meta-regression analysis*, Journal of Antimicrobial Chemotherapy, doi:10.1093/jac/dkac045.
56. **c19early.org (B)**, [c19early.org/timeline.html](https://c19early.org/timeline.html).
57. **Evering et al.**, *Safety and Efficacy of Combination SARS-CoV-2 Monoclonal Neutralizing Antibodies (mAb) BRIL-196 and BRIL-198 in Non-Hospitalized COVID-19 Patients*, Open Forum Infectious Diseases, doi:10.1093/ofid/ofab466.1643.
58. **Self et al.**, *Efficacy and safety of two neutralising monoclonal antibody therapies, sotrovimab and BRIL-196 plus BRIL-198, for adults hospitalised with COVID-19 (TICO): a randomised controlled trial*, The Lancet Infectious Diseases, doi:10.1016/S1473-3099(21)00751-9.
59. **Zhang et al.**, *What's the relative risk? A method of correcting the odds ratio in cohort studies of common outcomes*, JAMA, 80:19, 1690, doi:10.1001/jama.280.19.1690.
60. **Altman, D.**, *How to obtain the P value from a confidence interval*, BMJ, doi:10.1136/bmj.d2304.
61. **Altman (B) et al.**, *How to obtain the confidence interval from a P value*, BMJ, doi:10.1136/bmj.d2090.
62. **Sweeting et al.**, *What to add to nothing? Use and avoidance of continuity corrections in meta-analysis of sparse data*, Statistics in Medicine, doi:10.1002/sim.1761.
63. **Deng, H.**, *PyMeta, Python module for meta-analysis*, [www.pymeta.com/](http://www.pymeta.com/).