

Azelastine for COVID-19: real-time meta analysis of 3 studies

@CovidAnalysis, March 2025, Version 5

<https://c19early.org/azlmeta.html>

Abstract

Significantly lower risk is seen for cases and viral clearance. 3 studies from 3 independent teams in 3 countries show significant benefit.


Meta analysis using the most serious outcome reported shows 21% [-3-39%] lower risk, without reaching statistical significance. Results are worse for Randomized Controlled Trials.

Currently there is limited data, with only 310 patients in trials to date. Studies to date are from only 3 different groups.

No treatment is 100% effective. Protocols combine safe and effective options with individual risk/benefit analysis and monitoring. Other treatments are more effective. All data and sources to reproduce this analysis are in the appendix.


Azelastine for COVID-19

	Improvement	Studies	Patients
All studies	21%	3	310
Recovery	13%	2	310
Cases	45%	1	0
Viral clearance	7%	2	310
RCTs	13%	2	310
Prophylaxis	45%	1	0
Early	13%	1	59
Late	12%	1	251



Evolution of COVID-19 clinical evidence

Meta analysis results over time




AZELASTINE FOR COVID-19 – HIGHLIGHTS

Azelastine reduces risk with low confidence for cases, viral clearance, and in pooled analysis.

Real-time updates and corrections with a consistent protocol for 116 treatments. Outcome specific analysis and combined evidence from all studies including treatment delay, a primary confounding factor.

3 azelastine COVID-19 studies



Timeline of COVID-19 azelastine studies (pooled effects)




Figure 1. A. Random effects meta-analysis. This plot shows pooled effects, see the specific outcome analyses for individual outcomes. Analysis validating pooled outcomes for COVID-19 can be found below. Effect extraction is pre-specified, using the most serious outcome reported. For details see the appendix. **B. Timeline of results in azelastine studies.**

Introduction

Immediate treatment recommended

SARS-CoV-2 infection primarily begins in the upper respiratory tract and may progress to the lower respiratory tract, other tissues, and the nervous and cardiovascular systems, which may lead to cytokine storm, pneumonia, ARDS, neurological injury¹⁻¹³ and cognitive deficits^{4,9}, cardiovascular complications¹⁴⁻¹⁷, organ failure, and death. Minimizing replication as early as possible is recommended.

Many treatments are expected to modulate infection

SARS-CoV-2 infection and replication involves the complex interplay of 50+ host and viral proteins and other factors^{A,18-24}, providing many therapeutic targets for which many existing compounds have known activity. Scientists have predicted that over 8,000 compounds may reduce COVID-19 risk²⁵, either by directly minimizing infection or replication, by supporting immune system function, or by minimizing secondary complications.

Analysis

We analyze all significant controlled studies of azelastine for COVID-19. Search methods, inclusion criteria, effect extraction criteria (more serious outcomes have priority), all individual study data, PRISMA answers, and statistical methods are detailed in Appendix 1. We present random effects meta-analysis results for all studies, studies within each treatment stage, individual outcomes, and Randomized Controlled Trials (RCTs).

Treatment timing

Figure 2 shows stages of possible treatment for COVID-19. Prophylaxis refers to regularly taking medication before becoming sick, in order to prevent or minimize infection. Early Treatment refers to treatment immediately or soon after symptoms appear, while Late Treatment refers to more delayed treatment.




Figure 2. Treatment stages.

Preclinical Research

An *In Silico* study supports the efficacy of azelastine²⁶.

An *In Vitro* study supports the efficacy of azelastine²⁷.

Preclinical research is an important part of the development of treatments, however results may be very different in clinical trials. Preclinical results are not used in this paper.

Results

Table 1 summarizes the results for all stages combined, for Randomized Controlled Trials, and for specific outcomes. Table 2 shows results by treatment stage. Figure 3 plots individual results by treatment stage. Figure 4, 5, 6, and 7 show forest plots for random effects meta-analysis of all studies with pooled effects, recovery, cases, and viral clearance.

	Improvement	Studies	Patients	Authors
All studies	21% [-3-39%]	3	310	44
Randomized Controlled Trials	13% [-13-32%]	2	310	35
Recovery	13% [-13-32%]	2	310	35
Viral	7% [4-9%] ****	2	310	35

Table 1. Random effects meta-analysis for all stages combined, for Randomized Controlled Trials, and for specific outcomes. Results show the percentage improvement with treatment and the 95% confidence interval. * $p < 0.05$
**** $p < 0.0001$.

	Early treatment	Late treatment	Prophylaxis
All studies	13% [-33-43%]	12% [-20-36%]	45% [6-68%] *
Randomized Controlled Trials	13% [-33-43%]	12% [-20-36%]	
Recovery	13% [-33-43%]	12% [-20-36%]	
Viral	14% [-9-33%]	6% [4-9%] ****	

Table 2. Random effects meta-analysis results by treatment stage. Results show the percentage improvement with treatment, the 95% confidence interval, and the number of studies for the stage. * $p < 0.05$ **** $p < 0.0001$.




Figure 3. Scatter plot showing the most serious outcome in all studies, and for studies within each stage. Diamonds shows the results of random effects meta-analysis.

3 azelastine COVID-19 studies




Figure 4. Random effects meta-analysis for all studies. This plot shows pooled effects, see the specific outcome analyses for individual outcomes. Analysis validating pooled outcomes for COVID-19 can be found below. Effect extraction is pre-specified, using the most serious outcome reported. For details see the appendix.

2 azelastine COVID-19 recovery results

c19early.org
March 2025




Figure 5. Random effects meta-analysis for recovery.

1 azelastine COVID-19 case result

c19early.org
March 2025



Figure 6. Random effects meta-analysis for cases.

2 azelastine COVID-19 viral clearance results

c19early.org
March 2025




Figure 7. Random effects meta-analysis for viral clearance.

Randomized Controlled Trials (RCTs)

Figure 8 shows a comparison of results for RCTs and non-RCT studies. Figure 9 shows a forest plot for random effects meta-analysis of all Randomized Controlled Trials. RCT results are included in Table 1 and Table 2.




Figure 8. Results for RCTs and non-RCT studies.

2 azelastine COVID-19 Randomized Controlled Trials




Figure 9. Random effects meta-analysis for all Randomized Controlled Trials. This plot shows pooled effects, see the specific outcome analyses for individual outcomes. Analysis validating pooled outcomes for COVID-19 can be found below. Effect extraction is pre-specified, using the most serious outcome reported. For details see the appendix.

RCTs have many potential biases

RCTs help to make study groups more similar and can provide a higher level of evidence, however they are subject to many biases²⁸, and analysis of double-blind RCTs has identified extreme levels of bias²⁹. For COVID-19, the overhead may delay treatment, dramatically compromising efficacy; they may encourage monotherapy for simplicity at the cost of efficacy which may rely on combined or synergistic effects; the participants that sign up may not reflect real world usage or the population that benefits most in terms of age, comorbidities, severity of illness, or other factors; standard of care may be compromised and unable to evolve quickly based on emerging research for new diseases; errors may be made in randomization and medication delivery; and investigators may have hidden agendas or vested interests influencing design, operation, analysis, reporting, and the potential for fraud. All of these biases have been observed with COVID-19 RCTs. There is no guarantee that a specific RCT provides a higher level of evidence.

Conflicts of interest for COVID-19 RCTs

RCTs are expensive and many RCTs are funded by pharmaceutical companies or interests closely aligned with pharmaceutical companies. For COVID-19, this creates an incentive to show efficacy for patented commercial products, and an incentive to show a lack of efficacy for inexpensive treatments. The bias is expected to be significant, for example *Als-Nielsen et al.* analyzed 370 RCTs from Cochrane reviews, showing that trials funded by

for-profit organizations were 5 times more likely to recommend the experimental drug compared with those funded by nonprofit organizations. For COVID-19, some major philanthropic organizations are largely funded by investments with extreme conflicts of interest for and against specific COVID-19 interventions.

RCTs for novel acute diseases requiring rapid treatment

High quality RCTs for novel acute diseases are more challenging, with increased ethical issues due to the urgency of treatment, increased risk due to enrollment delays, and more difficult design with a rapidly evolving evidence base. For COVID-19, the most common site of initial infection is the upper respiratory tract. Immediate treatment is likely to be most successful and may prevent or slow progression to other parts of the body. For a non-prophylaxis RCT, it makes sense to provide treatment in advance and instruct patients to use it immediately on symptoms, just as some governments have done by providing medication kits in advance. Unfortunately, no RCTs have been done in this way. Every treatment RCT to date involves delayed treatment. Among the 116 treatments we have analyzed, 66% of RCTs involve very late treatment 5+ days after onset. No non-prophylaxis COVID-19 RCTs match the potential real-world use of early treatments. They may more accurately represent results for treatments that require visiting a medical facility, e.g., those requiring intravenous administration.

RCT bias for widely available treatments

RCTs have a bias against finding an effect for interventions that are widely available — patients that believe they need the intervention are more likely to decline participation and take the intervention. RCTs for azelastine are more likely to enroll low-risk participants that do not need treatment to recover, making the results less applicable to clinical practice. This bias is likely to be greater for widely known treatments, and may be greater when the risk of a serious outcome is overstated. This bias does not apply to the typical pharmaceutical trial of a new drug that is otherwise unavailable.

Observational studies have been shown to be reliable

Evidence shows that observational studies can also provide reliable results. *Concato et al.* found that well-designed observational studies do not systematically overestimate the magnitude of the effects of treatment compared to RCTs. *Anglemyer et al.* analyzed reviews comparing RCTs to observational studies and found little evidence for significant differences in effect estimates. We performed a similar analysis across the 116 treatments we cover, showing no significant difference in the results of RCTs compared to observational studies, RR 1.00 [0.93-1.08]. Similar results are found for all low-cost treatments, RR 1.02 [0.93-1.13]. High-cost treatments show a non-significant trend towards RCTs showing greater efficacy, RR 0.92 [0.82-1.03]. Details can be found in the supplementary data. *Lee et al.* showed that only 14% of the guidelines of the Infectious Diseases Society of America were based on RCTs. Evaluation of studies relies on an understanding of the study and potential biases. Limitations in an RCT can outweigh the benefits, for example excessive dosages, excessive treatment delays, or remote survey bias may have a greater effect on results. Ethical issues may also prevent running RCTs for known effective treatments. For more on issues with RCTs see ^{35,36}.

For COVID-19, observational study results do not systematically differ from RCTs, RR 1.00 [0.93-1.08] across 116 treatments³¹.

Using all studies identifies efficacy 7+ months faster (8+ months for low-cost treatments)

Currently, 48 of the treatments we analyze show statistically significant efficacy or harm, defined as $\geq 10\%$ decreased risk or $>0\%$ increased risk from ≥ 3 studies. Of these, 60% have been confirmed in RCTs, with a mean delay of 7.2 months (68% with 8.3 months delay for low-cost treatments). The remaining treatments either have no RCTs, or the point estimate is consistent.

Summary

We need to evaluate each trial on its own merits. RCTs for a given medication and disease may be more reliable, however they may also be less reliable. For off-patent medications, very high conflict of interest trials may be more likely to be RCTs, and more likely to be large trials that dominate meta analyses.

Heterogeneity

Heterogeneity in COVID-19 studies arises from many factors including:

Treatment delay

The time between infection or the onset of symptoms and treatment may critically affect how well a treatment works. For example an antiviral may be very effective when used early but may not be effective in late stage disease, and may even be harmful. Oseltamivir, for example, is generally only considered effective for influenza when used within 0-36 or 0-48 hours^{37,38}. Baloxavir marboxil studies for influenza also show that treatment delay is critical — *Ikematsu et al.* report an 86% reduction in cases for post-exposure prophylaxis, *Hayden et al.* show a 33 hour reduction in the time to alleviation of symptoms for treatment within 24 hours and a reduction of 13 hours for treatment within 24-48 hours, and *Kumar et al.* report only 2.5 hours improvement for inpatient treatment.

Treatment delay	Result
Post-exposure prophylaxis	86% fewer cases ³⁹
<24 hours	-33 hours symptoms ⁴⁰
24-48 hours	-13 hours symptoms ⁴⁰
Inpatients	-2.5 hours to improvement ⁴¹

Table 3. Studies of baloxavir marboxil for influenza show that early treatment is more effective.

Figure 10 shows a mixed-effects meta-regression for efficacy as a function of treatment delay in COVID-19 studies from 116 treatments, showing that efficacy declines rapidly with treatment delay. Early treatment is critical for COVID-19.




Figure 10. Early treatment is more effective. Meta-regression showing efficacy as a function of treatment delay in COVID-19 studies from 116 treatments.

Patient demographics

Details of the patient population including age and comorbidities may critically affect how well a treatment works. For example, many COVID-19 studies with relatively young low-comorbidity patients show all patients recovering quickly with or without treatment. In such cases, there is little room for an effective treatment to improve results, for example as in *López-Medina et al.*

SARS-CoV-2 variants

Efficacy may depend critically on the distribution of SARS-CoV-2 variants encountered by patients. Risk varies significantly across variants⁴³, for example the Gamma variant shows significantly different characteristics⁴⁴⁻⁴⁷. Different mechanisms of action may be more or less effective depending on variants, for example the degree to which TMPRSS2 contributes to viral entry can differ across variants^{48,49}.

Treatment regimen

Effectiveness may depend strongly on the dosage and treatment regimen.

Medication quality

The quality of medications may vary significantly between manufacturers and production batches, which may significantly affect efficacy and safety. *Williams et al.* analyze ivermectin from 11 different sources, showing highly variable antiparasitic efficacy across different manufacturers. *Xu et al.* analyze a treatment from two different manufacturers, showing 9 different impurities, with significantly different concentrations for each manufacturer.

Other treatments

The use of other treatments may significantly affect outcomes, including supplements, other medications, or other interventions such as prone positioning. Treatments may be synergistic⁵²⁻⁶³, therefore efficacy may depend strongly on combined treatments.

Effect measured

Across all studies there is a strong association between different outcomes, for example improved recovery is strongly associated with lower mortality. However, efficacy may differ depending on the effect measured, for example a treatment may be more effective against secondary complications and have minimal effect on viral clearance.

Meta analysis

The distribution of studies will alter the outcome of a meta analysis. Consider a simplified example where everything is equal except for the treatment delay, and effectiveness decreases to zero or below with increasing delay. If there are many studies using very late treatment, the outcome may be negative, even though early treatment is very effective. All meta analyses combine heterogeneous studies, varying in population, variants, and potentially all factors above, and therefore may obscure efficacy by including studies where treatment is less effective. Generally, we expect the estimated effect size from meta analysis to be less than that for the optimal case. Looking at all studies is valuable for providing an overview of all research, important to avoid cherry-picking, and informative when a positive result is found despite combining less-optimal situations. However, the resulting estimate does not apply to specific cases such as early treatment in high-risk populations. While we present results for all studies, we also present treatment time and individual outcome analyses, which may be more informative for specific use cases.

Pooled Effects

Combining studies is required

For COVID-19, delay in clinical results translates into additional death and morbidity, as well as additional economic and societal damage. Combining the results of studies reporting different outcomes is required. There may be no mortality in a trial with low-risk patients, however a reduction in severity or improved viral clearance may translate into lower mortality in a high-risk population. Different studies may report lower severity, improved recovery, and lower mortality, and the significance may be very high when combining the results. "*The studies reported different outcomes*" is not a good reason for disregarding results. Pooling the results of studies reporting different outcomes allows us to use more of the available information. Logically we should, and do, use additional information when evaluating treatments—for example dose-response and treatment delay-response relationships provide additional evidence of efficacy that is considered when reviewing the evidence for a treatment.

Specific outcome and pooled analyses

We present both specific outcome and pooled analyses. In order to combine the results of studies reporting different outcomes we use the most serious outcome reported in each study, based on the thesis that improvement in the most serious outcome provides comparable measures of efficacy for a treatment. A critical advantage of this approach is simplicity and transparency. There are many other ways to combine evidence for different outcomes, along with additional evidence such as dose-response relationships, however these increase complexity.

Ethical and practical issues limit high-risk trials

Trials with high-risk patients may be restricted due to ethics for treatments that are known or expected to be effective, and they increase difficulty for recruiting. Using less severe outcomes as a proxy for more serious outcomes allows faster and safer collection of evidence.

Validating pooled outcome analysis for COVID-19

For many COVID-19 treatments, a reduction in mortality logically follows from a reduction in hospitalization, which follows from a reduction in symptomatic cases, which follows from a reduction in PCR positivity. We can directly test this for COVID-19.

Analysis of the the association between different outcomes across studies from all 116 treatments we cover confirms the validity of pooled outcome analysis for COVID-19. Figure 11 shows that lower hospitalization is very strongly associated with lower mortality ($p < 0.000000000001$). Similarly, Figure 12 shows that improved recovery is very strongly associated with lower mortality ($p < 0.000000000001$). Considering the extremes, *Singh et al.* show an association between viral clearance and hospitalization or death, with $p = 0.003$ after excluding one large outlier from a mutagenic treatment, and based on 44 RCTs including 52,384 patients. Figure 13 shows that improved viral clearance is strongly associated with fewer serious outcomes. The association is very similar to *Singh et al.*, with higher confidence due to the larger number of studies. As with *Singh et al.*, the confidence increases when excluding the outlier treatment, from $p = 0.00000036$ to $p = 0.00000016$.




Figure 11. Lower hospitalization is associated with lower mortality, supporting pooled outcome analysis.




Figure 12. Improved recovery is associated with lower mortality, supporting pooled outcome analysis.




Figure 11. Improved viral clearance is associated with fewer serious outcomes, supporting pooled outcome analysis.

Pooled outcomes identify efficacy 5 months faster (7 months for RCTs)

Currently, 48 of the treatments we analyze show statistically significant efficacy or harm, defined as $\geq 10\%$ decreased risk or $>0\%$ increased risk from ≥ 3 studies. 89% of these have been confirmed with one or more specific outcomes, with a mean delay of 4.8 months. When restricting to RCTs only, 56% of treatments showing statistically significant efficacy/harm with pooled effects have been confirmed with one or more specific outcomes, with a mean delay of 6.5 months. Figure 14 shows when treatments were found effective during the pandemic. Pooled outcomes often resulted in earlier detection of efficacy.

Time when COVID-19 studies showed efficacy



Figure 14. The time when studies showed that treatments were effective, defined as statistically significant improvement of $\geq 10\%$ from ≥ 3 studies. Pooled results typically show efficacy earlier than specific outcome results. Results from all studies often shows efficacy much earlier than when restricting to RCTs. Results reflect conditions as used in trials to date, these depend on the population treated, treatment delay, and treatment regimen.

Limitations

Pooled analysis could hide efficacy, for example a treatment that is beneficial for late stage patients but has no effect on viral clearance may show no efficacy if most studies only examine viral clearance. In practice, it is rare for a non-antiviral treatment to report viral clearance and to not report clinical outcomes; and in practice other sources of heterogeneity such as difference in treatment delay is more likely to hide efficacy.

Summary

Analysis validates the use of pooled effects and shows significantly faster detection of efficacy on average. However, as with all meta analyses, it is important to review the different studies included. We also present individual outcome analyses, which may be more informative for specific use cases.

Discussion

Publication bias

Publishing is often biased towards positive results, however evidence suggests that there may be a negative bias for inexpensive treatments for COVID-19. Both negative and positive results are very important for COVID-19, media in many countries prioritizes negative results for inexpensive treatments (inverting the typical incentive for scientists that value media recognition), and there are many reports of difficulty publishing positive results⁶⁵⁻⁶⁸. For azelastine, there is currently not enough data to evaluate publication bias with high confidence.

One method to evaluate bias is to compare prospective vs. retrospective studies. Prospective studies are more likely to be published regardless of the result, while retrospective studies are more likely to exhibit bias. For example, researchers may perform preliminary analysis with minimal effort and the results may influence their decision to continue. Retrospective studies also provide more opportunities for the specifics of data extraction and adjustments to influence results.

Figure 15 shows a scatter plot of results for prospective and retrospective studies. The median effect size for retrospective studies is 45% improvement, compared to 13% for prospective studies, suggesting a potential bias towards publishing results showing higher efficacy.




Figure 15. Prospective vs. retrospective studies. The diamonds show the results of random effects meta-analysis.

Conflicts of interest

Pharmaceutical drug trials often have conflicts of interest whereby sponsors or trial staff have a financial interest in the outcome being positive. Azelastine for COVID-19 lacks this because it is off-patent, has multiple manufacturers, and is very low cost. In contrast, most COVID-19 azelastine trials have been run by physicians on the front lines with the primary goal of finding the best methods to save human lives and minimize the collateral damage caused by COVID-19. While pharmaceutical companies are careful to run trials under optimal conditions (for example, restricting patients to those most likely to benefit, only including patients that can be treated soon after onset when necessary, and ensuring accurate dosing), not all azelastine trials represent the optimal conditions for efficacy.

Limitations

Summary statistics from meta analysis necessarily lose information. As with all meta analyses, studies are heterogeneous, with differences in treatment delay, treatment regimen, patient demographics, variants, conflicts of interest, standard of care, and other factors. We provide analyses for specific outcomes and by treatment delay, and we aim to identify key characteristics in the forest plots and summaries. Results should be viewed in the context of study characteristics.

Some analyses classify treatment based on early or late administration, as done here, while others distinguish between mild, moderate, and severe cases. Viral load does not indicate degree of symptoms — for example patients may have a high viral load while being asymptomatic. With regard to treatments that have antiviral properties, timing of treatment is critical — late administration may be less helpful regardless of severity.

Details of treatment delay per patient is often not available. For example, a study may treat 90% of patients relatively early, but the events driving the outcome may come from 10% of patients treated very late. Our 5 day cutoff for early treatment may be too conservative, 5 days may be too late in many cases.

Comparison across treatments is confounded by differences in the studies performed, for example dose, variants, and conflicts of interest. Trials with conflicts of interest may use designs better suited to the preferred outcome.

In some cases, the most serious outcome has very few events, resulting in lower confidence results being used in pooled analysis, however the method is simpler and more transparent. This is less critical as the number of studies increases. Restriction to outcomes with sufficient power may be beneficial in pooled analysis and improve accuracy when there are few studies, however we maintain our pre-specified method to avoid any retrospective changes.

Studies show that combinations of treatments can be highly synergistic and may result in many times greater efficacy than individual treatments alone⁵²⁻⁶³. Therefore standard of care may be critical and benefits may diminish or disappear if standard of care does not include certain treatments.

This real-time analysis is constantly updated based on submissions. Accuracy benefits from widespread review and submission of updates and corrections from reviewers. Less popular treatments may receive fewer reviews.

No treatment or intervention is 100% available and effective for all current and future variants. Efficacy may vary significantly with different variants and within different populations. All treatments have potential side effects. Propensity to experience side effects may be predicted in advance by qualified physicians. We do not provide medical advice. Before taking any medication, consult a qualified physician who can compare all options, provide personalized advice, and provide details of risks and benefits based on individual medical history and situations.

Other studies

Additional preclinical or review papers suggesting potential benefits of azelastine for COVID-19 include⁷¹⁻⁸¹. We have not reviewed these studies in detail.

Perspective

Results compared with other treatments

SARS-CoV-2 infection and replication involves a complex interplay of 50+ host and viral proteins and other factors¹⁸⁻²⁴, providing many therapeutic targets. Over 8,000 compounds have been predicted to reduce COVID-19 risk²⁵, either by directly minimizing infection or replication, by supporting immune system function, or by minimizing secondary complications. Figure 16 shows an overview of the results for azelastine in the context of multiple COVID-19 treatments, and Figure 17 shows a plot of efficacy vs. cost for COVID-19 treatments.

Efficacy in COVID-19 studies (pooled effects)

c19early.org
March 2025




Figure 16. Scatter plot showing results within the context of multiple COVID-19 treatments. Diamonds shows the results of random effects meta-analysis. 0.5% of 8,000+ proposed treatments show efficacy⁸².

Efficacy vs. cost for COVID-19 treatments

c19early.org
March 2025
Regdanvimab \$2,100




Figure 17. Efficacy vs. cost for COVID-19 treatments.


Conclusion

Significantly lower risk is seen for cases and viral clearance. 3 studies from 3 independent teams in 3 countries show significant benefit. Meta analysis using the most serious outcome reported shows 21% [-3-39%] lower risk, without reaching statistical significance. Results are worse for Randomized Controlled Trials.

Currently there is limited data, with only 310 patients in trials to date. Studies to date are from only 3 different groups.

Study Notes


Klussmann



RCT 90 outpatients showing potential benefit of azelastine nasal spray for reducing viral load. Patients were randomized to receive placebo, 0.02%, or 0.1% azelastine nasal spray for 11 days. The 0.1% azelastine group showed greater viral load reduction compared to placebo ($p=0.007$) and achieved PCR negativity earlier (48.2% vs 23.1% negative by day 11). Symptoms improved across all groups, with the 0.1% group showing slightly greater improvement (12.74 point reduction vs 11.12 for placebo) and significantly better improvement in shortness of breath on days 3-4. No significant differences were found in quality of life measures or WHO clinical progression scores.

Meiser

Azelastine CARVIN-II LATE TREATMENT DB RCT



Is **late** treatment with azelastine beneficial for COVID-19?

Double-blind RCT 251 patients in India (September 2022 - May 2023)

Improved viral clearance with azelastine ($p < 0.000001$)

c19early.org Meiser et al., Viruses, October 2024

RCT 294 low-risk subjects with mild COVID-19 showing significantly greater reduction in viral load with azelastine 0.1% nasal spray vs. placebo. There was no COVID-19 related hospitalization in either group. The reduction in viral load was significantly higher in the azelastine group at day 3, 6 and 11.


Authors report that treatment was associated with significant improvement in fever, weakness, and hypoxia compared to placebo, however detailed results are not provided for these and other symptoms.

Treatment delay from symptom onset is not specified, however the baseline viral load, lack of progression, and recovery profile are consistent with relatively late treatment among low-risk patients.

Time to recovery results are not clear, with an inconsistent HR and 95% CI reported 0.32 [0.83-2.32].

Reznikov

Azelastine for COVID-19 Reznikov et al. Prophylaxis



Does azelastine reduce COVID-19 infections?

Retrospective study in the USA

Fewer cases with azelastine ($p = 0.028$)

c19early.org Reznikov et al., Biochemical and Bioph..., Jan 2021

Retrospective 219,000 patients showing lower risk of COVID-19 with antihistamine H1RA use.

In Vitro study showing these drugs exhibit direct antiviral activity against SARS-CoV-2. Molecular docking suggests hydroxyzine and azelastine may exert antiviral effects by binding ACE2 and the sigma-1 receptor.

Appendix 1. Methods and Data

We perform ongoing searches of PubMed, medRxiv, Europe PMC, ClinicalTrials.gov, The Cochrane Library, Google Scholar, Research Square, ScienceDirect, Oxford University Press, the reference lists of other studies and meta-analyses, and submissions to the site c19early.org. Search terms are azelastine and COVID-19 or SARS-CoV-2. Automated searches are performed twice daily, with all matches reviewed for inclusion. All studies regarding the use of azelastine for COVID-19 that report a comparison with a control group are included in the main analysis. This is a living analysis and is updated regularly.

We extracted effect sizes and associated data from all studies. If studies report multiple kinds of effects then the most serious outcome is used in pooled analysis, while other outcomes are included in the outcome specific analyses. For example, if effects for mortality and cases are both reported, the effect for mortality is used, this may be different to the effect that a study focused on. If symptomatic results are reported at multiple times, we used the latest time, for example if mortality results are provided at 14 days and 28 days, the results at 28 days have preference. Mortality alone is preferred over combined outcomes. Outcomes with zero events in both arms are not used, the next most serious outcome with one or more events is used. For example, in low-risk populations with no mortality, a reduction in mortality with treatment is not possible, however a reduction in hospitalization, for example, is still valuable. Clinical outcomes are considered more important than viral test status. When basically all patients recover in both treatment and control groups, preference for viral clearance and recovery is given to results mid-recovery where available. After most or all patients have recovered there is little or no room for an effective treatment to do better, however faster recovery is valuable. If only individual symptom data is available, the most serious symptom has priority, for example difficulty breathing or low SpO₂ is more important than cough. When results provide an odds ratio, we compute the relative risk when possible, or convert to a relative risk according to⁸³. Reported confidence intervals and *p*-values were used when available, using adjusted values when provided. If multiple types of adjustments are reported propensity score matching and multivariable regression has preference over propensity score matching or weighting, which has preference over multivariable regression. Adjusted results have preference over unadjusted results for a more serious outcome when the adjustments significantly alter results. When needed, conversion between reported *p*-values and confidence intervals followed *Altman, Altman (B)*, and Fisher's exact test was used to calculate *p*-values for event data. If continuity correction for zero values is required, we use the reciprocal of the opposite arm with the sum of the correction factors equal to 1⁸⁶. Results are expressed with RR < 1.0 favoring treatment, and using the risk of a negative outcome when applicable (for example, the risk of death rather than the risk of survival). If studies only report relative continuous values such as relative times, the ratio of the time for the treatment group versus the time for the control group is used. Calculations are done in Python (3.13.2) with *scipy* (1.15.2), *pythonmeta* (1.26), *numpy* (1.26.4), *statsmodels* (0.14.4), and *plotly* (6.0.0).

Forest plots are computed using *PythonMeta*⁸⁷ with the DerSimonian and Laird random effects model (the fixed effect assumption is not plausible in this case) and inverse variance weighting. Results are presented with 95% confidence intervals. Heterogeneity among studies was assessed using the *I*² statistic. Mixed-effects meta-regression results are computed with R (4.4.0) using the *metafor* (4.6-0) and *rms* (6.8-0) packages, and using the most serious sufficiently powered outcome. For all statistical tests, a *p*-value less than 0.05 was considered statistically significant. *Grobid* 0.8.0 is used to parse PDF documents.

We have classified studies as early treatment if most patients are not already at a severe stage at the time of treatment (for example based on oxygen status or lung involvement), and treatment started within 5 days of the onset of symptoms. If studies contain a mix of early treatment and late treatment patients, we consider the treatment time of patients contributing most to the events (for example, consider a study where most patients are treated early but late treatment patients are included, and all mortality events were observed with late treatment patients). We note that a shorter time may be preferable. Antivirals are typically only considered effective when used within a shorter timeframe, for example 0-36 or 0-48 hours for oseltamivir, with longer delays not being effective^{37,38}.

We received no funding, this research is done in our spare time. We have no affiliations with any pharmaceutical companies or political parties.

A summary of study results is below. Please submit updates and corrections at <https://c19early.org/azlmeta.html>.

Early treatment

Effect extraction follows pre-specified rules as detailed above and gives priority to more serious outcomes. For pooled analyses, the first (most serious) outcome is used, which may differ from the effect a paper focuses on. Other outcomes are used in outcome specific analyses.

<p><i>Klussmann, 4/26/2023, Double Blind Randomized Controlled Trial, placebo-controlled, Germany, peer-reviewed, 27 authors, CARVIN trial.</i></p>	<p>risk of no recovery, 12.7% lower, RR 0.87, $p = 0.54$, treatment mean 12.74 (± 10.74) $n=29$, control mean 11.12 (± 9.45) $n=30$, 0.1%.</p>
	<p>risk of no recovery, 32.7% higher, RR 1.33, $p = 0.26$, treatment mean 8.38 (± 9.42) $n=31$, control mean 11.12 (± 9.45) $n=30$, 0.02%.</p>
	<p>risk of no recovery, 66.3% lower, RR 0.34, $p = 1.00$, treatment 0 of 29 (0.0%), control 1 of 30 (3.3%), NNT 30, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm), 0.1%.</p>
	<p>risk of no recovery, 67.0% lower, RR 0.33, $p = 0.49$, treatment 0 of 31 (0.0%), control 1 of 30 (3.3%), NNT 30, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm), 0.02%.</p>
	<p>viral load, 14.2% lower, relative load 0.86, $p = 0.22$, treatment mean 4.45 (± 2.25) $n=29$, control mean 3.82 (± 1.61) $n=30$, 0.1%.</p>
	<p>viral load, 7.3% lower, relative load 0.93, $p = 0.52$, treatment mean 4.12 (± 2.01) $n=31$, control mean 3.82 (± 1.61) $n=30$, 0.02%.</p>
	<p>risk of no viral clearance, 21.7% lower, RR 0.78, $p = 0.04$, treatment mean 24.14 (± 13.12) $n=29$, control mean 18.89 (± 4.7) $n=30$, relative AUC, 0.1%.</p>
	<p>risk of no viral clearance, 6.2% lower, RR 0.94, $p = 0.54$, treatment mean 24.14 (± 12.56) $n=31$, control mean 22.64 (± 4.7) $n=30$, relative AUC, 0.02%.</p>

Late treatment

Effect extraction follows pre-specified rules as detailed above and gives priority to more serious outcomes. For pooled analyses, the first (most serious) outcome is used, which may differ from the effect a paper focuses on. Other outcomes are used in outcome specific analyses.

<p><i>Meiser, 10/31/2024, Double Blind Randomized Controlled Trial, placebo-controlled, India, peer-reviewed, 8 authors, study period 27 September, 2022 - 16 May, 2023, trial CTRI/2022/09 (CARVIN-II).</i></p>	<p>risk of no recovery, 12.5% lower, RR 0.87, $p = 0.42$, treatment mean 0.8 (± 0.99) $n=122$, control mean 0.7 (± 0.96) $n=129$, WHO scale improvement, mid-recovery, day 3.</p>
	<p>viral load, 6.4% lower, relative load 0.94, $p < 0.001$, treatment mean 2.8 (± 0.27) $n=122$, control mean 2.62 (± 0.26) $n=129$, relative viral load reduction, mid-recovery, day 3.</p>

	viral load, 2.2% lower, relative load 0.98, $p = 0.001$, treatment mean 5.05 (± 0.29) $n=122$, control mean 4.94 (± 0.25) $n=129$, relative viral load reduction, mid-recovery, day 6.
	viral load, 1.3% lower, relative load 0.99, $p = 0.003$, treatment mean 5.93 (± 0.22) $n=122$, control mean 5.85 (± 0.2) $n=129$, relative viral load reduction, mid-recovery, day 11.
	risk of no viral clearance, 3.2% lower, RR 0.97, $p < 0.001$, treatment mean 1014.96 (± 35.25) $n=122$, control mean 982.98 (± 31.68) $n=129$, relative AUC.

Prophylaxis

Effect extraction follows pre-specified rules as detailed above and gives priority to more serious outcomes. For pooled analyses, the first (most serious) outcome is used, which may differ from the effect a paper focuses on. Other outcomes are used in outcome specific analyses.

Reznikov, 1/31/2021, retrospective, USA, peer-reviewed, 9 authors.	risk of case, 45.5% lower, RR 0.55, $p = 0.03$, adjusted per study, age groups combined.
	risk of case, 58.8% lower, OR 0.41, $p < 0.001$, adjusted per study, inverted to make OR<1 favor treatment, azelastine, 61+, multivariable, RR approximated with OR.
	risk of case, 28.6% lower, OR 0.71, $p = 0.17$, adjusted per study, inverted to make OR<1 favor treatment, azelastine, 31-60, multivariable, RR approximated with OR.

Supplementary Data

Supplementary Data

Footnotes

- a. Viral infection and replication involves attachment, entry, uncoating and release, genome replication and transcription, translation and protein processing, assembly and budding, and release. Each step can be disrupted by therapeutics.

References

1. **Rong** et al., *Persistence of spike protein at the skull-meninges-brain axis may contribute to the neurological sequelae of COVID-19*, Cell Host & Microbe, doi:10.1016/j.chom.2024.11.007.
2. **Yang** et al., *SARS-CoV-2 infection causes dopaminergic neuron senescence*, Cell Stem Cell, doi:10.1016/j.stem.2023.12.012.
3. **Scardua-Silva** et al., *Microstructural brain abnormalities, fatigue, and cognitive dysfunction after mild COVID-19*, Scientific Reports, doi:10.1038/s41598-024-52005-7.
4. **Hampshire** et al., *Cognition and Memory after Covid-19 in a Large Community Sample*, New England Journal of Medicine, doi:10.1056/NEJMoa2311330.

5. **Duloquin** et al., *Is COVID-19 Infection a Multiorganic Disease? Focus on Extrapulmonary Involvement of SARS-CoV-2*, Journal of Clinical Medicine, doi:10.3390/jcm13051397.
6. **Sodagar** et al., *Pathological Features and Neuroinflammatory Mechanisms of SARS-CoV-2 in the Brain and Potential Therapeutic Approaches*, Biomolecules, doi:10.3390/biom12070971.
7. **Sagar** et al., *COVID-19-associated cerebral microbleeds in the general population*, Brain Communications, doi:10.1093/braincomms/fcae127.
8. **Verma** et al., *Persistent Neurological Deficits in Mouse PASC Reveal Antiviral Drug Limitations*, bioRxiv, doi:10.1101/2024.06.02.596989.
9. **Panagea** et al., *Neurocognitive Impairment in Long COVID: A Systematic Review*, Archives of Clinical Neuropsychology, doi:10.1093/arclin/aca042.
10. **Ariza** et al., *COVID-19: Unveiling the Neuropsychiatric Maze—From Acute to Long-Term Manifestations*, Biomedicine, doi:10.3390/biomedicine12061147.
11. **Vashisht** et al., *Neurological Complications of COVID-19: Unraveling the Pathophysiological Underpinnings and Therapeutic Implications*, Viruses, doi:10.3390/v16081183.
12. **Ahmad** et al., *Neurological Complications and Outcomes in Critically Ill Patients With COVID-19: Results From International Neurological Study Group From the COVID-19 Critical Care Consortium*, The Neurohospitalist, doi:10.1177/19418744241292487.
13. **Wang** et al., *SARS-CoV-2 membrane protein induces neurodegeneration via affecting Golgi-mitochondria interaction*, Translational Neurodegeneration, doi:10.1186/s40035-024-00458-1.
14. **Eberhardt** et al., *SARS-CoV-2 infection triggers pro-atherogenic inflammatory responses in human coronary vessels*, Nature Cardiovascular Research, doi:10.1038/s44161-023-00336-5.
15. **Van Tin** et al., *Spike Protein of SARS-CoV-2 Activates Cardiac Fibrogenesis through NLRP3 Inflammasomes and NF- κ B Signaling*, Cells, doi:10.3390/cells13161331.
16. **Borka Balas** et al., *COVID-19 and Cardiac Implications—Still a Mystery in Clinical Practice*, Reviews in Cardiovascular Medicine, doi:10.31083/j.rcm2405125.
17. **AlTaweel** et al., *An In-Depth Insight into Clinical, Cellular and Molecular Factors in COVID-19-Associated Cardiovascular Ailments for Identifying Novel Disease Biomarkers, Drug Targets and Clinical Management Strategies*, Archives of Microbiology & Immunology, doi:10.26502/ami.936500177.
18. **Malone** et al., *Structures and functions of coronavirus replication–transcription complexes and their relevance for SARS-CoV-2 drug design*, Nature Reviews Molecular Cell Biology, doi:10.1038/s41580-021-00432-z.
19. **Murigneux** et al., *Proteomic analysis of SARS-CoV-2 particles unveils a key role of G3BP proteins in viral assembly*, Nature Communications, doi:10.1038/s41467-024-44958-0.
20. **Lv** et al., *Host proviral and antiviral factors for SARS-CoV-2*, Virus Genes, doi:10.1007/s11262-021-01869-2.
21. **Lui** et al., *Nsp1 facilitates SARS-CoV-2 replication through calcineurin-NFAT signaling*, Virology, doi:10.1128/mbio.00392-24.
22. **Niarakis** et al., *Drug-target identification in COVID-19 disease mechanisms using computational systems biology approaches*, Frontiers in Immunology, doi:10.3389/fimmu.2023.1282859.
23. **Katiyar** et al., *SARS-CoV-2 Assembly: Gaining Infectivity and Beyond*, Viruses, doi:10.3390/v16111648.
24. **Wu** et al., *Decoding the genome of SARS-CoV-2: a pathway to drug development through translation inhibition*, RNA Biology, doi:10.1080/15476286.2024.2433830.
25. **c19early.org**, c19early.org/treatments.html.
26. **Hamdan** et al., *In silico Evaluation of H1-Antihistamine as Potential Inhibitors of SARS-CoV-2 RNA-dependent RNA Polymerase: Repurposing Study of COVID-19 Therapy*, Turkish Journal of Pharmaceutical Sciences, doi:10.4274/tjps.galenos.2024.49768.
27. **Reznikov** et al., *Identification of antiviral antihistamines for COVID-19 repurposing*, Biochemical and Biophysical Research Communications, doi:10.1016/j.bbrc.2020.11.095.
28. **Jadad** et al., *Randomized Controlled Trials: Questions, Answers, and Musings, Second Edition*, doi:10.1002/9780470691922.
29. **Götzsche**, P., *Bias in double-blind trials*, Doctoral Thesis, University of Copenhagen, www.scientificfreedom.dk/2023/05/16/bias-in-double-blind-trials-doctoral-thesis/.

30. **Als-Nielsen** et al., *Association of Funding and Conclusions in Randomized Drug Trials*, JAMA, doi:10.1001/jama.290.7.921.
31. **c19early.org (B)**, c19early.org/azlsupp.html#fig_rctobs.
32. **Concato** et al., NEJM, 342:1887-1892, doi:10.1056/NEJM200006223422507.
33. **Anglemyer** et al., *Healthcare outcomes assessed with observational study designs compared with those assessed in randomized trials*, Cochrane Database of Systematic Reviews 2014, Issue 4, doi:10.1002/14651858.MR000034.pub2.
34. **Lee** et al., *Analysis of Overall Level of Evidence Behind Infectious Diseases Society of America Practice Guidelines*, Arch Intern Med., 2011, 171:1, 18-22, doi:10.1001/archinternmed.2010.482.
35. **Deaton** et al., *Understanding and misunderstanding randomized controlled trials*, Social Science & Medicine, 210, doi:10.1016/j.socscimed.2017.12.005.
36. **Nichol** et al., *Challenging issues in randomised controlled trials*, Injury, 2010, doi: 10.1016/j.injury.2010.03.033, www.injuryjournal.com/article/S0020-1383(10)00233-0/fulltext.
37. **Treanor** et al., *Efficacy and Safety of the Oral Neuraminidase Inhibitor Oseltamivir in Treating Acute Influenza: A Randomized Controlled Trial*, JAMA, 2000, 283:8, 1016-1024, doi:10.1001/jama.283.8.1016.
38. **McLean** et al., *Impact of Late Oseltamivir Treatment on Influenza Symptoms in the Outpatient Setting: Results of a Randomized Trial*, Open Forum Infect. Dis. September 2015, 2:3, doi:10.1093/ofid/ofv100.
39. **Ikematsu** et al., *Baloxavir Marboxil for Prophylaxis against Influenza in Household Contacts*, New England Journal of Medicine, doi:10.1056/NEJMoa1915341.
40. **Hayden** et al., *Baloxavir Marboxil for Uncomplicated Influenza in Adults and Adolescents*, New England Journal of Medicine, doi:10.1056/NEJMoa1716197.
41. **Kumar** et al., *Combining baloxavir marboxil with standard-of-care neuraminidase inhibitor in patients hospitalised with severe influenza (FLAGSTONE): a randomised, parallel-group, double-blind, placebo-controlled, superiority trial*, The Lancet Infectious Diseases, doi:10.1016/S1473-3099(21)00469-2.
42. **López-Medina** et al., *Effect of Ivermectin on Time to Resolution of Symptoms Among Adults With Mild COVID-19: A Randomized Clinical Trial*, JAMA, doi:10.1001/jama.2021.3071.
43. **Korves** et al., *SARS-CoV-2 Genetic Variants and Patient Factors Associated with Hospitalization Risk*, medRxiv, doi:10.1101/2024.03.08.24303818.
44. **Faria** et al., *Genomics and epidemiology of the P.1 SARS-CoV-2 lineage in Manaus, Brazil*, Science, doi:10.1126/science.abh2644.
45. **Nonaka** et al., *SARS-CoV-2 variant of concern P.1 (Gamma) infection in young and middle-aged patients admitted to the intensive care units of a single hospital in Salvador, Northeast Brazil, February 2021*, International Journal of Infectious Diseases, doi:10.1016/j.ijid.2021.08.003.
46. **Karita** et al., *Trajectory of viral load in a prospective population-based cohort with incident SARS-CoV-2 G614 infection*, medRxiv, doi:10.1101/2021.08.27.21262754.
47. **Zavascki** et al., *Advanced ventilatory support and mortality in hospitalized patients with COVID-19 caused by Gamma (P.1) variant of concern compared to other lineages: cohort study at a reference center in Brazil*, Research Square, doi:10.21203/rs.3.rs-910467/v1.
48. **Willett** et al., *The hyper-transmissible SARS-CoV-2 Omicron variant exhibits significant antigenic change, vaccine escape and a switch in cell entry mechanism*, medRxiv, doi:10.1101/2022.01.03.21268111.
49. **Peacock** et al., *The SARS-CoV-2 variant, Omicron, shows rapid replication in human primary nasal epithelial cultures and efficiently uses the endosomal route of entry*, bioRxiv, doi:10.1101/2021.12.31.474653.
50. **Williams, T.**, *Not All Ivermectin Is Created Equal: Comparing The Quality of 11 Different Ivermectin Sources*, Do Your Own Research, doyourownresearch.substack.com/p/not-all-ivermectin-is-created-equal.
51. **Xu** et al., *A study of impurities in the repurposed COVID-19 drug hydroxychloroquine sulfate by UHPLC-Q/TOF-MS and LC-SPE-NMR*, Rapid Communications in Mass Spectrometry, doi:10.1002/rcm.9358.
52. **Jitobaom** et al., *Favipiravir and Ivermectin Showed in Vitro Synergistic Antiviral Activity against SARS-CoV-2*, Research Square, doi:10.21203/rs.3.rs-941811/v1.
53. **Jitobaom (B)** et al., *Synergistic anti-SARS-CoV-2 activity of repurposed anti-parasitic drug combinations*, BMC Pharmacology and Toxicology, doi:10.1186/s40360-022-00580-8.

54. **Jeffreys** et al., Remdesivir-ivermectin combination displays synergistic interaction with improved in vitro activity against SARS-CoV-2, *International Journal of Antimicrobial Agents*, doi:10.1016/j.ijantimicag.2022.106542.
55. **Ostrov** et al., *Highly Specific Sigma Receptor Ligands Exhibit Anti-Viral Properties in SARS-CoV-2 Infected Cells*, *Pathogens*, doi:10.3390/pathogens10111514.
56. **Alsaïdi** et al., *Griffithsin and Carrageenan Combination Results in Antiviral Synergy against SARS-CoV-1 and 2 in a Pseudoviral Model*, *Marine Drugs*, doi:10.3390/md19080418.
57. **Andreani** et al., *In vitro testing of combined hydroxychloroquine and azithromycin on SARS-CoV-2 shows synergistic effect*, *Microbial Pathogenesis*, doi:10.1016/j.micpath.2020.104228.
58. **De Forni** et al., *Synergistic drug combinations designed to fully suppress SARS-CoV-2 in the lung of COVID-19 patients*, *PLoS ONE*, doi:10.1371/journal.pone.0276751.
59. **Wan** et al., *Synergistic inhibition effects of andrographolide and baicalin on coronavirus mechanisms by downregulation of ACE2 protein level*, *Scientific Reports*, doi:10.1038/s41598-024-54722-5.
60. **Said** et al., *The effect of Nigella sativa and vitamin D3 supplementation on the clinical outcome in COVID-19 patients: A randomized controlled clinical trial*, *Frontiers in Pharmacology*, doi:10.3389/fphar.2022.1011522.
61. **Fiaschi** et al., *In Vitro Combinatorial Activity of Direct Acting Antivirals and Monoclonal Antibodies against the Ancestral B.1 and BQ.1.1 SARS-CoV-2 Viral Variants*, *Viruses*, doi:10.3390/v16020168.
62. **Xing** et al., *Published anti-SARS-CoV-2 in vitro hits share common mechanisms of action that synergize with antivirals*, *Briefings in Bioinformatics*, doi:10.1093/bib/bbab249.
63. **Thairu** et al., *A Comparison of Ivermectin and Non Ivermectin Based Regimen for COVID-19 in Abuja: Effects on Virus Clearance, Days-to-discharge and Mortality*, *Journal of Pharmaceutical Research International*, doi:10.9734/jpri/2022/v34i44A36328.
64. **Singh** et al., *The relationship between viral clearance rates and disease progression in early symptomatic COVID-19: a systematic review and meta-regression analysis*, *Journal of Antimicrobial Chemotherapy*, doi:10.1093/jac/dkae045.
65. **Meneguzzo**, A., *Médica defende tratamento precoce da Covid-19*, www.youtube.com/watch?v=X5FCrIm_19U.
66. **Boulware**, D., *Comments regarding paper rejection*, twitter.com/boulware_dr/status/1311331372884205570.
67. **Meeus**, G., *Online Comment*, twitter.com/gertmeeus_MD/status/1386636373889781761.
68. **twitter.com**, twitter.com/KashPrime/status/1768487878454124914.
69. **Klussmann** et al., *Early intervention with azelastine nasal spray may reduce viral load in SARS-CoV-2 infected patients*, *Scientific Reports*, doi:10.1038/s41598-023-32546-z.
70. **Meiser** et al., *Azelastine Nasal Spray in Non-Hospitalized Subjects with Mild COVID-19 Infection: A Randomized Placebo-Controlled, Parallel-Group, Multicentric, Phase II Clinical Trial*, *Viruses*, doi:10.3390/v16121914.
71. **Konrat** et al., *The Anti-histamine Azelastine, Identified by Computational Drug Repurposing, Inhibits SARS-CoV-2 Infection in Reconstituted Human Nasal Tissue In Vitro*, *bioRxiv*, doi:10.1101/2020.09.15.296228.
72. **Ge** et al., *Azelastine inhibits viropexis of SARS-CoV-2 spike pseudovirus by binding to SARS-CoV-2 entry receptor ACE2*, *Virology*, doi:10.1016/j.virol.2021.05.009.
73. **Klussmann (B)** et al., *COVID-19: Azelastine nasal spray Reduces Virus-load In Nasal swabs (CARVIN). Early intervention with azelastine nasal sprays reduces viral load in SARS-CoV-2 infected patients. First report on a double-blind placebo-controlled phase II clinical trial.*, *Research Square*, doi:10.21203/rs.3.rs-864566/v1.
74. **Fischhuber** et al., *Antiviral Potential of Azelastine against Major Respiratory Viruses*, *Viruses*, doi:10.3390/v15122300.
75. **Yu** et al., *The histamine receptor H1 acts as an alternative receptor for SARS-CoV-2*, *mBio*, doi:10.1128/mbio.01088-24.
76. **Loucera** et al., *Drug repurposing for COVID-19 using machine learning and mechanistic models of signal transduction circuits related to SARS-CoV-2 infection*, *Signal Transduction and Targeted Therapy*, doi:10.1038/s41392-020-00417-y.
77. **Gysi** et al., *Network Medicine Framework for Identifying Drug Repurposing Opportunities for COVID-19*, *arXiv*, doi:10.48550/arXiv.2004.07229.
78. **Odhar** et al., *Molecular docking and dynamics simulation of FDA approved drugs with the main protease from 2019 novel coronavirus*, *Bioinformatics*, doi:10.6026/97320630016236.
79. **Ghahremanpour** et al., *Identification of 14 Known Drugs as Inhibitors of the Main Protease of SARS-CoV-2*, *bioRxiv*, doi:10.1101/2020.08.28.271957.

80. **Ghahremanpour (B)** et al., *Identification of 14 Known Drugs as Inhibitors of the Main Protease of SARS-CoV-2*, ACS Medicinal Chemistry Letters, doi:10.1021/acsmchemlett.0c00521.
81. **Hosseini** et al., *Computational molecular docking and virtual screening revealed promising SARS-CoV-2 drugs*, Precision Clinical Medicine, doi:10.1093/pcmedi/pbab001.
82. **c19early.org (C)**, c19early.org/timeline.html.
83. **Zhang** et al., *What's the relative risk? A method of correcting the odds ratio in cohort studies of common outcomes*, JAMA, 80:19, 1690, doi:10.1001/jama.280.19.1690.
84. **Altman, D.**, *How to obtain the P value from a confidence interval*, BMJ, doi:10.1136/bmj.d2304.
85. **Altman (B)** et al., *How to obtain the confidence interval from a P value*, BMJ, doi:10.1136/bmj.d2090.
86. **Sweeting** et al., *What to add to nothing? Use and avoidance of continuity corrections in meta-analysis of sparse data*, Statistics in Medicine, doi:10.1002/sim.1761.
87. **Deng, H.**, *PyMeta, Python module for meta-analysis*, www.pymeta.com/.