

Vitamin D for COVID-19: real-time meta analysis of 302 studies (116 treatment studies and 186 sufficiency studies)

@CovidAnalysis, Oct 2, 2023, Version 260

<https://c19early.org/dmeta.html>

- 116 treatment studies show statistically significant lower risk for **mortality**, **ICU admission**, **hospitalization**, and **cases**. 58 studies from 54 independent teams in 20 countries show statistically significant lower risk.
- Random effects meta-analysis with pooled effects using the most serious outcome reported shows 60% [40-74%] and 36% [31-42%] lower risk for **early treatment** and for **all studies**. Results are similar for higher quality studies, peer-reviewed studies, and mortality: early treatment - 68% [45-82%], 57% [36-71%], 68% [39-84%]; all - 37% [31-42%], 40% [34-46%], 36% [27-43%].
- Late stage treatment with **calcifediol** or **calcitriol** is more effective than **cholecalciferol**: 65% [41-79%] vs. 38% [25-49%].
- 186 sufficiency studies show a strong association between vitamin D sufficiency and outcomes, with 52% [48-56%] lower risk for higher levels.
- No **treatment** or intervention is 100% effective. All practical, effective, and safe means should be used based on risk/benefit analysis. Multiple treatments are typically used in combination, and **other treatments** may be more effective. The quality of non-prescription supplements can vary widely *Crawford, Crighton*.
- All data and sources to reproduce this paper are in the **appendix**. Other meta analyses show significant improvements with vitamin D treatment for mortality *Argano, D'Ecclesiis, Hariyanto, Hosseini, Nikniaz, Shah, Xie*, mechanical ventilation *Hariyanto, Meng, Shah, Xie*, ICU admission *Hariyanto, Hosseini, Meng, Shah, Tentolouris, Xie*, hospitalization *Argano*, severity *D'Ecclesiis, Nikniaz, Varikasuvu, Xie*, and cases *Varikasuvu*.

Vitamin D for COVID-19

Improvement, Studies, Patients

All studies 36% 116 184,103

Mortality 36% 66 63,148

Ventilation 16% 18 8,140

ICU admission 47% 26 40,386

Hospitalization 20% 23 86,202

Cases 15% 27 134,493

RCTs 29% 28 42,124

RCT mortality 31% 15 1,949

RCT ICU 35% 11 36,116

Sufficiency 52% 186 233,340

Cholecalciferol 35% 102 174,764

Calcifediol 50% 14 9,339

Prophylaxis 31% 58 130,622

Early 60% 11 43,587


Late 44% 47 9,894

after exclusions

c19early.org

October 2023


Relative Risk



Evolution of COVID-19 clinical evidence

c19early.org

October 2023



HIGHLIGHTS

Vitamin D reduces risk for COVID-19 with very high confidence for [mortality](#), [ICU admission](#), [hospitalization](#), [recovery](#), [cases](#), [viral clearance](#), and in [pooled analysis](#), high confidence for [progression](#), and very low confidence for [ventilation](#).


We show traditional outcome specific analyses and combined evidence from all studies, incorporating treatment delay, a primary confounding factor in COVID-19 studies.


Real-time [updates and corrections](#), transparent analysis with all results in the same format, consistent protocol for [57 treatments](#).

All 116 vitamin D COVID-19 treatment studies

c19early.org

October 2023





¹ CT: study uses combined treatment

Tau² = 0.10, I² = 88.5%, p < 0.0001

Effect extraction pre-specified
(most serious outcome, see appendix)

0 0.25 0.5 0.75 1 1.25 1.5 1.75 2+
Favors vitamin D Favors contro **A**




Figure 1. A. Random effects meta-analysis of treatment studies. This plot shows pooled effects, analysis for individual outcomes is below, and more details on pooled effects can be found in the heterogeneity section. Effect extraction is pre-specified, using the most serious outcome reported. Simplified dosages are shown for comparison, these are the total dose in the first five days for treatment, and the monthly dose for prophylaxis. Calcifediol, calcitriol, and paricalcitol treatment are indicated with (c), (t), and (p). For details of effect extraction and full dosage information see the [appendix](#). **B. Scatter plot**

showing the distribution of effects reported in sufficiency studies and treatment studies. Diamonds show the results of random effects meta-analysis. **C. Scatter plot showing the most serious outcome in all studies in the context of multiple COVID-19 treatments.** Diamonds show the results of random effects meta-analysis for each treatment. **D. Timeline of results in vitamin D treatment studies.** The marked dates indicate the time when efficacy was known with a statistically significant improvement of $\geq 10\%$ from ≥ 3 studies for pooled outcomes, one or more specific outcome, pooled outcomes in RCTs, and one or more specific outcome in RCTs. Efficacy based on RCTs only was delayed by 10.8 months, compared to using all studies. Efficacy based on specific outcomes in RCTs was delayed by 2.2 months, compared to using pooled outcomes in RCTs.

Introduction

We analyze all significant controlled studies regarding vitamin D and COVID-19. Search methods, inclusion criteria, effect extraction criteria (more serious outcomes have priority), all individual study data, PRISMA answers, and statistical methods are detailed in Appendix 1. We perform random-effects meta analysis for all treatment studies, Randomized Controlled Trials, peer-reviewed studies, studies using cholecalciferol, studies using calcifediol/calcitriol, and for specific outcomes: mortality, mechanical ventilation, ICU admission, hospitalization, and case results. Results are presented for prophylaxis, early treatment, and late treatment. Separately, we perform random-effects meta analysis for studies that analyze outcomes based on vitamin D sufficiency (non-treatment studies).

Vitamin D. Vitamin D is a steroid hormone that helps regulate the immune system by binding to specific receptors and activating genes involved in immune defense. It increases the production of antimicrobial proteins, like cathelicidin and defensins, which fight a variety of pathogens, including bacteria, viruses, and fungi. Vitamin D supports the immune system by boosting our natural defenses and promoting healthy cell connections. It helps clear respiratory pathogens through processes like apoptosis and autophagy and regulates toll-like receptors, which play a key role in immunity. Vitamin D also aids in immune cell maturation, balances inflammation, and reduces the production of proinflammatory cytokines. Vitamin D has been shown to downregulate angiotensin-converting enzyme-2 (ACE-2) receptors, which play a role in COVID-19 infection. By suppressing the production of ACE-2 and related molecules, vitamin D increases antioxidant and anti-inflammatory effects, enhances antimicrobial defenses, reduces cytokine storms, and promotes a protective immune response, all of which help decrease the severity of the disease. The European Food Safety Authority has found evidence for a causal relationship between the intake of vitamin D and optimal immune system function *Galmés, Galmés (B)*.

Vitamin D undergoes two conversion steps before reaching the biologically active form as shown in Figure 2. The first step is conversion to calcidiol, or 25(OH)D, in the liver. The second is conversion to calcitriol, or 1,25(OH)₂D, which occurs in the kidneys, the immune system, and elsewhere. Calcitriol is the active, steroid-hormone form of vitamin D, which binds with vitamin D receptors found in most cells in the body. Vitamin D was first identified in relation to bone health, but is now known to have multiple functions, including an important role in the immune system *Carlberg, Martens*. For example, *Quraishi* show a strong association between pre-operative vitamin D levels and hospital-acquired infections, as shown in Figure 3. There is a significant delay involved in the conversion from cholecalciferol, therefore calcifediol (calcidiol) or calcitriol may be preferable for treatment.




Figure 2. Simplified view of vitamin D sources and conversion.




Figure 3. Risk of hospital-acquired infections as a function of pre-operative vitamin D levels, from *Quraishi*.

Sufficiency. Many vitamin D studies analyze outcomes based on serum vitamin D levels which may be maintained via sun exposure, diet, or supplementation. We refer to these studies as sufficiency studies, as they typically present outcomes based on vitamin D sufficiency. These studies do not establish a causal link between vitamin D and outcomes. In general, low vitamin D levels are correlated with many other factors that may influence COVID-19 susceptibility and severity. Therefore, beneficial effects found in these studies may be due to factors other than vitamin D. On the other hand, if vitamin D is causally linked to the observed benefits, it is possible that adjustments for correlated factors could obscure this relationship. COVID-19 disease may also affect vitamin D levels *Silva*, suggesting additional caution in interpreting results for studies where the vitamin D levels are measured during the disease. For these reasons, we analyze sufficiency studies separately from treatment studies. We include all sufficiency studies that provide a comparison between two groups with low and high levels. Some studies only provide results as a function of change in vitamin D levels *Butler-Laporte, Gupta, Raisi-Estabragh*, which may not be indicative of results for deficiency/insufficiency versus sufficiency (increasing already sufficient levels may be less useful for example). Some studies show the average vitamin D level for patients in different groups, *Abdulameer, Aci, Al-Daghri, Alarслан, Azadeh, Beheshti, Chodick, D'Avolio, Desai, di Filippo, Ersöz, Hosseini (B), Jabbar, Jain, Kerget, Kumar, Latifi-Pupovci, Mansour, Mardani, Morad, Nicolescu, Pop-Kostova, Qu, Ranjbar, Rathod, Saeed, Saeed (B), Schmitt, Shannak, Sinnberg, Soltani-Zangbar, Takase, Vassiliou*, most of which show lower D levels for worse outcomes. Other studies analyze vitamin D status and outcomes in geographic regions, *Bakaloudi, Jayawardena, Marik, Papadimitriou, Rhodes, Sooriyaarachchi, Walrand, Yadav*, all finding worse outcomes to be more likely with lower D levels.

Sufficiency studies vary widely in terms of when vitamin D levels were measured, the cutoff level used, and the population analyzed (for example studies with hospitalized patients exclude the effect of vitamin D on the risk of hospitalization). We do not analyze sufficiency studies in more detail because there are many controlled treatment studies that provide better information on the use of vitamin D as a treatment for COVID-19. A more detailed analysis of sufficiency studies can be found in *Chiodini*. *Mishra* present a systematic review and meta analysis showing that vitamin D levels are significantly associated with COVID-19 cases.

Treatment. For studies regarding treatment with vitamin D, we distinguish three stages as shown in Figure 4. Prophylaxis refers to regularly taking vitamin D before being infected in order to minimize the severity of infection. Due to the mechanism of action, vitamin D is unlikely to completely prevent infection, although it may prevent infection from reaching a level detectable by PCR. Early Treatment refers to treatment immediately or soon after symptoms appear, while Late Treatment refers to more delayed treatment.




Figure 4. Treatment stages.

Preclinical Research

6 *In Silico* studies support the efficacy of vitamin D *Al-Mazaideh, Chellasamy, Mansouri, Pandya, Qayyum, Song*.

5 *In Vitro* studies support the efficacy of vitamin D *DiGuilio, Mok, Pickard, Rybakovsky, Sposito*.

An *In Vivo* animal study supports the efficacy of vitamin D *Fernandes de Souza*.

Preclinical research is an important part of the development of treatments, however results may be very different in clinical trials. Preclinical results are not used in this paper.

Results

Table 1 summarizes the results for all stages combined, with different exclusions, for specific outcomes, and for sufficiency (non-treatment) studies. Table 2 shows results by treatment stage. Figure 5 plots individual results by treatment stage. Figure 6, 7, 8, 9, 10, 11, 12, 13, and 14 show forest plots for treatment studies with pooled effects, peer-reviewed studies, cholecalciferol studies, calcifediol/calcitriol studies, and for studies reporting mortality, mechanical ventilation, ICU admission, hospitalization, and case results only. Figure 15 shows a forest plot for random effects meta-analysis of sufficiency (non-treatment) studies.

	<i>Improvement</i>	<i>Studies</i>	<i>Patients</i>	<i>Authors</i>
All studies	36% [31-42%] $p < 0.0001$ ****	116	184,103	1,184
After exclusions	40% [34-46%] $p < 0.0001$ ****	80	160,805	850
Peer-reviewed studies	37% [31-42%] $p < 0.0001$ ****	109	182,283	1,113
Randomized Controlled Trials	29% [18-39%] $p < 0.0001$ ****	28	42,124	332
RCTs after exclusions	38% [18-52%] $p = 0.00064$ ***	18	40,711	234
Cholecalciferol	35% [28-41%] $p < 0.0001$ ****	102	174,764	1,013
Calcifediol/calcitriol	50% [30-64%] $p < 0.0001$ ****	14	9,339	171
Mortality	36% [27-43%] $p < 0.0001$ ****	66	63,148	645
Ventilation	16% [-8-35%] $p = 0.18$	18	8,140	209
ICU admission	47% [29-61%] $p < 0.0001$ ****	26	40,386	299
Hospitalization	20% [8-30%] $p = 0.0015$ **	23	86,202	236
Recovery	26% [12-38%] $p = 0.00072$ ***	11	840	102
Cases	15% [6-23%] $p = 0.0011$ **	27	134,493	317
Viral	52% [30-67%] $p = 0.00014$ ***	4	200	26
RCT mortality	31% [6-50%] $p = 0.02$ *	15	1,949	174
RCT ventilation	20% [-0-36%] $p = 0.051$	9	5,362	119
RCT ICU admission	35% [8-55%] $p = 0.015$ *	11	36,116	159
RCT hospitalization	21% [-4-40%] $p = 0.087$	8	39,713	107
Sufficiency	52% [48-56%] $p < 0.0001$ ****	186	233,340	1,620

Table 1. Random effects meta-analysis for all stages combined, with different exclusions, for specific outcomes, and for sufficiency (non-treatment) studies. Results show the percentage improvement with treatment and the 95% confidence interval. * $p < 0.05$ ** $p < 0.01$ *** $p < 0.001$ **** $p < 0.0001$.

	Early treatment	Late treatment	Prophylaxis
All studies	60% [40-74%] ****	44% [32-53%] ****	31% [23-38%] ****
After exclusions	68% [45-82%] ****	64% [51-74%] ****	29% [21-36%] ****
Peer-reviewed studies	57% [36-71%] ****	43% [31-53%] ****	32% [24-39%] ****
Randomized Controlled Trials	32% [8-50%] *	34% [14-49%] **	23% [-12-47%]
RCTs after exclusions	65% [-65-92%]	50% [25-67%] ***	23% [-12-47%]
Cholecalciferol	60% [40-74%] ****	38% [25-49%] ****	30% [22-38%] ****
Calcifediol/calcitriol		65% [41-79%] ***	36% [13-54%] **
Mortality	68% [39-84%] ***	42% [29-53%] ****	23% [9-34%] **
Ventilation	97% [56-100%] *	6% [-21-26%]	38% [-3-63%]
ICU admission	87% [-143-99%]	48% [25-64%] ***	46% [22-63%] **
Hospitalization	90% [-453-100%]	20% [6-32%] **	13% [-4-27%]
Recovery	31% [7-49%] *	27% [4-45%] *	
Cases			15% [6-23%] **
Viral	52% [24-70%] **	53% [8-76%] *	
RCT mortality		31% [6-50%] *	
RCT ventilation		21% [0-37%] *	-95% [-3010-88%]
RCT ICU admission		38% [10-58%] *	-0% [-301-75%]
RCT hospitalization		29% [10-44%] **	-26% [-92-17%]

Table 2. Random effects meta-analysis results by treatment stage. Results show the percentage improvement with treatment, the 95% confidence interval, and the number of studies for the stage. * $p < 0.05$ ** $p < 0.01$ *** $p < 0.001$ **** $p < 0.0001$.





Figure 5. Results by treatment stage.

All 116 vitamin D COVID-19 treatment studies

c19early.org

October 2023






Figure 6. Random effects meta-analysis for treatment studies. Effect extraction is pre-specified, using the most serious outcome reported, see the [appendix](#) for details.

All 109 vitamin D COVID-19 peer reviewed studies

c19early.org

October 2023

	Improvement, RR [CI]	Treatment	Control	Dose (5d)	
Annweiler	89% 0.11 [0.03-0.48]	death	10/57	5/9	80,000IU
Annweiler	63% 0.37 [0.06-2.21]	death	3/16	10/32	80,000IU
Burahee	93% 0.07 [0.01-0.54]	death	0/12	2/2	400,000IU
Sánchez-Zuno (RCT)	89% 0.11 [0.01-0.91]	severe case	0/22	4/20	50,000IU
Efrid	49% 0.51 [0.23-1.17]	death	11/544	413/15,794	varies
Valecha	87% 0.13 [0.01-2.43]	ICU	0/30	3/25	
Khan (RCT)	33% 0.67 [0.37-1.19]	no recov.	10/25	15/25	1,800IU
Hunt	47% 0.53 [0.37-0.77]	death	43/1,019	1,569/25,489	n/a
Said (RCT)	42% 0.58 [0.09-3.47]	recovery	30 (n)	30 (n)	10,000IU
Din Ujjan (RCT)	29% 0.71 [0.50-1.03]	no recov.	15/25	21/25	1,800IU
Early treatment 57% 0.43 [0.29-0.64]					57% lower risk

Tau² = 0.17, I² = 59.6%, p < 0.0001

	Improvement, RR [CI]	Treatment	Control	Dose (5d)	
Tan	80% 0.20 [0.04-0.93]	oxygen	3/17	16/26	5,000IU
Krishnan	19% 0.81 [0.49-1.34]	death	8/16	84/136	n/a
Castillo (RCT)	85% 0.15 [0.01-2.93]	death	0/50	2/26	0.8mg (c)
Rastogi (RCT)	53% 0.47 [0.24-0.92]	viral+	6/16	19/24	300,000IU
Murai (DB RCT)	-49% 1.49 [0.55-4.05]	death	9/119	6/118	200,000IU
Ling	80% 0.20 [0.08-0.48]	death	73 (n)	253 (n)	40,000IU
Jevalikar	82% 0.18 [0.02-1.69]	death	1/128	3/69	60,000IU
Giannini	37% 0.63 [0.35-1.09]	death/ICU	14/36	29/55	400,000IU
Nogués (QR)	79% 0.21 [0.10-0.43]	death	21/447	62/391	0.8mg (c)
Lohia	11% 0.89 [0.32-1.89]	death	26 (n)	69 (n)	n/a
Mazziotti	19% 0.81 [0.45-1.47]	death	116 (n)	232 (n)	varies
Elhadi (ICU)	23% 0.77 [0.44-1.32]	death	7/15	274/450	n/a
Alcala-Diaz	81% 0.19 [0.04-0.83]	death	4/79	90/458	0.8mg (c)
Güven (ICU)	25% 0.75 [0.37-1.24]	death	43/113	30/62	300,000IU
Assiri (ICU)	-66% 1.66 [0.25-7.87]	death	12/90	2/28	n/a
Soliman (RCT)	63% 0.37 [0.09-2.78]	death	7/40	3/16	200,000IU
Elamir (RCT)	86% 0.14 [0.01-2.63]	death	0/25	3/25	2.5µg (t)
Yildiz	81% 0.19 [0.04-0.91]	death	1/37	24/170	300,000IU
Maghbooli (DB RCT)	40% 0.60 [0.15-2.38]	death	3/53	5/53	125µg (c)
Leal-Martínez (RCT)	86% 0.14 [0.03-0.80]	death	1/40	7/40	20,000IU
Beigm.. (SB RCT)	89% 0.11 [0.01-1.98]	death	0/30	4/30	600,000IU
Mahmood	30% 0.70 [0.47-1.04]	death	45/238	31/114	varies
Bishop (DB RCT)	34% 0.66 [0.23-1.92]	no recov.	5/65	8/69	1020µg (c)
Cannata-An.. (RCT)	-44% 1.44 [0.76-2.72]	death	22/274	15/269	100,000IU
Zangeneh (ICU)	-26% 1.26 [0.73-2.16]	death	n/a	n/a	n/a
Fiore	93% 0.07 [0.07-0.63]	death	3/58	11/58	200,000IU
Mariani (DB RCT)	-124% 2.24 [0.44-11.3]	death	5/115	2/103	500,000IU
Baykal	22% 0.78 [0.41-1.47]	death	7/18	28/56	300,000IU
Singh (DB RCT)	45% 0.55 [0.31-0.99]	death	11/45	20/45	600,000IU
Shahid	38% 0.62 [0.47-0.82]	death	705 (n)	773 (n)	n/a
Karonova (RCT)	86% 0.14 [0.01-2.66]	ICU	0/56	3/54	50,000IU
Zurita-C.. (SB RCT)	79% 0.21 [0.03-1.59]	death	1/20	6/25	10,000IU
De Niet (DB RCT)	65% 0.35 [0.04-3.10]	death	1/21	3/22	100,000IU
Fairfield	-9% 1.09 [1.04-1.12]	death	population-based cohort	n/a	
Lakkireddy (RCT)	61% 0.39 [0.08-1.91]	death	2/44	5/43	300,000IU
Hafez	94% 0.06 [0.00-1.29]	death	0/7	12/30	150,000IU
Saheb Shari.. (ICU)	36% 0.64 [0.46-0.90]	ICU	20 (n)	25 (n)	50,000IU
Karimpour-Razke..	79% 0.21 [0.10-0.45]	death	10/124	93/329	n/a
Hafezi (ICU)	63% 0.37 [0.14-0.94]	death	8/43	12/37	50,000IU
Bychinin (DB RCT)	27% 0.73 [0.47-1.14]	death	19/52	27/54	80,000IU
Domazet B.. (RCT)	21% 0.79 [0.55-1.13]	death	30/75	39/77	50,000IU
Shamsi	58% 0.42 [0.06-2.95]	death	1/17	23/166	n/a
Mingiano	39% 0.61 [0.38-0.99]	death	13/56	88/232	900µg (c)
Al Sulaiman (ICU)	-22% 1.22 [0.87-1.71]	death	72/144	62/144	n/a
Ogasawara	67% 0.33 [0.01-8.01]	death	0/54	1/54	5µg (p)
Seely (DB RCT)	48% 0.52 [0.10-2.71]	progression	2/42	4/44	55,000IU
Late treatment 43% 0.57 [0.47-0.69]					43% lower risk

Tau² = 0.21, I² = 80.0%, p < 0.0001

	Improvement, RR [CI]	Treatment	Control	Dose (1m)	
Blanch-Rubió	8% 0.92 [0.63-1.36]	cases	62/1,303	47/799	n/a
Sainz-Amo	33% 0.67 [0.27-1.67]	severe case	case control	n/a	
Hernández	-4% 1.04 [0.26-4.10]	death	2/19	20/197	varies
Annweiler	93% 0.07 [0.01-0.61]	death	2/29	10/32	50,000IU
Cereda	-73% 1.73 [0.81-2.74]	death	7/18	40/152	varies
Louca	8% 0.92 [0.88-0.97]	cases	population-based cohort	n/a	




Figure 7. Random effects meta-analysis for peer-reviewed treatment studies. Zeraatkar analyze 356 COVID-19 trials, finding no significant evidence that preprint results are inconsistent with peer-reviewed studies. They also show extremely long peer-review delays, with a median of 6 months to journal publication. A six month delay was equivalent to around 1.5 million deaths during the first two years of the pandemic. Authors recommend using preprint evidence, with appropriate checks for potential falsified data, which provides higher certainty much earlier. Effect extraction is pre-specified, using the most serious outcome reported, see the [appendix](#) for details.

All 102 cholecalciferol COVID-19 studies

c19early.org

October 2023

	Improvement, RR [CI]	Treatment	Control	Dose (5d)	
Annweiler	89% 0.11 [0.03-0.48] death	10/57	5/9	80,000IU	
Annweiler	63% 0.37 [0.06-2.21] death	3/16	10/32	80,000IU	
Burahee	93% 0.07 [0.01-0.54] death	0/12	2/2	400,000IU	
Asimi	97% 0.03 [0.00-0.44] ventilation	0/270	9/86	10,000IU	CT ¹
Sánchez-Zuno (RCT)	89% 0.11 [0.01-0.91] severe case	0/22	4/20	50,000IU	
Efird	49% 0.51 [0.23-1.17] death	11/544	413/15,794	varies	
Valecha	87% 0.13 [0.01-2.43] ICU	0/30	3/25		CT ¹
Khan (RCT)	33% 0.67 [0.37-1.19] no recov.	10/25	15/25	1,800IU	CT ¹
Hunt	47% 0.53 [0.37-0.77] death	43/1,019	1,569/25,489	n/a	
Said (RCT)	42% 0.58 [0.09-3.47] recovery	30 (n)	30 (n)	10,000IU	
Din Ujjan (RCT)	29% 0.71 [0.50-1.03] no recov.	15/25	21/25	1,800IU	CT ¹

Early treatment 60% 0.40 [0.26-0.60]

92/2,050 2,051/41,537

Tau² = 0.21, I² = 62.3%, p < 0.0001

	Improvement, RR [CI]	Treatment	Control	Dose (5d)	
Tan	80% 0.20 [0.04-0.93] oxygen	3/17	16/26	5,000IU	CT ¹
Krishnan	19% 0.81 [0.49-1.34] death	8/16	84/136	n/a	
Rastogi (RCT)	53% 0.47 [0.24-0.92] viral+	6/16	19/24	300,000IU	SHADE
Murai (DB RCT)	-49% 1.49 [0.55-4.05] death	9/119	6/118	200,000IU	
Ling	80% 0.20 [0.08-0.48] death	73 (n)	253 (n)	40,000IU	
Jevalikar	82% 0.18 [0.02-1.69] death	1/128	3/69	60,000IU	
Giannini	37% 0.63 [0.35-1.09] death/ICU	14/36	29/55	400,000IU	
Lohia	11% 0.89 [0.32-1.89] death	26 (n)	69 (n)	n/a	
Mazziotti	19% 0.81 [0.45-1.47] death	116 (n)	232 (n)	varies	
Elhadi (ICU)	23% 0.77 [0.44-1.32] death	7/15	274/450	n/a	ICU patients
Güven (ICU)	25% 0.75 [0.37-1.24] death	43/113	30/62	300,000IU	ICU patients
Assiri (ICU)	-66% 1.66 [0.25-7.87] death	12/90	2/28	n/a	ICU patients
Soliman (RCT)	63% 0.37 [0.09-2.78] death	7/40	3/16	200,000IU	
Yildiz	81% 0.19 [0.04-0.91] death	1/37	24/170	300,000IU	
Leal-Martínez (RCT)	86% 0.14 [0.03-0.80] death	1/40	7/40	20,000IU	CT ¹
Beigm.. (SB RCT)	89% 0.11 [0.01-1.98] death	0/30	4/30	600,000IU	ICU patients CT ¹
Baguma	97% 0.03 [0.00-0.54] death	23 (n)	458 (n)	n/a	
Mahmood	30% 0.70 [0.47-1.04] death	45/238	31/114	varies	
Cannata-An.. (RCT)	-44% 1.44 [0.76-2.72] death	22/274	15/269	100,000IU	COVID-VIT-D
Zangeneh (ICU)	-26% 1.26 [0.73-2.16] death	n/a	n/a	n/a	ICU patients
Fiore	93% 0.07 [0.07-0.63] death	3/58	11/58	200,000IU	
Mariani (DB RCT)	-124% 2.24 [0.44-11.3] death	5/115	2/103	500,000IU	CARED
Baykal	22% 0.78 [0.41-1.47] death	7/18	28/56	300,000IU	Shade-S
Singh (DB RCT)	45% 0.55 [0.31-0.99] death	11/45	20/45	600,000IU	
Shahid	38% 0.62 [0.47-0.82] death	705 (n)	773 (n)	n/a	
Karonova (RCT)	86% 0.14 [0.01-2.66] ICU	0/56	3/54	50,000IU	
Zurita-C.. (SB RCT)	79% 0.21 [0.03-1.59] death	1/20	6/25	10,000IU	
De Niet (DB RCT)	65% 0.35 [0.04-3.10] death	1/21	3/22	100,000IU	
Fairfield	-9% 1.09 [1.04-1.12] death	population-based cohort		n/a	
Lakkireddy (RCT)	61% 0.39 [0.08-1.91] death	2/44	5/43	300,000IU	see notes
Hafez	94% 0.06 [0.00-1.29] death	0/7	12/30	150,000IU	
Saheb Shari.. (ICU)	36% 0.64 [0.46-0.90] ICU	20 (n)	25 (n)	50,000IU	ICU patients
Karimpour-Razke..	79% 0.21 [0.10-0.45] death	10/124	93/329	n/a	
Hafezi (ICU)	63% 0.37 [0.14-0.94] death	8/43	12/37	50,000IU	ICU patients
Bychinin (DB RCT)	27% 0.73 [0.47-1.14] death	19/52	27/54	80,000IU	COVID-VIT
Domazet B.. (RCT)	21% 0.79 [0.55-1.13] death	30/75	39/77	50,000IU	ICU patients
Shamsi	58% 0.42 [0.06-2.95] death	1/17	23/166	n/a	
Al Sulaiman (ICU)	-22% 1.22 [0.87-1.71] death	72/144	62/144	n/a	ICU patients
Seely (DB RCT)	48% 0.52 [0.10-2.71] progression	2/42	4/44	55,000IU	CT ¹

Late treatment 38% 0.62 [0.51-0.75]

351/3,053 897/4,704

Tau² = 0.17, I² = 77.2%, p < 0.0001

	Improvement, RR [CI]	Treatment	Control	Dose (1m)	
Blanch-Rubió	8% 0.92 [0.63-1.36] cases	62/1,303	47/799	n/a	
Sainz-Amo	33% 0.67 [0.27-1.67] severe case	case control		n/a	
Hernández	-4% 1.04 [0.26-4.10] death	2/19	20/197	varies	
Annweiler	93% 0.07 [0.01-0.61] death	2/29	10/32	50,000IU	
Cereda	-73% 1.73 [0.81-2.74] death	7/18	40/152	varies	
Louca	8% 0.92 [0.88-0.97] cases	population-based cohort		n/a	
Cangiano	70% 0.30 [0.10-0.87] death	3/20	39/78	50,000IU	
Vasheghani	30% 0.70 [0.33-1.49] death	7/88	48/420	n/a	
Ma	30% 0.70 [0.50-0.97] cases	49/363	1,329/7,934	n/a	
Sulli	76% 0.24 [0.17-0.36] cases	case control		n/a	
Ullah	-42% 1.42 [0.74-2.37] death	21/64	26/135	n/a	
Maltzer	36% 0.64 [0.29-1.41] cases	6/131	239/3,338	n/a	



Figure 8. Random effects meta-analysis for cholecalciferol treatment studies. Effect extraction is pre-specified, using the most serious outcome reported, see the [appendix](#) for details.

All 14 calcifediol/calcitriol COVID-19 studies

c19early.org

October 2023




Figure 9. Random effects meta-analysis for calcifediol/calcitriol treatment studies. Effect extraction is pre-specified, using the most serious outcome reported, see the [appendix](#) for details.

All 66 vitamin D COVID-19 treatment mortality results

c19early.org

October 2023

	Improvement, RR [CI]	Treatment	Control	Dose (5d)	
Annweiler	89% 0.11 [0.03-0.48]	10/57	5/9	80,000IU	
Annweiler	63% 0.37 [0.06-2.21]	3/16	10/32	80,000IU	
Burahee	93% 0.07 [0.01-0.54]	0/12	2/2	400,000IU	
Efird	49% 0.51 [0.23-1.17]	11/544	413/15,794	varies	
Hunt	47% 0.53 [0.37-0.77]	43/1,019	1,569/25,489	n/a	

Early treatment 68% 0.32 [0.16-0.61] 67/1,648 1,999/41,326 **68% lower risk**

Tau² = 0.34, I² = 73.1%, p = 0.00067

	Improvement, RR [CI]	Treatment	Control	Dose (5d)	
Krishnan	19% 0.81 [0.49-1.34]	8/16	84/136	n/a	
Castillo (RCT)	85% 0.15 [0.01-2.93]	0/50	2/26	0.8mg (c)	COVIDIOL
Murai (DB RCT)	-49% 1.49 [0.55-4.05]	9/119	6/118	200,000IU	
Ling	80% 0.20 [0.08-0.48]	73 (n)	253 (n)	40,000IU	
Jevalikar	82% 0.18 [0.02-1.69]	1/128	3/69	60,000IU	
Nogués (QR)	79% 0.21 [0.10-0.43]	21/447	62/391	0.8mg (c)	
Lohia	11% 0.89 [0.32-1.89]	26 (n)	69 (n)	n/a	
Mazziotti	19% 0.81 [0.45-1.47]	116 (n)	232 (n)	varies	
Elhadi (ICU)	23% 0.77 [0.44-1.32]	7/15	274/450	n/a	ICU patients
Alcala-Diaz	81% 0.19 [0.04-0.83]	4/79	90/458	0.8mg (c)	
Güven (ICU)	25% 0.75 [0.37-1.24]	43/113	30/62	300,000IU	ICU patients
Assiri (ICU)	-66% 1.66 [0.25-7.87]	12/90	2/28	n/a	ICU patients
Soliman (RCT)	63% 0.37 [0.09-2.78]	7/40	3/16	200,000IU	
Elamir (RCT)	86% 0.14 [0.01-2.63]	0/25	3/25	2.5µg (t)	
Yildiz	81% 0.19 [0.04-0.91]	1/37	24/170	300,000IU	
Maghbooli (DB RCT)	40% 0.60 [0.15-2.38]	3/53	5/53	125µg (c)	
Leal-Martínez (RCT)	86% 0.14 [0.03-0.80]	1/40	7/40	20,000IU	CT ¹
Beigm.. (SB RCT)	89% 0.11 [0.01-1.98]	0/30	4/30	600,000IU	ICU patients CT ¹
Baguma	97% 0.03 [0.00-0.54]	23 (n)	458 (n)	n/a	
Mahmood	30% 0.70 [0.47-1.04]	45/238	31/114	varies	
Cannata-An.. (RCT)	-44% 1.44 [0.76-2.72]	22/274	15/269	100,000IU	COVID-VIT-D
Zangeneh (ICU)	-26% 1.26 [0.73-2.16]	n/a	n/a	n/a	ICU patients
Fiore	93% 0.07 [0.07-0.63]	3/58	11/58	200,000IU	
Mariani (DB RCT)	-124% 2.24 [0.44-11.3]	5/115	2/103	500,000IU	CARED
Baykal	22% 0.78 [0.41-1.47]	7/18	28/56	300,000IU	
Singh (DB RCT)	45% 0.55 [0.31-0.99]	11/45	20/45	600,000IU	Shade-S
Shahid	38% 0.62 [0.47-0.82]	705 (n)	773 (n)	n/a	
Zurita-C.. (SB RCT)	79% 0.21 [0.03-1.59]	1/20	6/25	10,000IU	
De Niet (DB RCT)	65% 0.35 [0.04-3.10]	1/21	3/22	100,000IU	
Fairfield	-9% 1.09 [1.04-1.12]	population-based cohort	n/a	n/a	
Lakkireddy (RCT)	61% 0.39 [0.08-1.91]	2/44	5/43	300,000IU	see notes
Hafez	94% 0.06 [0.00-1.29]	0/7	12/30	150,000IU	
Karimpour-Razke..	79% 0.21 [0.10-0.45]	10/124	93/329	n/a	
Hafezi (ICU)	63% 0.37 [0.14-0.94]	8/43	12/37	50,000IU	ICU patients
Bychinin (DB RCT)	27% 0.73 [0.47-1.14]	19/52	27/54	80,000IU	COVID-VIT
Domazet B.. (RCT)	21% 0.79 [0.55-1.13]	30/75	39/77	50,000IU	ICU patients
Shamsi	58% 0.42 [0.06-2.95]	1/17	23/166	n/a	ICU patients
Mingiano	39% 0.61 [0.38-0.99]	13/56	88/232	900µg (c)	
Al Sulaiman (ICU)	-22% 1.22 [0.87-1.71]	72/144	62/144	n/a	ICU patients
Ogasawara	67% 0.33 [0.01-8.01]	0/54	1/54	5µg (p)	

Late treatment 42% 0.58 [0.47-0.71] 367/3,630 1,077/5,715 **42% lower risk**

Tau² = 0.22, I² = 80.6%, p < 0.0001

	Improvement, RR [CI]	Treatment	Control	Dose (1m)	
Hernández	-4% 1.04 [0.26-4.10]	2/19	20/197	varies	
Annweiler	93% 0.07 [0.01-0.61]	2/29	10/32	50,000IU	
Cereda	-73% 1.73 [0.81-2.74]	7/18	40/152	varies	
Cangiano	70% 0.30 [0.10-0.87]	3/20	39/78	50,000IU	
Vasheghani	30% 0.70 [0.33-1.49]	7/88	48/420	n/a	
Ullah	-42% 1.42 [0.74-2.37]	21/64	26/135	n/a	
Oristrell	43% 0.57 [0.41-0.80]	2,296 (n)	3,407 (n)	7.4µg (t)	
Loucera (PSM)	33% 0.67 [0.50-0.91]	374 (n)	374 (n)	varies (c)	
Oristrell	-1% 1.01 [0.93-1.09]	population-based cohort	varies (c)	varies (c)	
Jimenez	50% 0.50 [0.28-0.90]	16/94	65/191	3.7µg (p)	
Pecina	-70% 1.70 [0.36-8.20]	29 (n)	63 (n)	n/a	
Arroyo-Díaz	-12% 1.12 [0.73-1.66]	50/189	167/1,078	n/a	
Ahmed	10% 0.90 [0.72-1.07]	n/a	n/a	n/a	
Mahmood	9% 0.91 [0.60-1.38]	34/138	31/114	varies	
Tylicki	14% 0.86 [0.40-1.38]	28/85	25/48	n/a	
Subramanian	27% 0.73 [0.47-1.09]	31/131	80/336	n/a	
Junior	22% 0.78 [0.30-1.99]	8/113	8/88	n/a	




Figure 10. Random effects meta-analysis for treatment mortality results only.




Figure 11. Random effects meta-analysis for treatment mechanical ventilation results only.

All 26 vitamin D COVID-19 treatment ICU results

c19early.org

October 2023




Figure 12. Random effects meta-analysis for treatment ICU admission results only.

All 23 vitamin D COVID-19 treatment hospitalization results

c19early.org

October 2023




Figure 13. Random effects meta-analysis for treatment hospitalization results only.

All 27 vitamin D COVID-19 treatment case results

c19early.org
October 2023





Figure 14. Random effects meta-analysis for treatment COVID-19 case results only.


All 186 vitamin D COVID-19 sufficiency studies

c19early.org

October 2023



Amend (2020)	27%	0.71 [0.10-0.76]	death	5/17	420
Diaz-Curiel	73%	0.27 [0.07-0.67]	ICU	3/214	91/1,017
Dror	85%	0.15 [0.04-0.44]	severe case	109/120	76/133
Campi	24%	0.76 [0.31-1.83]	death	6/39	13/64
Jude	72%	0.28 [0.25-0.32]	hosp.	n/a	n/a
Al-Jarallah	-88%	1.88 [0.33-6.97]	death	8/120	9/119
Zelzer	46%	0.54 [0.27-1.07]	death	24/121	10/27
Nasiri	-9%	1.09 [0.31-3.83]	death	238 (n)	43 (n)
Bianconi	18%	0.82 [0.41-1.65]	death	94 (n)	106 (n)
González-Estevez	25%	0.75 [0.57-0.98]	symp. case	6/8	32/32
Jimenez	-8%	1.08 [0.59-1.98]	death	50 (n)	110 (n)
Cozier	39%	0.61 [0.39-0.96]	cases	94/1,601	33/373
Al-Salman	44%	0.56 [0.33-0.95]	ICU	113 (n)	337 (n)
Matin	66%	0.34 [0.21-0.56]	cases	case control	
Nimavat	50%	0.50 [0.19-1.27]	death	13/131	5/25
Ribeiro	50%	0.50 [0.28-0.87]	cases	n/a	n/a
Eden (ICU)	64%	0.36 [0.11-1.21]	death	3/26	8/25
Alpcan	73%	0.27 [0.13-0.56]	cases	case control	
Sinaci	79%	0.21 [0.10-0.43]	m/s case	8/100	23/59
di Filippo	11%	0.89 [0.35-2.29]	death	5/28	12/60
Connolly	90%	0.10 [0.01-1.06]	death	65 (n)	49 (n)
Breslin	56%	0.44 [0.22-0.91]	progression	106 (n)	32 (n)
Parra-Ortega	99%	0.01 [0.00-0.20]	death	0/15	63/79
Golabi	90%	0.10 [0.04-0.24]	symp.	34 (n)	10 (n)
Pecina	36%	0.64 [0.04-6.25]	death	6/77	1/15
Karonova	78%	0.22 [0.07-0.67]	death	8/96	10/37
Zafar	-43%	1.43 [0.38-5.39]	death	12/42	2/10
Derakhshanian	45%	0.55 [0.30-0.98]	death	148 (n)	142 (n)
Israel	34%	0.66 [0.54-0.81]	severe case	case control	
Afaghi	55%	0.45 [0.34-0.59]	death	97/537	51/109
Ramirez-Sandoval	32%	0.68 [0.57-0.83]	death	2,337 (n)	571 (n)
Hurst	68%	0.32 [0.13-0.73]	death	68 (n)	191 (n)
Atanasovska	41%	0.59 [0.16-2.23]	death	2/9	9/24
Asghar	53%	0.47 [0.22-0.99]	death	73 (n)	18 (n)
Gönen	66%	0.34 [0.04-3.22]	death	1/80	3/82
Ramos	46%	0.54 [0.25-1.19]	cases	4/9	9/11
Asgari	73%	0.27 [0.09-0.86]	death	n/a	n/a
Seven	47%	0.53 [0.34-0.84]	severe case	n/a	n/a
Ranjbar	42%	0.58 [0.32-1.04]	death	16/163	26/154
Kaur	90%	0.10 [0.04-0.25]	death	5/64	13/17
Fatemi	42%	0.58 [0.30-1.05]	death	18/139	25/109
Ma	67%	0.33 [0.08-1.30]	hosp.	7,893 (n)	7,823 (n)
Putra	26%	0.74 [0.26-2.17]	hosp.	case control	
Seal	45%	0.55 [0.38-0.79]	death	n/a	n/a
Juraj	19%	0.81 [0.64-1.03]	death	127/283	41/74
Saponaro	36%	0.64 [0.25-1.59]	ARDS	5/32	15/61
Subramanian	50%	0.50 [0.27-0.89]	death	16/115	33/118
AlKhafaji	39%	0.61 [0.14-2.17]	death	2/76	13/127
Bushnaq	32%	0.68 [0.37-1.26]	ventilation	10/53	40/144
Junior	84%	0.16 [0.03-0.83]	ventilation	n/a	n/a
Cannata-Andía	-117%	2.17 [0.66-7.17]	death	87 (n)	96 (n)
Sanson	64%	0.36 [0.14-0.91]	death/vent.	2/9	37/60
Zidrou	26%	0.74 [0.15-3.52]	death	2/25	5/46
Rodríguez-Vidales	39%	0.61 [0.22-0.99]	severe case	89/265	27/32
Karonova	22%	0.78 [0.72-0.83]	severe case	n/a	n/a
Pande	93%	0.07 [0.03-0.14]	severe case	7/116	85/93
Ghanei	42%	0.58 [0.31-1.10]	cases	case control	
Ferrer-Sánchez	82%	0.18 [0.01-3.14]	ICU	0/9	4/73
Hafez	98%	0.02 [0.00-0.33]	death	6/116	3/10
Martínez-Rodríguez	52%	0.48 [0.24-0.97]	death	n/a	n/a
Kalichuran	60%	0.40 [0.27-0.60]	symp. case	56 (n)	44 (n)
Voelkle	23%	0.77 [0.28-1.66]	death/ICU	8/34	7/23
Nguyen	81%	0.19 [0.05-0.65]	death	n/a	n/a
Charkowick	73%	0.27 [0.09-0.78]	death	140 (n)	68 (n)
Kazemi	76%	0.24 [0.03-1.93]	death	1/75	7/127
Ozturk	46%	0.54 [0.26-1.09]	severe case	9/110	29/190
Baykal	-8%	1.08 [0.67-1.74]	death	11/20	28/55
Neves	57%	0.43 [0.20-0.91]	death	12/87	9/28
Alzahrani	43%	0.57 [0.17-1.96]	death	179 (n)	78 (n)
Bogliolo	15%	0.85 [0.62-1.16]	death	361 (all patients)	
Charla	11%	0.89 [0.56-1.43]	death	24/91	26/88
Gholi (ICU)	75%	0.25 [0.12-0.56]	death	157 (n)	38 (n)
Doğan	64%	0.36 [0.18-0.72]	cases	case control	






Figure 15. Random effects meta-analysis for sufficiency studies. This plot pools studies with different effects, different vitamin D cutoff levels and measurement times, and studies may be within hospitalized patients, excluding the risk of hospitalization. However, the prevalence of positive effects is notable.

Randomized Controlled Trials (RCTs)

Results restricted to Randomized Controlled Trials (RCTs), after exclusions, and for specific outcomes are shown in Figure 16, 17, 18, 19, and 20.

RCTs have many potential biases. Bias in clinical research may be defined as something that tends to make conclusions differ systematically from the truth. RCTs help to make study groups more similar and can provide a higher level of evidence, however they are subject to many biases ^{Jadad}, and analysis of double-blind RCTs has identified extreme levels of bias ^{Gotzsche}. For COVID-19, the overhead may delay treatment, dramatically compromising efficacy; they may encourage monotherapy for simplicity at the cost of efficacy which may rely on combined or synergistic effects; the participants that sign up may not reflect real world usage or the population that benefits most

in terms of age, comorbidities, severity of illness, or other factors; standard of care may be compromised and unable to evolve quickly based on emerging research for new diseases; errors may be made in randomization and medication delivery; and investigators may have hidden agendas or vested interests influencing design, operation, analysis, and the potential for fraud. All of these biases have been observed with COVID-19 RCTs. There is no guarantee that a specific RCT provides a higher level of evidence.

RCTs for novel acute diseases requiring rapid treatment. High quality RCTs for novel acute diseases are more challenging, with increased ethical issues due to the urgency of treatment, increased risk due to enrollment delays, and more difficult design with a rapidly evolving evidence base. For COVID-19, the most common site of initial infection is the upper respiratory tract. Immediate treatment is likely to be most successful and may prevent or slow progression to other parts of the body. For a non-prophylaxis RCT, it makes sense to provide treatment in advance and instruct patients to use it immediately on symptoms, just as some governments have done by providing medication kits in advance. Unfortunately, no RCTs have been done in this way. Every treatment RCT to date involves delayed treatment. Among the 57 treatments we have analyzed, 64% of RCTs involve very late treatment 5+ days after onset. No non-prophylaxis COVID-19 RCTs match the potential real-world use of early treatments (they may more accurately represent results for treatments that require visiting a medical facility, e.g., those requiring intravenous administration).

RCT bias for widely available treatments. RCTs have a bias against finding an effect for interventions that are widely available — patients that believe they need the intervention are more likely to decline participation and take the intervention. RCTs for vitamin D are more likely to enroll low-risk participants that do not need treatment to recover, making the results less applicable to clinical practice. This bias is likely to be greater for widely known treatments, and may be greater when the risk of a serious outcome is overstated. This bias does not apply to the typical pharmaceutical trial of a new drug that is otherwise unavailable.

Non-RCT studies have been shown to be reliable. Evidence shows that non-RCT trials can also provide reliable results. [Concato](#) find that well-designed observational studies do not systematically overestimate the magnitude of the effects of treatment compared to RCTs. [Anglemyer](#) summarized reviews comparing RCTs to observational studies and found little evidence for significant differences in effect estimates. [Lee](#) shows that only 14% of the guidelines of the Infectious Diseases Society of America were based on RCTs. Evaluation of studies relies on an understanding of the study and potential biases. Limitations in an RCT can outweigh the benefits, for example excessive dosages, excessive treatment delays, or Internet survey bias could have a greater effect on results. Ethical issues may also prevent running RCTs for known effective treatments. For more on issues with RCTs see [Deaton, Nichol](#).

Using all studies identifies efficacy 5.7+ months faster for COVID-19. Currently, 38 of the treatments we analyze show statistically significant efficacy or harm, defined as $\geq 10\%$ decreased risk or $> 0\%$ increased risk from ≥ 3 studies. Of the 38 treatments with statistically significant efficacy/harm, 24 have been confirmed in RCTs, with a mean delay of 5.7 months. For the 14 unconfirmed treatments, 4 have zero RCTs to date. The point estimates for the remaining 10 are all consistent with the overall results (benefit or harm), with 8 showing $> 20\%$. The only treatments showing $> 10\%$ efficacy for all studies, but $< 10\%$ for RCTs are sotrovimab and aspirin.

Summary. We need to evaluate each trial on its own merits. RCTs for a given medication and disease may be more reliable, however they may also be less reliable. For off-patent medications, very high conflict of interest trials may be more likely to be RCTs, and more likely to be large trials that dominate meta analyses.

All 28 vitamin D COVID-19 Randomized Controlled Trials

c19early.org

October 2023




Figure 16. Random effects meta-analysis for Randomized Controlled Trials only. Effect extraction is pre-specified, using the most serious outcome reported, see the [appendix](#) for details.

18 vitamin D COVID-19 Randomized Controlled Trials after exclusions

c19early.org

October 2023



¹ CT: study uses combined treatment

Tau² = 0.00, I² = 0.0%, p = 0.00064

Figure 17. Random effects meta-analysis for RCTs after exclusions. Effect extraction is pre-specified, using the most serious outcome reported, see the [appendix](#) for details.

All 15 vitamin D COVID-19 RCT mortality results

c19early.org

October 2023




Figure 18. Random effects meta-analysis for RCT mortality results.

All 11 vitamin D COVID-19 RCT ICU results

c19early.org

October 2023




Figure 19. Random effects meta-analysis for RCT ICU admission results.

All 8 vitamin D COVID-19 RCT hospitalization results

c19early.org
October 2023




Figure 20. Random effects meta-analysis for RCT hospitalization results.

Exclusions

To avoid bias in the selection of studies, we include all studies in the main analysis, with the exception of *Espitia-Hernandez*. This study uses a combined protocol with another medication that shows high effectiveness when used alone. Authors report on viral clearance, showing 100% clearance with treatment and 0% for the control group. Based on the known mechanisms of action, the combined medication is likely to contribute more to the improvement.

Here we show the results after excluding studies with critical issues.

Murai is a very late stage study (mean 10 days from symptom onset, with 90% on oxygen at baseline), with poorly matched arms in terms of gender, ethnicity, hypertension, diabetes, and baseline ventilation, all of which favor the control group. Further, this study uses cholecalciferol, which may be especially poorly suited for such a late stage. *Cannata-Andía*, *Mariani* are also very late stage studies using cholecalciferol.

The studies excluded are as follows, and the resulting forest plot is shown in Figure 21.

Abdulateef, unadjusted results with no group details.

Al Sulaiman, very late stage study using cholecalciferol instead of calcifediol or calcitriol.

Asimi, excessive unadjusted differences between groups.

Assiri, unadjusted results with no group details.

Aweimer, unadjusted results with no group details.

Baykal, unadjusted results with no group details; significant confounding by time possible due to separation of groups in different time periods.

Beigmohammadi, very late stage study using cholecalciferol instead of calcifediol or calcitriol.

Bychinin, very late stage study using cholecalciferol instead of calcifediol or calcitriol.

Campi, significant unadjusted differences between groups.

Cannata-Andía, very late stage study using cholecalciferol instead of calcifediol or calcitriol.

Din Ujjan, based on dosages and previous research, combined treatments may contribute more to the effect seen.

Domazet Bugarin, very late stage study using cholecalciferol instead of calcifediol or calcitriol.

Elhadi, unadjusted results with no group details.

Fairfield, substantial unadjusted confounding by indication likely.

Guldemir, unadjusted results with no group details.

Güven, very late stage, ICU patients.

Hafezi, very late stage study using cholecalciferol instead of calcifediol or calcitriol.

Holt, significant unadjusted confounding possible.

Junior, unadjusted results with no group details.

Khan, based on dosages and previous research, combined treatments may contribute more to the effect seen.

Krishnan, unadjusted results with no group details.

Leal-Martínez, combined treatments may contribute more to the effect seen.

Lázaro, very few events; unadjusted results with no group details; minimal details provided.

Mahmood, unadjusted results with no group details; substantial unadjusted confounding by indication likely.

Mahmood, unadjusted results with no group details; substantial unadjusted confounding by indication likely.

Mohseni, unadjusted results with no group details.

Murai, very late stage, >50% on oxygen/ventilation at baseline; very late stage study using cholecalciferol instead of calcifediol or calcitriol.

Pecina, unadjusted results with no group details.

Saheb Sharif-Askari, very late stage study using cholecalciferol instead of calcifediol or calcitriol.

Shahid, minimal details provided.

Shamsi, unadjusted results with no group details.

Shehab, unadjusted results with no group details.

Singh, minimal details provided.

Ullah, significant unadjusted confounding possible.

Zangeneh, very late stage study using cholecalciferol instead of calcifediol or calcitriol.

Zurita-Cruz, randomization resulted in significant baseline differences that were not adjusted for.

80 vitamin D COVID-19 treatment studies after exclusions

c19early.org

October 2023

	Improvement, RR [CI]	Treatment	Control	Dose (5d)	
Annweiler	89% 0.11 [0.03-0.48]	death	10/57	5/9	80,000IU
Annweiler	63% 0.37 [0.06-2.21]	death	3/16	10/32	80,000IU
Burahee	93% 0.07 [0.01-0.54]	death	0/12	2/2	400,000IU
Sánchez-Zuno (RCT)	89% 0.11 [0.01-0.91]	severe case	0/22	4/20	50,000IU
Efrid	49% 0.51 [0.23-1.17]	death	11/544	413/15,794	varies
Valecha	87% 0.13 [0.01-2.43]	ICU	0/30	3/25	
Hunt	47% 0.53 [0.37-0.77]	death	43/1,019	1,569/25,489	n/a
Said (RCT)	42% 0.58 [0.09-3.47]	recovery	30 (n)	30 (n)	10,000IU

Early treatment	68%	0.32 [0.18-0.55]	67/1,730	2,006/41,401	68% lower risk
------------------------	------------	-------------------------	-----------------	---------------------	-----------------------

Tau² = 0.28, I² = 57.7%, p < 0.0001

	Improvement, RR [CI]	Treatment	Control	Dose (5d)	
Tan	80% 0.20 [0.04-0.93]	oxygen	3/17	16/26	5,000IU
Castillo (RCT)	85% 0.15 [0.01-2.93]	death	0/50	2/26	0.8mg (c)
Rastogi (RCT)	53% 0.47 [0.24-0.92]	viral+	6/16	19/24	300,000IU
Ling	80% 0.20 [0.08-0.48]	death	73 (n)	253 (n)	40,000IU
Jevalikar	82% 0.18 [0.02-1.69]	death	1/128	3/69	60,000IU
Giannini	37% 0.63 [0.35-1.09]	death/ICU	14/36	29/55	400,000IU
Nogués (QR)	79% 0.21 [0.10-0.43]	death	21/447	62/391	0.8mg (c)
Lohia	11% 0.89 [0.32-1.89]	death	26 (n)	69 (n)	n/a
Mazziotti	19% 0.81 [0.45-1.47]	death	116 (n)	232 (n)	varies
Alcala-Diaz	81% 0.19 [0.04-0.83]	death	4/79	90/458	0.8mg (c)
Soliman (RCT)	63% 0.37 [0.09-2.78]	death	7/40	3/16	200,000IU
Elamir (RCT)	86% 0.14 [0.01-2.63]	death	0/25	3/25	2.5µg (t)
Yildiz	81% 0.19 [0.04-0.91]	death	1/37	24/170	300,000IU
Maghbooli (DB RCT)	40% 0.60 [0.15-2.38]	death	3/53	5/53	125µg (c)
Baguma	97% 0.03 [0.00-0.54]	death	23 (n)	458 (n)	n/a
Bishop (DB RCT)	34% 0.66 [0.23-1.92]	no recov.	5/65	8/69	1020µg (c)
Fiore	93% 0.07 [0.07-0.63]	death	3/58	11/58	200,000IU
Mariani (DB RCT)	-124% 2.24 [0.44-11.3]	death	5/115	2/103	500,000IU
Karonova (RCT)	86% 0.14 [0.01-2.66]	ICU	0/56	3/54	50,000IU
De Niet (DB RCT)	65% 0.35 [0.04-3.10]	death	1/21	3/22	100,000IU
Lakkireddy (RCT)	61% 0.39 [0.08-1.91]	death	2/44	5/43	300,000IU
Hafez	94% 0.06 [0.00-1.29]	death	0/7	12/30	150,000IU
Karimpour-Razke..	79% 0.21 [0.10-0.45]	death	10/124	93/329	n/a
Mingiano	39% 0.61 [0.38-0.99]	death	13/56	88/232	900µg (c)
Ogasawara	67% 0.33 [0.01-8.01]	death	0/54	1/54	5µg (p)
Seely (DB RCT)	48% 0.52 [0.10-2.71]	progression	2/42	4/44	55,000IU

Late treatment	64%	0.36 [0.26-0.49]	101/1,808	486/3,363	64% lower risk
-----------------------	------------	-------------------------	------------------	------------------	-----------------------

Tau² = 0.27, I² = 53.0%, p < 0.0001

	Improvement, RR [CI]	Treatment	Control	Dose (1m)	
Blanch-Rubió	8% 0.92 [0.63-1.36]	cases	62/1,303	47/799	n/a
Sainz-Amo	33% 0.67 [0.27-1.67]	severe case	case control		n/a
Hernández	-4% 1.04 [0.26-4.10]	death	2/19	20/197	varies
Annweiler	93% 0.07 [0.01-0.61]	death	2/29	10/32	50,000IU
Cereda	-73% 1.73 [0.81-2.74]	death	7/18	40/152	varies
Louca	8% 0.92 [0.88-0.97]	cases	population-based cohort		n/a
Cangiano	70% 0.30 [0.10-0.87]	death	3/20	39/78	50,000IU
Vasheghani	30% 0.70 [0.33-1.49]	death	7/88	48/420	n/a
Ma	30% 0.70 [0.50-0.97]	cases	49/363	1,329/7,934	n/a
Sulli	76% 0.24 [0.17-0.36]	cases	case control		n/a
Meltzer	36% 0.64 [0.29-1.41]	cases	6/131	239/3,338	n/a
Ünsal	71% 0.29 [0.11-0.76]	pneumonia	4/28	14/28	varies
Oristrell	43% 0.57 [0.41-0.80]	death	2,296 (n)	3,407 (n)	7.4µg (t)
Loucera (PSM)	33% 0.67 [0.50-0.91]	death	374 (n)	374 (n)	varies (c)
Levitus	31% 0.69 [0.37-1.24]	severe case	65 (n)	64 (n)	varies
Aldwihi	-49% 1.49 [1.13-1.87]	hosp.	94/259	143/479	n/a
Dudley	22% 0.78 [0.23-2.61]	symp. case	15/58	2/6	22,400IU
Fasano	42% 0.58 [0.34-0.99]	cases	13/329	92/1,157	n/a
Oristrell	-1% 1.01 [0.93-1.09]	death	population-based cohort		varies (c)
Jimenez	50% 0.50 [0.28-0.90]	death	16/94	65/191	3.7µg (p)
Israel	13% 0.87 [0.79-0.95]	hosp.	case control		n/a
Sinaci	90% 0.10 [0.01-1.70]	severe case	0/36	7/123	n/a
Golabi	-25% 1.25 [0.86-1.84]	cases	case control		n/a
Bagheri	71% 0.29 [0.10-0.83]	severe case	131 (n)	379 (n)	n/a
Arroyo-Díaz	-12% 1.12 [0.73-1.66]	death	50/189	167/1,078	n/a
Ahmed	49% 0.90 [0.72-1.07]	death	n/a	n/a	n/a
Ma	40% 0.51 [0.29-0.91]	hosp.	26,605 (n)	12,710 (n)	varies
Tylicki	14% 0.86 [0.40-1.38]	death	28/85	25/48	n/a




Figure 21. Random effects meta-analysis excluding studies with significant issues. Effect extraction is pre-specified, using the most serious outcome reported, see the [appendix](#) for details.

Heterogeneity

Heterogeneity in COVID-19 studies arises from many factors including:

Treatment delay. The time between infection or the onset of symptoms and treatment may critically affect how well a treatment works. For example an antiviral may be very effective when used early but may not be effective in late stage disease, and may even be harmful. Oseltamivir, for example, is generally only considered effective for influenza when used within 0-36 or 0-48 hours [McLean, Treanor](#). Baloxavir studies for influenza also show that treatment delay is critical — [Ikematsu](#) report an 86% reduction in cases for post-exposure prophylaxis, [Hayden](#) show a 33 hour reduction in the time to alleviation of symptoms for treatment within 24 hours and a reduction of 13 hours for treatment within 24-48 hours, and [Kumar \(B\)](#) report only 2.5 hours improvement for inpatient treatment.

Treatment delay	Result
Post exposure prophylaxis	86% fewer cases Ikematsu
<24 hours	-33 hours symptoms Hayden
24-48 hours	-13 hours symptoms Hayden
Inpatients	-2.5 hours to improvement Kumar (B)

Table 3. Studies of baloxavir for influenza show that early treatment is more effective.

Figure 22 shows a mixed-effects meta-regression for efficacy as a function of treatment delay in COVID-19 studies from 57 treatments, showing that efficacy declines rapidly with treatment delay. Early treatment is critical for COVID-19.




Figure 22. Early treatment is more effective. Meta-regression showing efficacy as a function of treatment delay in COVID-19 studies from 57 treatments.

Patient demographics. Details of the patient population including age and comorbidities may critically affect how well a treatment works. For example, many COVID-19 studies with relatively young low-comorbidity patients show all patients recovering quickly with or without treatment. In such cases, there is little room for an effective treatment to improve results (as in *López-Medina*).

Effect measured. Efficacy may differ significantly depending on the effect measured, for example a treatment may be very effective at reducing mortality, but less effective at minimizing cases or hospitalization. Or a treatment may have no effect on viral clearance while still being effective at reducing mortality.

Variants. There are many different variants of SARS-CoV-2 and efficacy may depend critically on the distribution of variants encountered by the patients in a study. For example, the Gamma variant shows significantly different characteristics *Faria, Karita, Nonaka, Zavascki*. Different mechanisms of action may be more or less effective depending on variants, for example the viral entry process for the omicron variant has moved towards TMPRSS2-independent fusion, suggesting that TMPRSS2 inhibitors may be less effective *Peacock, Willett*.

Regimen. Effectiveness may depend strongly on the dosage, treatment regimen, and the form of vitamin D used (cholecalciferol, calcifediol, or calcitriol).

Other treatments. The use of other treatments may significantly affect outcomes, including anything from supplements, other medications, or other kinds of treatment such as prone positioning.

Medication quality. The quality of medications may vary significantly between manufacturers and production batches, which may significantly affect efficacy and safety. *Williams* analyze ivermectin from 11 different sources, showing highly variable antiparasitic efficacy across different manufacturers. *Xu* analyze a treatment from two different manufacturers, showing 9 different impurities, with significantly different concentrations for each manufacturer. Non-prescription supplements may show very wide variations in quality *Crawford, Crighton*.

Pooled outcome analysis. We present both pooled analyses and specific outcome analyses. Notably, pooled analysis often results in earlier detection of efficacy as shown in Figure 23. For many COVID-19 treatments, a reduction in mortality logically follows from a reduction in hospitalization, which follows from a reduction in symptomatic cases,

etc. An antiviral tested with a low-risk population may report zero mortality in both arms, however a reduction in severity and improved viral clearance may translate into lower mortality among a high-risk population, and including these results in pooled analysis allows faster detection of efficacy. Trials with high-risk patients may also be restricted due to ethical concerns for treatments that are known or expected to be effective.

Pooled analysis enables using more of the available information. While there is much more information available, for example dose-response relationships, the advantage of the method used here is simplicity and transparency. Note that pooled analysis could hide efficacy, for example a treatment that is beneficial for late stage patients but has no effect on viral replication or early stage disease could show no efficacy in pooled analysis if most studies only examine viral clearance. While we present pooled results, we also present individual outcome analyses, which may be more informative for specific use cases.

Pooled outcomes identify efficacy faster. Currently, 38 of the treatments we analyze show statistically significant efficacy or harm, defined as $\geq 10\%$ decreased risk or $>0\%$ increased risk from ≥ 3 studies. 91% of treatments showing statistically significant efficacy/harm with pooled effects have been confirmed with one or more specific outcomes, with a mean delay of 3.3 months. When restricting to RCTs only, 57% of treatments showing statistically significant efficacy/harm with pooled effects have been confirmed with one or more specific outcomes, with a mean delay of 3.9 months.

Time when COVID-19 studies showed efficacy




Figure 23. The time when studies showed that treatments were effective, defined as statistically significant improvement of $\geq 10\%$ from ≥ 3 studies. Pooled results typically show efficacy earlier than specific outcome results. Results from all studies often shows efficacy much earlier than when restricting to RCTs. Results reflect conditions as used in trials to date, these depend on the population treated, treatment delay, and treatment regimen.

Meta analysis. The distribution of studies will alter the outcome of a meta analysis. Consider a simplified example where everything is equal except for the treatment delay, and effectiveness decreases to zero or below with increasing delay. If there are many studies using very late treatment, the outcome may be negative, even though early treatment is very effective. This may have a greater effect than pooling different outcomes such as mortality and hospitalization. For example a treatment may have 50% efficacy for mortality but only 40% for hospitalization when used within 48 hours. However efficacy could be 0% when used late.

All meta analyses combine heterogeneous studies, varying in population, variants, and potentially all factors above, and therefore may obscure efficacy by including studies where treatment is less effective. Generally, we expect the estimated effect size from meta analysis to be less than that for the optimal case. Looking at all studies is valuable for providing an overview of all research, important to avoid cherry-picking, and informative when a positive result is found despite combining less-optimal situations. However, the resulting estimate does not apply to specific cases such as early treatment in high-risk populations with a specific form and dosage of vitamin D. While we present results for all studies, we also present treatment time and individual outcome analyses, which may be more informative for specific use cases.

Vitamin D studies vary widely in all the factors above, which makes the consistently positive results even more remarkable. A failure to detect an association after combining heterogeneous studies does not mean the treatment is not effective (it may only work in certain cases), however the reverse is not true — an identified association is valid, although the magnitude of the effect may be larger for more optimal cases, and lower for less optimal cases. While we present results for all studies in this paper, the individual outcome, form of vitamin D, and treatment time analyses are more relevant for specific use cases.

Discussion

Sufficiency studies. For sufficiency studies, different studies use different levels as the threshold of sufficiency, vitamin D levels were measured at different times, and some studies measure risk only within hospitalized patients, which excludes the risk of a serious enough case to be hospitalized. However, 173 of 186 studies present positive effects.

Sufficiency studies show a strong correlation between low vitamin D levels and worse COVID-19 outcomes, however they do not provide information on vitamin D treatment. Studies with vitamin D levels measured after admission may show lower levels because COVID-19 infection reduces vitamin D levels. Studies with levels measured before infection also show significant benefit, however the cause could be one or more correlated factors. For example, sunlight exposure increases vitamin D levels, but also increases intracellular melatonin [Zimmerman](#), and melatonin shows significant benefit for COVID-19 [c19early.org](#). Sun exposure is also correlated with physical exercise, which also shows benefit for COVID-19 [c19early.org \(B\)](#).

Treatment studies. 99 of 116 treatment studies report positive effects. Studies vary significantly in terms of treatment delay, treatment regimen, patients characteristics, and (for the pooled effects analysis) outcomes, as reflected in the high degree of heterogeneity. However treatment consistently shows a significant benefit. The treatment studies not showing positive effects are mostly prophylaxis studies with unknown dosages. The only non-prophylaxis studies reporting negative effects are a small unadjusted retrospective [Assiri](#), [Zangeneh](#) with no details of treatment, and [Cannata-Andía](#), [Mariani](#), [Murai](#) which are very late stage studies using cholecalciferol. For [Murai](#), the result also has very low statistical significance due to the small number of events, and the other reported outcomes of ventilation and ICU admission, which have slightly more events and higher confidence, show benefits for vitamin D. Calcifediol or calcitriol, which avoids several days delay in conversion, may be more successful, especially with very late stage usage.

Acute treatment shows higher efficacy than long-term supplementation. Acute treatment shows greater efficacy than chronic prophylaxis for mortality (and in pooled analysis). One hypothesis is that long-term supplementation may affect normal biological processing. A key component of vitamin D processing is regulation via the enzyme CYP24A1, which breaks down active vitamin D. Long-term supplementation may lead to upregulation of CYP24A1, and potentially lower availability of active vitamin D where needed during infection. The prophylaxis RCTs to date [Jolliffe](#), [Villasis-Keever](#) are consistent with this possibility, with the shorter-term supplementation in [Villasis-Keever](#) showing better results compared to the longer-term high adherence daily supplementation in [Jolliffe](#). Specific forms and administration of vitamin D may minimize upregulation of CYP24A1 [Petkovich](#). [Bader](#) performed an RCT showing high-dose cholecalciferol (50,000 IU/week) significantly increased IL-6, however other studies have shown no significant difference in IL-6 [El Hajj](#), [Mousa](#) (30,000IU/wk and 100,000IU bolus + 4,000IU/day).

Other factors may be responsible for the observed lower efficacy in prophylaxis studies. For example, analysis of hospitalized patients is subject to selection bias because long-term accurate-dosage supplementing individuals may be significantly less likely to be hospitalized. Studies spanning higher-UV months are subject to confounding. Note that prophylaxis studies include case results, whereas we may expect vitamin D to be more effective against serious outcomes. Comparison of acute treatment versus long-term supplementation should use the specific outcome analyses rather than the pooled outcome analyses.

Publication bias. Publishing is often biased towards positive results, however evidence suggests that there may be a negative bias for inexpensive treatments for COVID-19. Both negative and positive results are very important for COVID-19, media in many countries prioritizes negative results for inexpensive treatments (inverting the typical incentive for scientists that value media recognition), and there are many reports of difficulty publishing positive results *Boulware, Meeus, Meneguesso*.

One method to evaluate bias is to compare prospective vs. retrospective studies. Prospective studies are more likely to be published regardless of the result, while retrospective studies are more likely to exhibit bias. For example, researchers may perform preliminary analysis with minimal effort and the results may influence their decision to continue. Retrospective studies also provide more opportunities for the specifics of data extraction and adjustments to influence results.

Figure 24 shows a scatter plot of results for prospective and retrospective treatment studies. Prospective studies show 51% [34-63%] improvement in meta analysis, compared to 31% [25-37%] for retrospective studies, suggesting possible negative publication bias, with a non-significant trend towards retrospective studies reporting lower efficacy. This gives us further confidence in the significant efficacy seen in all studies.




Figure 24. Prospective vs. retrospective studies. The diamonds show the results of random effects meta-analysis.

Funnel plot analysis. Funnel plots have traditionally been used for analyzing publication bias. This is invalid for COVID-19 acute treatment trials — the underlying assumptions are invalid, which we can demonstrate with a simple example. Consider a set of hypothetical perfect trials with no bias. Figure 25 plot A shows a funnel plot for a simulation of 80 perfect trials, with random group sizes, and each patient's outcome randomly sampled (10% control event probability, and a 30% effect size for treatment). Analysis shows no asymmetry ($p > 0.05$). In plot B, we add a single typical variation in COVID-19 treatment trials — treatment delay. Consider that efficacy varies from 90% for treatment within 24 hours, reducing to 10% when treatment is delayed 3 days. In plot B, each trial's treatment delay is randomly selected. Analysis now shows highly significant asymmetry, $p < 0.0001$, with six variants of Egger's test all showing $p < 0.05$ *Egger, Harbord, Macaskill, Moreno, Peters, Rothstein, Rücker, Stanley*. Note that these tests fail even though treatment delay is uniformly distributed. In reality treatment delay is more complex — each trial has a different distribution of delays across patients, and the distribution across trials may be biased (e.g., late treatment trials may be more common). Similarly, many other variations in trials may produce asymmetry, including dose, administration, duration of treatment, differences in SOC, comorbidities, age, variants, and bias in design, implementation, analysis, and reporting.




Figure 25. Example funnel plot analysis for simulated perfect trials.

Conflicts of interest. Pharmaceutical drug trials often have conflicts of interest whereby sponsors or trial staff have a financial interest in the outcome being positive. Vitamin D for COVID-19 lacks this because it is an inexpensive and widely available supplement. In contrast, most COVID-19 vitamin D trials have been run by physicians on the front lines with the primary goal of finding the best methods to save human lives and minimize the collateral damage caused by COVID-19. While pharmaceutical companies are careful to run trials under optimal conditions (for example, restricting patients to those most likely to benefit, only including patients that can be treated soon after onset when necessary, and ensuring accurate dosing), not all vitamin D trials represent the optimal conditions for efficacy.

Other meta analyses. Other meta analyses show significant improvements with vitamin D treatment for mortality *Argano, D'Ecclesiis, Hariyanto, Hosseini, Nikniaz, Shah, Xie*, mechanical ventilation *Hariyanto, Meng, Shah, Xie*, ICU admission *Hariyanto, Hosseini, Meng, Shah, Tentolouris, Xie*, hospitalization *Argano*, severity *D'Ecclesiis, Nikniaz, Varikasuvu, Xie*, and cases *Varikasuvu*.

Lakkireddy. The first version of *Lakkireddy* was censored based on incorrect claims from an anti-treatment researcher. For example, the author claims that the gender difference between arms (7/44 vs. 15/43 female) indicates randomization failure, however by simulation, using the group sizes and overall gender ratio, the difference between the number of female patients in each arm is expected to be ≥ 8 6.4% of the time (2.7% with ≥ 8 in the control arm, and 3.7% with ≥ 8 in the treatment arm).

Author claims that the difference in CRP would only happen about one in a billion times. This is incorrect. CRP is not normally distributed, and the observed values could be due to a very small number of outliers with very large CRP in one group.

A response from the study authors can be found at c19early.org (C). The study was republished.

Limitations. Summary statistics from meta analysis necessarily lose information. As with all meta analyses, studies are *heterogeneous*, with differences in treatment delay, treatment regimen, patient demographics, variants, conflicts of interest, standard of care, and *other factors*. We provide analyses by specific outcomes and by treatment delay, and we aim to identify key characteristics in the forest plots and summaries. Results should be viewed in the context of study characteristics.

Some analyses classify treatment based on early or late administration, as done here, while others distinguish between mild, moderate, and severe cases. Viral load does not indicate degree of symptoms — for example patients may have a high viral load while being asymptomatic. With regard to treatments that have antiviral properties, timing of treatment is critical — late administration may be less helpful regardless of severity.

Details of treatment delay per patient is often not available. For example, a study may treat 90% of patients relatively early, but the events driving the outcome may come from 10% of patients treated very late. Our 5 day cutoff for early treatment may be too conservative, [5 days may be too late in many cases](#).

Comparison across treatments is confounded by differences in the studies performed, for example dose, variants, and conflicts of interest. Trials affiliated with special interests may use designs better suited to the preferred outcome.

In some cases, the most serious outcome has very few events, resulting in lower confidence results being used in pooled analysis, however the method is simpler and more transparent. This is less critical as the number of studies increases. Restriction to outcomes with sufficient power may be beneficial in pooled analysis and improve accuracy when there are few studies, however we maintain our pre-specified method to avoid any retrospective changes.

Studies show that combinations of treatments can be highly synergistic and may result in many times greater efficacy than individual treatments alone [Alsaïdi, Andreani, Biancatelli, De Forni, Gasmi, Jeffreys, Jitobaom, Jitobaom \(B\), Ostrov, Thairu](#). Therefore standard of care may be critical and benefits may diminish or disappear if standard of care does not include certain treatments.

This real-time analysis is constantly updated based on submissions. Accuracy benefits from widespread review and submission of updates and corrections from reviewers. Less popular treatments may receive fewer reviews.

No treatment, vaccine, or intervention is 100% available and effective for all current and future variants. Efficacy may vary significantly with different variants and within different populations. All treatments have potential side effects. Propensity to experience side effects may be predicted in advance by qualified physicians. We do not provide medical advice. Before taking any medication, consult a qualified physician who can compare all options, provide personalized advice, and provide details of risks and benefits based on individual medical history and situations.

Physician case series results. Table 4 shows the reported results of physicians that use early treatments for COVID-19, compared to the results for a non-treating physician. The treatments used vary. Physicians typically use a combination of treatments, with almost all reporting use of ivermectin and/or HCQ, and most using additional treatments, including vitamin D. These results are subject to selection and ascertainment bias and more accurate analysis requires details of the patient populations and followup, however results are consistently better across many teams, and consistent with the extensive controlled trial evidence that shows a significant reduction in risk with many early treatments, and improved results with the use of multiple treatments in combination.

LATE TREATMENT						
<i>Physician / Team</i>	<i>Location</i>	<i>Patients</i>	<i>Hospitalization</i>		<i>Mortality</i>	
Dr. David Uip (*)	Brazil	2,200	38.6% (850)	Ref.	2.5% (54)	Ref.
EARLY TREATMENT - 39 physicians/teams						
<i>Physician / Team</i>	<i>Location</i>	<i>Patients</i>	<i>Hospitalization</i>	<i>Improvement</i>	<i>Mortality</i>	<i>Improvement</i>
Dr. Roberto Alfonso Accinelli 0/360 deaths for treatment within 3 days	Peru	1,265			0.6% (7)	77.5%
Dr. Mohammed Tarek Alam patients up to 84 years old	Bangladesh	100			0.0% (0)	100.0%
Dr. Oluwagbenga Alonge	Nigeria	310			0.0% (0)	100.0%
Dr. Raja Bhattacharya up to 88yo, 81% comorbidities	India	148			1.4% (2)	44.9%
Dr. Flavio Cadegiani	Brazil	3,450	0.1% (4)	99.7%	0.0% (0)	100.0%
Dr. Alessandro Capucci	Italy	350	4.6% (16)	88.2%		
Dr. Shankara Chetty	South Africa	8,000			0.0% (0)	100.0%
Dr. Deborah Chisholm	USA	100			0.0% (0)	100.0%
Dr. Ryan Cole	USA	400	0.0% (0)	100.0%	0.0% (0)	100.0%
Dr. Marco Cosentino vs. 3-3.8% mortality during period; earlier treatment better	Italy	392	6.4% (25)	83.5%	0.3% (1)	89.6%
Dr. Jeff Davis	USA	6,000			0.0% (0)	100.0%
Dr. Dhanajay	India	500			0.0% (0)	100.0%
Dr. Bryan Tyson & Dr. George Fareed	USA	20,000	0.0% (6)	99.9%	0.0% (4)	99.2%
Dr. Raphael Furtado	Brazil	170	0.6% (1)	98.5%	0.0% (0)	100.0%
Dr. Heather Gessling	USA	1,500			0.1% (1)	97.3%
Dr. Ellen Guimarães	Brazil	500	1.6% (8)	95.9%	0.4% (2)	83.7%
Dr. Syed Haider	USA	4,000	0.1% (5)	99.7%	0.0% (0)	100.0%
Dr. Mark Hancock	USA	24			0.0% (0)	100.0%
Dr. Sabine Hazan	USA	1,000			0.0% (0)	100.0%
Dr. Mollie James	USA	3,500	1.1% (40)	97.0%	0.0% (1)	98.8%
Dr. Roberta Lacerda	Brazil	550	1.5% (8)	96.2%	0.4% (2)	85.2%
Dr. Katarina Lindley	USA	100	5.0% (5)	87.1%	0.0% (0)	100.0%
Dr. Ben Marble	USA	150,000			0.0% (4)	99.9%
Dr. Edimilson Migowski	Brazil	2,000	0.3% (7)	99.1%	0.1% (2)	95.9%
Dr. Abdulrahman Mohana	Saudi Arabia	2,733			0.0% (0)	100.0%
Dr. Carlos Nigro	Brazil	5,000	0.9% (45)	97.7%	0.5% (23)	81.3%
Dr. Benoit Ochs	Luxembourg	800			0.0% (0)	100.0%
Dr. Ortore	Italy	240	1.2% (3)	96.8%	0.0% (0)	100.0%
Dr. Valerio Pascua one death for a patient presenting on the 5th day in need of supplemental oxygen	Honduras	415	6.3% (26)	83.8%	0.2% (1)	90.2%
Dr. Sebastian Pop	Romania	300			0.0% (0)	100.0%
Dr. Brian Proctor	USA	869	2.3% (20)	94.0%	0.2% (2)	90.6%

Dr. Anastacio Queiroz	Brazil	700			0.0% (0)	100.0%
Dr. Didier Raoult	France	8,315	2.6% (214)	93.3%	0.1% (5)	97.6%
Dr. Karin Ried up to 99yo, 73% comorbidities, av. age 63	Turkey	237			0.4% (1)	82.8%
Dr. Roman Rozencwaig patients up to 86 years old	Canada	80			0.0% (0)	100.0%
Dr. Vipul Shah	India	8,000			0.1% (5)	97.5%
Dr. Silvestre Sobrinho	Brazil	116	8.6% (10)	77.7%	0.0% (0)	100.0%
Dr. Unknown	Brazil	957	1.7% (16)	95.7%	0.2% (2)	91.5%
Dr. Vladimir Zelenko	USA	2,200	0.5% (12)	98.6%	0.1% (2)	96.3%
Mean improvement with early treatment protocols		237,521	Hospitalization	94.1%	Mortality	94.7%

Table 4. Physician results with early treatment protocols compared to no early treatment. (*) Dr. Uip reportedly prescribed early treatment for himself, but not for patients medicospelavidacovid19.com.br.

NIH

NIH provides an analysis of vitamin D for COVID-19 covid19treatmentguidelines.nih.gov, concluding that there is insufficient evidence to recommend for or against use. However, they appear not to have looked at the majority of the evidence. For example, considering RCTs providing clinical results for COVID-19 and vitamin D, they reference only [Elamir](#), [Mariani](#), [Murai](#), [Villasis-Keever](#), and appear not to know about 24 other RCTs [Beigmohammadi](#), [Bishop](#), [Brunvoll](#), [Bychinin](#), [Cannata-Andia](#), [Castillo](#), [De Niet](#), [Din Ujjan](#), [Domazet Bugarin](#), [Hosseini \(C\)](#), [Jolliffe](#), [Karonova](#), [Khan](#), [Lakkireddy](#), [Leal-Martinez](#), [Maghbooli](#), [Rastogi](#), [Said](#), [Seely](#), [Singh](#), [Soliman](#), [Sánchez-Zuno](#), [Wang](#), [Zurita-Cruz](#) as shown in Figure 26. Notably, the NIH selection does not correspond to the most relevant and highest quality studies, for example including [Murai](#), which studies very late treatment (10 days from symptom onset, with 90% on oxygen at baseline) using cholecalciferol. Calcifediol or calcitriol, which avoids several days delay in conversion, may be more appropriate, especially with this very late stage usage. They include none of the early treatment RCTs.

Vitamin D RCTs missing in NIH analysis

c19early.org

October 2023



Figure 26. Analysis by NIH is missing 24 RCTs.

Conclusion

Random effects meta-analysis with pooled effects using the most serious outcome reported shows 60% [40-74%] and 36% [31-42%] lower risk for **early treatment and for all studies**. Results are similar for higher quality studies, peer-reviewed studies, and mortality: early treatment - 68% [45-82%], 57% [36-71%], 68% [39-84%]; all - 37% [31-42%], 40% [34-46%], 36% [27-43%].

116 treatment studies show statistically significant lower risk for **mortality, ICU admission, hospitalization, and cases**. 58 studies from 54 independent teams in 20 countries show statistically significant lower risk.

Acute treatment (early 60% [40-74%], late 44% [32-53%]) shows greater efficacy than **chronic prophylaxis** (31% [23-38%]).

Late stage treatment with **calcifediol or calcitriol** is more effective than **cholecalciferol**: 65% [41-79%] vs. 38% [25-49%].

Other meta analyses show significant improvements with vitamin D treatment for mortality *Argano, D'Ecclesiis, Hariyanto, Hosseini, Nikniaz, Shah, Xie*, mechanical ventilation *Hariyanto, Meng, Shah, Xie*, ICU admission *Hariyanto, Hosseini, Meng, Shah, Tentolouris, Xie*, hospitalization *Argano*, severity *D'Ecclesiis, Nikniaz, Varikasuvu, Xie*, and cases *Varikasuvu*.

Revisions

This paper is data driven, all graphs and numbers are dynamically generated. Please submit updates and corrections at <https://c19early.org/dmeta.html>.

10/2: We added *Bogomaz*.

9/24: We added *Seely (B)*.

9/8: We added *Ogasawara*.

8/29: We added *Shamsi*.

8/24: We added *Al Sulaiman*.

8/21: We added *Mayurathan*.

8/13: We added *Mingiano*.

8/10: We added *Connolly*.

8/5: We added analysis for RCT ICU outcomes.

6/24: We added *Frish*.

6/4: We added *Manojlovic*.

6/4: We added *Jalavu*.

6/4: We added *Wani*.

5/8: We added *Hogarth*.

5/6: We added *Ritsinger*.

5/5: We added *Regalia*.

5/2: We added *Sanamandra*.

4/25: We added *AlKhafaji, Baralić, Hafez*.

4/20: We added *Allami*.

4/19: We added *Cetin Ozbek, Zafar*.

4/16: We added *Rachman*.

4/12: We added *Basińska-Lewandowska*.

4/7: We added *Hermawan, Protas*.

4/6: We added *Bayrak*.

4/5: We added *Aweimer, Khalil, Wang*.

4/2: We added *Gonzalez*.

4/1: We added *Arabadzhiyska*.

3/28: We added *Schmidt*.

3/28: We added *Huang, Nasiri*.

3/23: We added *Davran*.

3/15: We added *Bucurica*.

3/15: We added *Topan*.

3/14: We added *Domazet Bugarin, Siuka*.

3/4: We added *Şengül*.

3/4: We added *Chen*.

3/2: We added *Tan*.

2/18: We added *Ortatatli*.

2/8: We added *Arabi*.

1/28: We added *Batur*.

1/20: We added *Mostafa*.

1/19: We added *Din Ujjan*.

1/17: We added *Valecha*.

1/8: We updated *van Helmond* to the journal version.

1/7/2023: We updated discussion of acute treatment vs. long-term supplementation.

12/31: We added *De Nicolò*.

12/20: We added *Abdrabbo AlYafei*.

12/20: We updated the discussion of heterogeneity and RCTs.

12/12: We added *Vásquez-Procopio*.

12/3: We added *Tallon*.

11/27: We added *Guldemir (B)*.

11/26: We added *Sharif*.

11/13: We added *Gibbons*.

11/8: We added *Said*.

11/4: We added *Bychinin*.

10/28: We added *Álvarez*.

10/26: We added *Hafezi*.

10/15: We added *Charla*.

10/8: We added *Karimpour-Razkenari*.

10/1: We added *Singh*.

9/20: We added *Shahid*.

9/19: We added *van Helmond*.

9/15: We added *Brunvoll*.

9/11: We added *Zeidan*.

8/25: We added *Hafez (B)*.

8/24: We added *Aldwihi, Saheb Sharif-Askari*.

8/23: We added *Doğan*.

8/21: We added *Reyes Pérez*.

8/19: We added *Kalichuran*.

8/16: We updated *Lakkireddy* to the new version (post censorship of the previous version).

8/12: We added *Dana, Zurita-Cruz*.

8/10: We added *Barrett*.

8/5: We added *Bogliolo*.

8/3: We added *Alzahrani*.

7/27: We added *De Niet*.

7/26: We added *Neves*.

7/24: We added *Gholi*.

7/19: We added *Baykal*.

7/2: We added *Hunt*.

6/24: We added *Karonova*.

5/28: We added *Mariani*.

5/25: We added *Kazemi, Zangeneh*.

5/24: We added *Ghanei*.

5/23: We added *Fiore*.

5/20: We added *Hosseini (C)*.

5/19: We added *Jabeen*.

5/19: We added *Ozturk*.

5/8: We added *Charkowick*.

5/5: We added *Nguyen*.

5/1: We added *Khan*.

4/30: We added *Voelkle*.

4/24: We added *Davoudi*.

4/22: We added discussion of *Lakkireddy*.

4/18: We added *Villasis-Keever*.

4/17: We added a section on preclinical research.

4/15: We added *Parant*.

4/12: We added *Martínez-Rodríguez*.

4/5: We added preprint discussion based on *Zeraatkar*.

4/2: We added *Ferrer-Sánchez*.

3/31: We added *Ramos*.

3/27: We added *Pande*.

3/25: We added *Elhadi*.

3/23: We added *Jolliffe*.

3/20: We added *Bushnaq*.

3/19: We added *Shehab*.

3/7: We added *Rodríguez-Vidales*.

3/5: We added *Reis*.

3/4: We added *Nimer*.

3/3: We added *Karonova (B)*.

2/24: We added *Zidrou*.

2/20: We added *Sanson*.

2/19: We added *Cannata-Andía*.

2/18: We added *González-Estevez, Junior*.

2/17: We added *Mahmood*.

2/15: We updated *Vanegas-Cedillo* to the journal version.

2/11: We added *Bychinin (B)*.

2/8: We added *Subramanian*.

2/8: We added *Ranjbar*.

2/7: We added *Tylicki, Ullah*.

2/6: We added *Bishop*.

2/4: We added *Ahmed*.

2/4: We updated *Dror* to the journal version.

1/30: We updated *Leal-Martínez* to the journal version.

1/29: We added *Ansari*.

1/28: We added *Anjum*.

1/25: We added *Saponaro*.

1/23: We added *Juraj*.

1/14: We added *Baguma (B)*.

1/13: We updated *Israel* to the journal version.

1/8: We added *Seal*.

1/5: We added *Pepkowitz*.

1/3/2022: We added *Efird*.

12/26: We added *Abdulateef*.

12/21: We added *Beigmohammadi, Sainz-Amo*.

12/20: We added *Galaznik*.

12/17: We added *Seven*.

12/16: We added *Parra-Ortega*.

12/14: We added *Putra*.

12/9: We added analysis of the number of independent research groups reporting statistically significant positive results.

12/7: We added *Ma*.

12/5: We added *Asgari*.

12/3: We updated *Loucera* to the journal version.

12/3: We added *Fatemi*.

12/3: We added *Kaur*.

11/22: Added discussion related to sufficiency studies.

11/14: We added *Gönen*.

11/12: We added *Asghar*.

11/7: We added *Holt*.

11/3: We added *Atanasovska*.

11/2: We added *Al-Salman, Eden*.

11/1: We updated *Golabi* to the journal version.

10/31: We added *Assiri, Bianconi, Leal-Martínez*.

10/30: We added *Campi, Gaudio*.

10/27: We added *Hurst, Lázaro*.

10/19: We added *Jimenez*.

10/19: We added *Sinaci, Zelzer*.

10/18: We added *Mohseni*.

10/18: We added *Basaran, Dudley*.

10/16: We added a summary plot for all results.

10/15: We added *Ramirez-Sandoval*.

10/15: We added *Maghbooli*.

10/14: We added *Arroyo-Díaz, Burahee* and analysis of treatment mechanical ventilation, ICU admission, and hospitalization results.

9/28: We added *Yildiz*.

9/27: We added *Derakhshanian*.

9/22: We added *Bagheri*.

9/14: We added *Ribeiro*.

9/14: We updated *Vasheghani (B)* to the journal version of the article.

9/14: We added *Elamir*.

9/10: We added *Tomas-Irrigable*.

9/7: We added *Karonova (C)*, *Pecina*.

9/6: We added *Soliman*.

9/1: We added *Golabi*.

8/23: We corrected *Jain (B)* to include the mortality outcome.

8/15: We added *Nimavat*.

8/13: We added *di Filippo (B)* and updated *Louca* to the journal version of the article.

8/12: We added *Alpcan*.

8/10: We added discussion of the immune system and vitamin D.

8/2: We added *Matin*.

8/1: We added *Pimental*.

7/28: We added *Israel (B)*.

7/27: We added *Cozier*.

7/26: We added *Güven*.

7/25: We added *Asimi*.

7/24: We added *Orchard*.

7/21: We added *Savitri*.

7/19: We added *Oristrell*.

7/11: We added *Krishnan*.

6/25: We added *Cereda (B)*.

6/19: We added *Jude*.

6/16: We added *Campi*.

6/12: We added *Levitus*.

6/11: We updated *Oristrell (B)* to the journal version.

6/9: We added *Fasano*.

6/8: We updated *Nogués* to the journal version.

6/7: We added *Diaz-Curiel*, *Dror*.

5/29: We added *Sánchez-Zuno*.

5/22: We added analysis restricted to cholecalciferol studies.

5/21: We added *Alcala-Diaz, Li*.

5/20: We updated *Lakkireddy* to the journal version.

5/19: We added *AlSafar*.

5/10: We added additional information in the abstract.

5/9: We clarified terminology for prophylaxis and added discussion of heterogeneity.

5/8: We added analysis for treatment studies restricted to peer-reviewed articles.

4/30: We added *Loucera*.

4/29: We corrected the treatment group counts for the early treatment group in *Annweiler* (there was no change in the relative risk).

4/24: We added analysis restricted to RCT studies and to calcifediol/calcitriol studies. We have excluded *Espitia-Hernandez* in the treatment analysis because they use a combined protocol with another medication that shows high effectiveness when used alone.

4/14: We added *Blanch-Rubió*.

4/13: We added *Lohia, Oristrell (B)*.

4/12: We added *Barassi*.

4/10: We added *Szeto*.

4/9: We added *Ünsal*.

4/5: We added *Bayramoğlu, Livingston*.

4/4: We added event counts to the forest plots.

3/31: We added *Mendy*.

3/30: We added *Macaya*.

3/29: We added *Im*.

3/28: We added *Freitas*.

3/22: We added *Meltzer*.

3/15: We added *Vanegas-Cedillo*.

3/14: We added *Cereda*.

3/12: We added *Charoenngam*.

3/10: We added *Mazziotti*.

3/6: We added *Ricci*.

2/26: We added *Lakkireddy*.

2/25: We added *Sulli (B)*.

2/20: We added *Gavioli*.

2/20: We added *Infante*.

2/18: *Murai* was updated to the journal version of the paper.

2/17: We corrected an error in the effect extraction for *Angelidi*, and we added treatment case and viral clearance forest plots.

2/16: We added *Susianti*.

2/10: We added *Nogués*.

2/10: We added *Karonova (D)*.

2/9: We added *Karahan*.

2/7: We added *Li (B)*.

2/5: We added *Yilmaz*.

1/31: We added *Demir*.

1/30: We added *Ma (B)*.

1/22: We added *Giannini*.

1/21: We added *Bennouar*.

1/19: We added *Amin*.

1/18: We added *Vasheghani (B)*.

1/16: We moved the analysis with exclusions to the main text, and added additional commentary.

1/15: We added the effect measured for each study in the forest plots.

1/10: We added *Angelidi*.

1/7: We added direct links to the study details in the chronological plots.

1/5: We added direct links to the study details in the forest plots.

1/2/2021: We added dosage information and we added the number of patients to the forest plots.

12/31: We added additional details about the studies in the appendix.

12/28: We added *Jevalikar*.

12/27: We added the total number of authors and patients.

12/23: We added *Cangiano*.

Appendix 1. Methods and Data

We performed ongoing searches of PubMed, medRxiv, ClinicalTrials.gov, The Cochrane Library, Google Scholar, Collabovid, Research Square, ScienceDirect, Oxford University Press, the reference lists of other studies and meta-analyses, and submissions to the site c19early.org/d. Search terms were vitamin D, cholecalciferol, and calcitriol, and COVID-19 or SARS-CoV-2. Automated searches are performed every hour with notification of new matches. All studies that report a result for vitamin D treatment of COVID-19 patients compared to a control group, and all studies comparing COVID-19 outcomes in groups of patients with low and high vitamin D levels are included. A few studies only provide results as a function of change in vitamin D levels, which may not be indicative of results for deficiency/insufficiency versus sufficiency (if levels are already sufficient then further increase may be less useful). This is a living analysis and is updated regularly.

We extracted effect sizes and associated data from all studies. If studies report multiple kinds of effects then the most serious outcome is used in pooled analysis, while other outcomes are included in the outcome specific analyses. For example, if effects for mortality and cases are both reported, the effect for mortality is used, this may be different to the effect that a study focused on. If symptomatic results are reported at multiple times, we used the latest time, for example if mortality results are provided at 14 days and 28 days, the results at 28 days are used. Mortality alone is preferred over combined outcomes. Outcomes with zero events in both arms were not used (the next most serious outcome is used — no studies were excluded). For example, in low-risk populations with no mortality, a reduction in mortality with treatment is not possible, however a reduction in hospitalization, for example, is still valuable. Clinical outcome is considered more important than PCR testing status. When basically all patients recover in both treatment and control groups, preference for viral clearance and recovery is given to results mid-recovery where available (after most or all patients have recovered there is no room for an effective treatment to do better). If only individual symptom data is available, the most serious symptom has priority, for example difficulty breathing or low SpO₂ is more important than cough. When results provide an odds ratio, we computed the relative risk when possible, or converted to a relative risk according to [Zhang](#). Reported confidence intervals and *p*-values were used when available, using adjusted values when provided. If multiple types of adjustments are reported including propensity score matching (PSM), the PSM results are used. Adjusted primary outcome results have preference over unadjusted results for a more serious outcome when the adjustments significantly alter results. When needed, conversion between reported *p*-values and confidence intervals followed [Altman, Altman \(B\)](#), and Fisher's exact test was used to calculate *p*-values for event data. If continuity correction for zero values is required, we use the reciprocal of the opposite arm with the sum of the correction factors equal to 1 [Sweeting](#). Results are expressed with RR < 1.0 favoring treatment, and using the risk of a negative outcome when applicable (for example, the risk of death rather than the risk of survival). If studies only report relative continuous values such as relative times, the ratio of the time for the treatment group versus the time for the control group is used. Calculations are done in Python (3.11.6) with [scipy](#) (1.11.1), [pythonmeta](#) (1.26), [numpy](#) (1.25.0), [statsmodels](#) (0.14.0), and [plotly](#) (5.15.0).

Forest plots are computed using [PythonMeta](#) [Deng](#) with the DerSimonian and Laird random effects model (the fixed effect assumption is not plausible in this case) and inverse variance weighting. Mixed-effects meta-regression results are computed with R (4.1.2) using the [metafor](#) (3.0-2) and [rms](#) (6.2-0) packages, and using the most serious sufficiently powered outcome. Forest plots show simplified dosages for comparison, these are the total dose in the first five days for treatment, and the monthly dose for prophylaxis. Calcifediol, calcitriol, and paricalcitol treatment are indicated with (c), (t), and (p). For full dosage details see below.

We received no funding, this research is done in our spare time. We have no affiliations with any pharmaceutical companies or political parties.

We have classified studies as early treatment if most patients are not already at a severe stage at the time of treatment (for example based on oxygen status or lung involvement), and treatment started within 5 days of the onset of symptoms. If studies contain a mix of early treatment and late treatment patients, we consider the treatment time of patients contributing most to the events (for example, consider a study where most patients are treated early but late treatment patients are included, and all mortality events were observed with late treatment patients).

Analysis of outcomes based on sufficiency

Effect extraction follows pre-specified rules as detailed above and gives priority to more serious outcomes. Only the first (most serious) outcome is used in pooled analysis, which may differ from the effect a paper focuses on. Other outcomes are used in outcome specific analyses.

<i>Abdollahi</i> , 12/12/2020, retrospective, Iran, peer-reviewed, 7 authors.	risk of case, 53.9% lower, RR 0.46, $p = 0.001$, high D levels 108, low D levels 294, >30ng/ml.
<i>Abdrabbo AlYafei</i> , 12/5/2022, retrospective, Qatar, peer-reviewed, mean age 19.0, 5 authors.	risk of case, 23.2% lower, OR 0.77, $p < 0.001$, cutoff 10ng/mL, adjusted per study, inverted to make OR<1 favor high D levels ($\geq 10\text{ng/mL}$), case control OR, severe deficiency vs. optimal, multivariable.
	risk of case, 21.5% lower, OR 0.78, $p < 0.001$, cutoff 20ng/mL, adjusted per study, inverted to make OR<1 favor high D levels ($\geq 20\text{ng/mL}$), case control OR, mild/moderate deficiency vs. optimal, multivariable.
<i>Abdulrahman</i> , 4/17/2023, retrospective, United Kingdom, peer-reviewed, mean age 69.0, 7 authors, study period April 2020 - May 2021.	risk of death, 90.1% lower, OR 0.10, $p = 0.048$, high D levels ($\geq 25\text{nmol/L}$) 76, low D levels ($< 25\text{nmol/L}$) 5, adjusted per study, inverted to make OR<1 favor high D levels ($\geq 25\text{nmol/L}$), multivariable, RR approximated with OR.
	risk of progression, 82.5% lower, OR 0.18, $p = 0.09$, high D levels ($\geq 25\text{nmol/L}$) 76, low D levels ($< 25\text{nmol/L}$) 5, adjusted per study, inverted to make OR<1 favor high D levels ($\geq 25\text{nmol/L}$), hospitalization, ICU, or death, multivariable, RR approximated with OR.
<i>Abrishami</i> , 10/30/2020, retrospective, Iran, peer-reviewed, mean age 55.2, 7 authors.	risk of death, 75.9% lower, RR 0.24, $p = 0.04$, high D levels ($\geq 25\text{ng/mL}$) 3 of 47 (6.4%), low D levels ($< 25\text{ng/mL}$) 9 of 26 (34.6%), NNT 3.5, adjusted per study, inverted to make RR<1 favor high D levels ($\geq 25\text{ng/mL}$), Cox model 2.
<i>Afaghi</i> , 10/12/2021, retrospective, Iran, peer-reviewed, 7 authors.	risk of death, 55.0% lower, RR 0.45, $p = 0.002$, high D levels 97 of 537 (18.1%), low D levels 51 of 109 (46.8%), NNT 3.5, adjusted per study, inverted to make RR<1 favor high D levels, odds ratio converted to relative risk, >20ng/mL, multivariate.
	risk of mechanical ventilation, 55.9% lower, RR 0.44, $p < 0.001$, high D levels 89 of 537 (16.6%), low D levels 41 of 109 (37.6%), NNT 4.8, >20ng/mL, unadjusted.
	risk of ICU admission, 34.1% lower, RR 0.66, $p < 0.001$, high D levels 211 of 537 (39.3%), low D levels 65 of 109 (59.6%), NNT 4.9, >20ng/mL, unadjusted.
<i>Al-Salman</i> , 7/29/2021, retrospective, Bahrain, peer-reviewed, 5 authors.	risk of ICU admission, 44.4% lower, OR 0.56, $p = 0.03$, high D levels ($\geq 50\text{nmol/L}$) 113, low D levels ($< 50\text{nmol/L}$) 337, inverted to make OR<1 favor high D levels ($\geq 50\text{nmol/L}$), multinomial regression, RR approximated with OR.

<i>Alguwaihies</i> , 12/5/2020, retrospective, Saudi Arabia, peer-reviewed, 10 authors.	risk of death, 85.7% lower, RR 0.14, $p = 0.007$, high D levels 111, low D levels 328, inverted to make $RR < 1$ favor high D levels, > 12.5 nmol/L.
<i>AlKhafaji</i> , 1/31/2022, retrospective, Saudi Arabia, peer-reviewed, mean age 56.8, 16 authors, study period January 2021 - August 2021.	risk of death, 38.6% lower, RR 0.61, $p = 0.50$, high D levels (≥ 20 ng/mL) 2 of 76 (2.6%), low D levels (< 20 ng/mL) 13 of 127 (10.2%), inverted to make $RR < 1$ favor high D levels (≥ 20 ng/mL), odds ratio converted to relative risk.
	risk of mechanical ventilation, 31.0% lower, RR 0.69, $p = 0.51$, high D levels (≥ 20 ng/mL) 2 of 76 (2.6%), low D levels (< 20 ng/mL) 13 of 127 (10.2%), inverted to make $RR < 1$ favor high D levels (≥ 20 ng/mL), odds ratio converted to relative risk.
	risk of ICU admission, 41.8% lower, RR 0.58, $p = 0.20$, high D levels (≥ 20 ng/mL) 2 of 76 (2.6%), low D levels (< 20 ng/mL) 13 of 127 (10.2%), inverted to make $RR < 1$ favor high D levels (≥ 20 ng/mL), odds ratio converted to relative risk.
<i>Allami</i> , 11/8/2022, retrospective, Iraq, peer-reviewed, 6 authors.	risk of hospitalization, 92.5% lower, OR 0.07, $p < 0.001$, high D levels (≥ 10 ng/mL) 91, low D levels (< 10 ng/mL) 80, adjusted per study, inverted to make $OR < 1$ favor high D levels (≥ 10 ng/mL), case control OR, multivariable.
<i>Alpcan</i> , 8/10/2021, retrospective, Turkey, peer-reviewed, 3 authors.	risk of case, 73.0% lower, OR 0.27, $p < 0.001$, high D levels 42 of 75 (56.0%) cases, 66 of 80 (82.5%) controls, NNT 3.2, case control OR, > 20 ng/mL.
<i>AlSafar</i> , 5/19/2021, retrospective, United Arab Emirates, peer-reviewed, 8 authors.	risk of death, 59.3% lower, RR 0.41, $p = 0.048$, high D levels 16 of 337 (4.7%), low D levels 10 of 127 (7.9%), adjusted per study, inverted to make $RR < 1$ favor high D levels, odds ratio converted to relative risk, ≥ 12 ng/mL.
	risk of severe case, 33.2% lower, RR 0.67, $p = 0.005$, high D levels 337, low D levels 127, adjusted per study, inverted to make $RR < 1$ favor high D levels, odds ratio converted to relative risk, ≥ 12 ng/mL.
<i>Alzahrani</i> , 6/23/2022, retrospective, Saudi Arabia, peer-reviewed, mean age 54.3, 9 authors, study period March 2020 - July 2021.	risk of death, 42.5% lower, OR 0.57, $p = 0.46$, high D levels (≥ 25 ng/mL) 179, low D levels (< 25 ng/mL) 78, adjusted per study, inverted to make $OR < 1$ favor high D levels (≥ 25 ng/mL), multivariable, RR approximated with OR.
	risk of ICU admission, 7.4% lower, OR 0.93, $p = 0.80$, high D levels (≥ 25 ng/mL) 179, low D levels (< 25 ng/mL) 78, adjusted per study, inverted to make $OR < 1$ favor high D levels (≥ 25 ng/mL), multivariable, RR approximated with OR.
<i>Al-Jarallah</i> , 6/20/2021, retrospective, Kuwait, peer-reviewed, 20 authors.	risk of death, 88.3% higher, RR 1.88, $p = 0.45$, high D levels 8 of 120 (6.7%), low D levels 9 of 119 (7.6%), odds ratio converted to relative risk.
<i>Amin</i> , 1/7/2021, retrospective, population-based cohort, United Kingdom, peer-reviewed, 2 authors.	COVID-19 severity, 32.3% higher, RR 1.32, $p = 0.20$, high D levels 140,898, low D levels 35,079, inverted to make $RR < 1$ favor high D levels, odds ratio converted to relative risk, ≥ 50 nmol/L vs. < 25 nmol/L, MR Egger, baseline risk approximated with overall risk.

	<p>risk of case, 7.6% higher, RR 1.08, $p = 0.14$, high D levels 140,898, low D levels 35,079, inverted to make $RR < 1$ favor high D levels, odds ratio converted to relative risk, ≥ 50 nmol/L vs. < 25 nmol/L, MR Egger, baseline risk approximated with overall risk.</p>
<p><i>Angelidi</i>, 1/9/2021, retrospective, USA, peer-reviewed, 8 authors.</p>	<p>risk of death, 88.0% lower, RR 0.12, $p = 0.01$, high D levels 6 of 65 (9.2%), low D levels 20 of 79 (25.3%), NNT 6.2, adjusted per study, > 30 ng/mL, supplementary table 2, multivariable logistic regression model 5.</p>
<p><i>Anjum</i>, 7/31/2020, prospective, Pakistan, peer-reviewed, 6 authors, study period March 2020 - June 2020, excluded in exclusion analyses: unadjusted results with no group details.</p>	<p>risk of death, 62.5% lower, RR 0.38, $p = 0.02$, high D levels (≥ 25 nmol/L) 8 of 80 (10.0%), low D levels (< 25 nmol/L) 16 of 60 (26.7%), NNT 6.0.</p>
<p><i>Ansari</i>, 12/31/2020, prospective, Pakistan, peer-reviewed, 6 authors, study period 1 March, 2020 - 31 August, 2020, excluded in exclusion analyses: unadjusted results with no group details.</p>	<p>risk of death, 86.0% lower, RR 0.14, $p = 0.02$, high D levels (≥ 25 nmol/L) 2 of 68 (2.9%), low D levels (< 25 nmol/L) 12 of 57 (21.1%), NNT 5.5.</p>
<p><i>Arabadzhiyska</i>, 2/28/2023, retrospective, Bulgaria, peer-reviewed, mean age 53.7, 2 authors, study period October 2021 - December 2021.</p>	<p>risk of severe case, 29.8% lower, RR 0.70, $p = 0.16$, high D levels (≥ 20 ng/mL) 16 of 44 (36.4%), low D levels (< 20 ng/mL) 29 of 56 (51.8%), NNT 6.5.</p>
<p><i>Arabi</i>, 1/22/2023, retrospective, Iran, peer-reviewed, 7 authors.</p>	<p>risk of death, 40.0% lower, RR 0.60, $p = 0.28$, high D levels (≥ 20 ng/mL) 6 of 30 (20.0%), low D levels (< 20 ng/mL) 13 of 39 (33.3%), NNT 7.5.</p>
	<p>risk of ICU admission, 39.3% lower, RR 0.61, $p = 0.20$, high D levels (≥ 20 ng/mL) 7 of 30 (23.3%), low D levels (< 20 ng/mL) 15 of 39 (38.5%), NNT 6.6.</p>
	<p>risk of AKI, 42.2% lower, RR 0.58, $p = 0.13$, high D levels (≥ 20 ng/mL) 8 of 30 (26.7%), low D levels (< 20 ng/mL) 18 of 39 (46.2%), NNT 5.1.</p>
<p><i>Asgari</i>, 11/21/2021, retrospective, Iran, peer-reviewed, 6 authors, study period 21 May, 2020 - 4 September, 2020.</p>	<p>risk of death, 72.5% lower, OR 0.27, $p = 0.03$, cutoff 25 ng/mL, adjusted per study, inverted to make $OR < 1$ favor high D levels (≥ 25 ng/mL), RR approximated with OR.</p>
	<p>risk of progression, 65.6% lower, OR 0.34, $p = 0.02$, cutoff 25 ng/mL, adjusted per study, inverted to make $OR < 1$ favor high D levels (≥ 25 ng/mL), RR approximated with OR.</p>
<p><i>Asghar</i>, 11/10/2021, retrospective, Pakistan, peer-reviewed, 8 authors.</p>	<p>risk of death, 53.1% lower, HR 0.47, $p = 0.046$, high D levels (≥ 10 ng/mL) 73, low D levels (< 10 ng/mL) 18, inverted to make $HR < 1$ favor high D levels (≥ 10 ng/mL), multivariate Cox regression.</p>
	<p>risk of mechanical ventilation, 19.4% lower, HR 0.81, $p = 0.32$, high D levels (≥ 10 ng/mL) 5 of 73 (6.8%), low D levels (< 10 ng/mL) 6 of 18 (33.3%), NNT 3.8, adjusted per study, inverted to make $HR < 1$ favor high D levels (≥ 10 ng/mL), multivariate Cox regression.</p>

	<p>risk of ICU admission, 32.9% lower, HR 0.67, $p = 0.54$, high D levels ($\geq 10\text{ng/mL}$) 73, low D levels ($< 10\text{ng/mL}$) 18, inverted to make $\text{HR} < 1$ favor high D levels ($\geq 10\text{ng/mL}$), multivariate Cox regression.</p>
<p><i>Atanasovska</i>, 11/2/2021, retrospective, North Macedonia, peer-reviewed, 8 authors.</p>	<p>risk of death, 40.7% lower, RR 0.59, $p = 0.68$, high D levels ($\geq 30\text{ng/mL}$) 2 of 9 (22.2%), low D levels ($< 30\text{ng/mL}$) 9 of 24 (37.5%), NNT 6.5.</p>
	<p>risk of severe case, 59.0% lower, RR 0.41, $p = 0.13$, high D levels ($\geq 30\text{ng/mL}$) 2 of 9 (22.2%), low D levels ($< 30\text{ng/mL}$) 13 of 24 (54.2%), NNT 3.1.</p>
<p><i>Baktash</i>, 8/27/2020, prospective, United Kingdom, peer-reviewed, 8 authors.</p>	<p>risk of death, 28.6% lower, RR 0.71, $p = 0.50$, high D levels 4 of 31 (12.9%), low D levels 6 of 39 (15.4%), adjusted per study, inverted to make $\text{RR} < 1$ favor high D levels, $> 30\text{nmol/L}$.</p>
<p><i>Barassi</i>, 1/25/2021, retrospective, Italy, peer-reviewed, 8 authors.</p>	<p>risk of death, 64.9% lower, RR 0.35, $p = 0.44$, high D levels 1 of 31 (3.2%), low D levels 8 of 87 (9.2%), NNT 17, $> 20\text{ng/mL}$.</p>
	<p>risk of mechanical ventilation, 64.9% lower, RR 0.35, $p = 0.15$, high D levels 2 of 31 (6.5%), low D levels 16 of 87 (18.4%), NNT 8.4, $> 20\text{ng/mL}$.</p>
<p><i>Barrett</i>, 8/9/2022, prospective, Ireland, peer-reviewed, mean age 56.0, 19 authors, study period March 2020 - April 2021.</p>	<p>risk of death, 78.4% lower, OR 0.22, $p = 0.006$, high D levels ($\geq 30\text{nmol/L}$) 144, low D levels ($< 30\text{nmol/L}$) 88, adjusted per study, inverted to make $\text{OR} < 1$ favor high D levels ($\geq 30\text{nmol/L}$), multivariable, RR approximated with OR.</p>
	<p>risk of ICU admission, 15.3% lower, OR 0.85, $p = 0.63$, high D levels ($\geq 30\text{nmol/L}$) 144, low D levels ($< 30\text{nmol/L}$) 88, adjusted per study, inverted to make $\text{OR} < 1$ favor high D levels ($\geq 30\text{nmol/L}$), multivariable, RR approximated with OR.</p>
	<p>risk of progression, 52.6% lower, OR 0.47, $p = 0.12$, high D levels ($\geq 30\text{nmol/L}$) 144, low D levels ($< 30\text{nmol/L}$) 88, adjusted per study, inverted to make $\text{OR} < 1$ favor high D levels ($\geq 30\text{nmol/L}$), extended oxygen requirement, multivariable, RR approximated with OR.</p>
<p><i>Basaran</i>, 2/12/2021, retrospective, Turkey, peer-reviewed, 6 authors.</p>	<p>risk of severe case, 68.6% lower, RR 0.31, $p = 0.005$, high D levels 82 of 119 (68.9%), low D levels 80 of 85 (94.1%), NNT 4.0, inverted to make $\text{RR} < 1$ favor high D levels, odds ratio converted to relative risk, $> 10\mu\text{g/L}$, per standard deviation increase in levels.</p>
<p><i>Basińska-Lewandowska</i>, 3/24/2023, retrospective, Poland, peer-reviewed, 5 authors, excluded in exclusion analyses: unadjusted results with no group details.</p>	<p>risk of case, 58.3% lower, RR 0.42, $p = 0.02$, high D levels ($\geq 12\text{ng/mL}$) 20 of 109 (18.3%), low D levels ($< 12\text{ng/mL}$) 11 of 25 (44.0%), NNT 3.9.</p>
<p><i>Batur</i>, 12/26/2022, retrospective, Turkey, peer-reviewed, 2 authors, study period March 2020 - June 2021, excluded in exclusion analyses: unadjusted differences between groups.</p>	<p>risk of death, 71.9% lower, RR 0.28, $p < 0.001$, high D levels ($\geq 20\text{ng/mL}$) 17 of 76 (22.4%), low D levels ($< 20\text{ng/mL}$) 94 of 118 (79.7%), NNT 1.7.</p>

	secondary infection, 23.3% lower, RR 0.77, $p = 0.03$, high D levels ($\geq 20\text{ng/mL}$) 40 of 76 (52.6%), low D levels ($< 20\text{ng/mL}$) 81 of 118 (68.6%), NNT 6.2, growth in culture.
<i>Baykal</i> , 5/30/2022, retrospective, Turkey, peer-reviewed, 2 authors, study period 1 April, 2020 - 1 March, 2021, dosage 300,000IU single dose.	risk of death, 8.0% higher, RR 1.08, $p = 0.80$, high D levels ($\geq 20\text{ng/mL}$) 11 of 20 (55.0%), low D levels ($< 20\text{ng/mL}$) 28 of 55 (50.9%), outcome based on serum levels.
	risk of ICU admission, 4.8% lower, RR 0.95, $p = 1.00$, high D levels ($\geq 20\text{ng/mL}$) 9 of 20 (45.0%), low D levels ($< 20\text{ng/mL}$) 26 of 55 (47.3%), NNT 44, outcome based on serum levels.
	risk of progression, 6.1% lower, RR 0.94, $p = 0.77$, high D levels ($\geq 20\text{ng/mL}$) 14 of 20 (70.0%), low D levels ($< 20\text{ng/mL}$) 41 of 55 (74.5%), NNT 22, severe/critical, outcome based on serum levels.
<i>Bayrak</i> , 4/5/2023, retrospective, Turkey, peer-reviewed, mean age 19.0, 8 authors, study period November 2020 - January 2021.	risk of moderate/severe case, 26.5% lower, RR 0.73, $p = 1.00$, high D levels ($\geq 20\text{ng/mL}$) 3 of 49 (6.1%), low D levels ($< 20\text{ng/mL}$) 2 of 24 (8.3%), NNT 45.
	risk of case, 33.4% lower, OR 0.67, $p = 0.23$, high D levels ($\geq 20\text{ng/mL}$) 41 of 73 (56.2%) cases, 50 of 76 (65.8%) controls, NNT 9.9, case control OR.
<i>Bayramoğlu</i> , 3/31/2021, retrospective, Turkey, peer-reviewed, 7 authors.	risk of moderate/severe case, 69.5% lower, RR 0.30, $p = 0.03$, high D levels 10 of 60 (16.7%), low D levels 24 of 43 (55.8%), NNT 2.6, adjusted per study, inverted to make $\text{RR} < 1$ favor high D levels, odds ratio converted to relative risk, $> 12 \text{ ng/mL}$, multivariate logistic regression.
<i>Bennouar</i> , 1/12/2021, prospective, Algeria, peer-reviewed, 4 authors.	risk of death, 85.5% lower, RR 0.14, $p = 0.002$, high D levels 4 of 30 (13.3%), low D levels 15 of 32 (46.9%), NNT 3.0, adjusted per study, inverted to make $\text{RR} < 1$ favor high D levels, $> 30 \mu\text{g/L}$ vs. $< 10 \mu\text{g/L}$, proportional Cox regression.
	risk of death, 63.0% lower, RR 0.37, $p = 0.10$, high D levels 4 of 30 (13.3%), low D levels 14 of 35 (40.0%), NNT 3.7, adjusted per study, inverted to make $\text{RR} < 1$ favor high D levels, $> 30 \mu\text{g/L}$ vs. $10\text{-}19 \mu\text{g/L}$, proportional Cox regression.
	risk of death, 23.1% lower, RR 0.77, $p = 0.73$, high D levels 4 of 30 (13.3%), low D levels 4 of 23 (17.4%), NNT 25, adjusted per study, inverted to make $\text{RR} < 1$ favor high D levels, $> 30 \mu\text{g/L}$ vs. $20\text{-}29 \mu\text{g/L}$, proportional Cox regression.
<i>Bianconi</i> , 7/1/2021, prospective, Italy, peer-reviewed, 12 authors.	risk of death, 17.5% lower, HR 0.82, $p = 0.58$, high D levels ($\geq 12\text{ng/ml}$) 94, low D levels ($< 12\text{ng/ml}$) 106, model 3, Table S2, Cox proportional hazards.
	risk of death, 13.9% lower, HR 0.86, $p = 0.73$, high D levels ($\geq 20\text{ng/ml}$) 40, low D levels ($< 20\text{ng/ml}$) 160, model 3, Table S2, Cox proportional hazards.
	risk of death/ICU, 15.9% lower, HR 0.84, $p = 0.53$, high D levels ($\geq 12\text{ng/ml}$) 94, low D levels ($< 12\text{ng/ml}$) 106, model 3, Cox

	proportional hazards.
	risk of death/ICU, 10.9% lower, HR 0.89, $p = 0.73$, high D levels ($\geq 20\text{ng/ml}$) 40, low D levels ($< 20\text{ng/ml}$) 160, model 3, Cox proportional hazards.
<i>Bogliolo</i> , 7/5/2022, prospective, Italy, peer-reviewed, median age 73.0, 16 authors, study period March 2020 - August 2020.	risk of death, 15.3% lower, HR 0.85, $p = 0.29$, cutoff 20ng/mL, inverted to make $\text{HR} < 1$ favor high D levels ($\geq 20\text{ng/mL}$).
<i>Bogomaz</i> , 8/24/2023, retrospective, Ukraine, peer-reviewed, median age 62.0, 2 authors.	risk of death, 70.0% lower, RR 0.30, $p = 0.24$, high D levels ($\geq 30\text{ng/ml}$) 1 of 28 (3.6%), low D levels ($< 30\text{ng/ml}$) 5 of 42 (11.9%), NNT 12, inverted to make $\text{RR} < 1$ favor high D levels ($\geq 30\text{ng/ml}$), odds ratio converted to relative risk.
	risk of mechanical ventilation, 75.0% lower, RR 0.25, $p = 0.23$, high D levels ($\geq 30\text{ng/ml}$) 1 of 28 (3.6%), low D levels ($< 30\text{ng/ml}$) 6 of 42 (14.3%), NNT 9.3.
	risk of progression, 62.5% lower, RR 0.38, $p = 0.30$, high D levels ($\geq 30\text{ng/ml}$) 2 of 28 (7.1%), low D levels ($< 30\text{ng/ml}$) 8 of 42 (19.0%), NNT 8.4, critical case.
	risk of oxygen therapy, 27.0% lower, RR 0.73, $p = 0.24$, high D levels ($\geq 30\text{ng/ml}$) 10 of 28 (35.7%), low D levels ($< 30\text{ng/ml}$) 28 of 42 (66.7%), NNT 3.2, adjusted per study, inverted to make $\text{RR} < 1$ favor high D levels ($\geq 30\text{ng/ml}$), odds ratio converted to relative risk, multivariable.
<i>Breslin</i> , 8/17/2021, retrospective, Ireland, peer-reviewed, 4 authors.	risk of progression, 55.6% lower, OR 0.44, $p = 0.03$, high D levels ($\geq 30\text{nmol/l}$) 106, low D levels ($< 30\text{nmol/l}$) 32, adjusted per study, inverted to make $\text{OR} < 1$ favor high D levels ($\geq 30\text{nmol/l}$), infiltrates on chest X-ray, multivariable, RR approximated with OR.
<i>Bucurica</i> , 3/6/2023, retrospective, Romania, peer-reviewed, mean age 55.2, 9 authors, study period 1 June, 2020 - 31 May, 2022.	risk of case, 27.6% lower, OR 0.72, $p < 0.001$, high D levels ($\geq 20\text{ng/mL}$) 7,958, low D levels ($< 20\text{ng/mL}$) 3,224, inverted to make $\text{OR} < 1$ favor high D levels ($\geq 20\text{ng/mL}$), RR approximated with OR.
	risk of case, 7.4% higher, OR 1.07, $p = 0.19$, high D levels ($\geq 30\text{ng/mL}$) 4,367, low D levels ($< 30\text{ng/mL}$) 6,815, inverted to make $\text{OR} < 1$ favor high D levels ($\geq 30\text{ng/mL}$), RR approximated with OR.
<i>Bushnaq</i> , 2/8/2022, retrospective, Saudi Arabia, peer-reviewed, 7 authors, excluded in exclusion analyses: unadjusted results with no group details.	risk of mechanical ventilation, 32.1% lower, RR 0.68, $p = 0.27$, high D levels ($\geq 20\text{ng/mL}$) 10 of 53 (18.9%), low D levels ($< 20\text{ng/mL}$) 40 of 144 (27.8%), NNT 11, unadjusted.
	risk of ICU admission, 3.9% lower, RR 0.96, $p = 0.87$, high D levels ($\geq 20\text{ng/mL}$) 23 of 53 (43.4%), low D levels ($< 20\text{ng/mL}$) 65 of 144 (45.1%), NNT 57, unadjusted.
<i>Bychinin (B)</i> , 5/7/2021, retrospective, Russia, peer-reviewed, 5 authors, excluded in exclusion analyses: excessive unadjusted differences between groups.	risk of death, 36.2% lower, RR 0.64, $p = 0.03$, high D levels ($\geq 10\text{ng/mL}$) 16 of 38 (42.1%), low D levels ($< 10\text{ng/mL}$) 31 of 47 (66.0%), NNT 4.2.

<p><i>Campi</i>, 6/14/2021, prospective, Italy, peer-reviewed, 21 authors, dosage not specified.</p>	<p>risk of death for severe patients, 24.3% lower, RR 0.76, $p = 0.53$, high D levels ($\geq 20\text{ng/ml}$) 6 of 39 (15.4%), low D levels ($< 20\text{ng/ml}$) 13 of 64 (20.3%), NNT 20, hospitalized patients, outcome based on serum levels.</p>
	<p>risk of ICU for severe patients, 53.1% lower, RR 0.47, $p < 0.001$, high D levels ($\geq 20\text{ng/ml}$) 12 of 39 (30.8%), low D levels ($< 20\text{ng/ml}$) 42 of 64 (65.6%), NNT 2.9, hospitalized patients, outcome based on serum levels.</p>
<p><i>Cannata-Andía</i>, 2/18/2022, prospective, multiple countries, peer-reviewed, median age 59.0, 22 authors, study period 4 April, 2020 - 22 April, 2021, dosage 100,000IU single dose, trial NCT04552951 (history) (COVID-VIT-D), excluded in exclusion analyses: very late stage study using cholecalciferol instead of calcifediol or calcitriol.</p>	<p>risk of death, 117.0% higher, RR 2.17, $p = 0.20$, high D levels 87, low D levels 96, >25 vs. ≤ 10 ng/mL, adjusted by demographics, comorbidities, and laboratory parameters, outcome based on serum levels.</p>
	<p>risk of ICU admission, 65.0% lower, RR 0.35, $p = 0.04$, high D levels 87, low D levels 96, >25 vs. ≤ 10 ng/mL, adjusted by demographics, comorbidities, and laboratory parameters, outcome based on serum levels.</p>
	<p>risk of progression, 79.0% lower, RR 0.21, $p = 0.003$, high D levels 87, low D levels 96, pulmonary involvement at admission, >25 vs. ≤ 10 ng/mL, adjusted by demographics, comorbidities, and laboratory parameters, outcome based on serum levels.</p>
<p><i>Carpagnano</i>, 8/9/2020, retrospective, Italy, peer-reviewed, 10 authors.</p>	<p>risk of death at day 26, 70.6% lower, RR 0.29, $p = 0.0499$, high D levels 5 of 34 (14.7%), low D levels 4 of 8 (50.0%), NNT 2.8, >30 ng/mL.</p>
	<p>risk of death at day 10, 90.0% lower, RR 0.10, $p = 0.02$, high D levels 2 of 34 (5.9%), low D levels 4 of 8 (50.0%), NNT 2.3, adjusted per study, >30 ng/mL.</p>
<p><i>Cereda</i>, 11/1/2020, prospective, Italy, peer-reviewed, 13 authors.</p>	<p>risk of death, 120.0% higher, RR 2.20, $p = 0.04$, high D levels 10 of 30 (33.3%), low D levels 24 of 99 (24.2%), inverted to make $\text{RR} < 1$ favor high D levels, odds ratio converted to relative risk, $>20\text{ng/mL}$.</p>
	<p>risk of ICU admission, 86.7% lower, RR 0.13, $p = 0.59$, high D levels 0 of 30 (0.0%), low D levels 5 of 99 (5.1%), NNT 20, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm).</p>
<p><i>Cetin Ozbek</i>, 3/24/2023, retrospective, Turkey, peer-reviewed, mean age 63.4, 6 authors, study period 1 August, 2021 - 31 October, 2021.</p>	<p>risk of death, 50.9% lower, RR 0.49, $p = 0.07$, high D levels ($\geq 20\text{ng/mL}$) 7 of 61 (11.5%), low D levels ($< 20\text{ng/mL}$) 25 of 107 (23.4%), NNT 8.4.</p>
	<p>risk of death, 3.0% lower, OR 0.97, $p = 0.32$, adjusted per study, continuous values, multivariable, RR approximated with OR.</p>
<p><i>Charkowick</i>, 5/5/2022, retrospective, USA, peer-reviewed, 10 authors, study period 1 January, 2020 - 5 February, 2021.</p>	<p>risk of death, 73.4% lower, OR 0.27, $p = 0.02$, high D levels 140, low D levels 68, adjusted per study, inverted to make $\text{OR} < 1$ favor high D levels, multivariable, RR approximated with OR.</p>
	<p>risk of ICU admission, 67.2% lower, OR 0.33, $p = 0.001$, high D levels 140, low D levels 68, adjusted per study, inverted to make $\text{OR} < 1$ favor high D levels, multivariable, RR approximated with</p>

	OR.
<i>Charla</i> , 7/13/2022, retrospective, India, preprint, 8 authors, study period 1 April, 2020 - 30 April, 2021, excluded in exclusion analyses: excessive unadjusted differences between groups.	risk of death, 10.7% lower, RR 0.89, $p = 0.74$, high D levels ($\geq 20\text{ng/ml}$) 24 of 91 (26.4%), low D levels ($< 20\text{ng/ml}$) 26 of 88 (29.5%), NNT 32.
<i>Charoenngam</i> , 3/8/2021, retrospective, USA, peer-reviewed, 6 authors.	risk of death, 34.1% lower, RR 0.66, $p = 0.26$, high D levels 12 of 100 (12.0%), low D levels 29 of 187 (15.5%), adjusted per study, odds ratio converted to relative risk, $\geq 30\text{ng/mL}$.
	risk of mechanical ventilation, 37.2% lower, RR 0.63, $p = 0.17$, high D levels 14 of 100 (14.0%), low D levels 34 of 187 (18.2%), adjusted per study, odds ratio converted to relative risk, $\geq 30\text{ng/mL}$.
	risk of ICU admission, 23.1% lower, RR 0.77, $p = 0.28$, high D levels 25 of 100 (25.0%), low D levels 56 of 187 (29.9%), NNT 20, adjusted per study, odds ratio converted to relative risk, $\geq 30\text{ng/mL}$.
	risk of death, 58.1% lower, RR 0.42, $p = 0.05$, high D levels 7 of 57 (12.3%), low D levels 25 of 79 (31.6%), NNT 5.2, adjusted per study, odds ratio converted to relative risk, > 65 years old, $\geq 30\text{ng/mL}$.
<i>Chen</i> , 2/28/2023, retrospective, China, peer-reviewed, 9 authors, study period 1 June, 2022 - 5 July, 2022.	viral clearance, 40.0% improved, HR 0.60, $p = 0.01$, high D levels ($\geq 41.07\text{ng/mL}$) 52, low D levels ($< 27.5\text{ng/mL}$) 53, adjusted per study, tertile 3 vs. tertile 1, multivariable, Cox proportional hazards.
<i>Connolly</i> , 8/17/2021, retrospective, Ireland, peer-reviewed, 8 authors, study period March 2020 - May 2020.	risk of death, 90.4% lower, OR 0.10, $p = 0.06$, high D levels ($\geq 30\text{nmol/l}$) 65, low D levels ($< 30\text{nmol/l}$) 49, adjusted per study, inverted to make $\text{OR} < 1$ favor high D levels ($\geq 30\text{nmol/l}$), multivariable, RR approximated with OR.
	risk of oxygen therapy, 73.3% lower, OR 0.27, $p = 0.048$, high D levels ($\geq 30\text{nmol/l}$) 65, low D levels ($< 30\text{nmol/l}$) 49, adjusted per study, inverted to make $\text{OR} < 1$ favor high D levels ($\geq 30\text{nmol/l}$), multivariable, RR approximated with OR.
<i>Cozier</i> , 7/27/2021, prospective, USA, peer-reviewed, 6 authors.	risk of case, 38.6% lower, RR 0.61, $p = 0.04$, high D levels 94 of 1,601 (5.9%), low D levels 33 of 373 (8.8%), NNT 34, adjusted per study, inverted to make $\text{RR} < 1$ favor high D levels, odds ratio converted to relative risk, $> 20\text{ng/mL}$, multivariable.
<i>Dana</i> , 8/11/2022, retrospective, Iran, peer-reviewed, 16 authors, study period March 2020 - November 2020.	risk of death, 33.1% lower, RR 0.67, $p = 0.29$, high D levels ($\geq 10\text{ng/mL}$) 49 of 376 (13.0%), low D levels ($< 10\text{ng/mL}$) 8 of 46 (17.4%), NNT 23, adjusted per study, inverted to make $\text{RR} < 1$ favor high D levels ($\geq 10\text{ng/mL}$), odds ratio converted to relative risk, sufficiency vs. severe deficiency, multivariable.
	risk of death, 15.7% lower, RR 0.84, $p = 0.44$, high D levels ($\geq 20\text{ng/mL}$) 49 of 376 (13.0%), low D levels ($< 20\text{ng/mL}$) 30 of 197 (15.2%), NNT 46, adjusted per study, inverted to make $\text{RR} < 1$ favor high D levels ($\geq 20\text{ng/mL}$), odds ratio converted to relative risk, sufficiency vs. deficiency, multivariable.

	<p>risk of severe case, no change, RR 1.00, $p = 1.00$, high D levels ($\geq 10\text{ng/mL}$) 59 of 376 (15.7%), low D levels ($< 10\text{ng/mL}$) 7 of 46 (15.2%), adjusted per study, inverted to make $\text{RR} < 1$ favor high D levels ($\geq 10\text{ng/mL}$), odds ratio converted to relative risk, sufficiency vs. severe deficiency, multivariable.</p>
	<p>risk of severe case, 11.6% lower, RR 0.88, $p = 0.45$, high D levels ($\geq 20\text{ng/mL}$) 59 of 376 (15.7%), low D levels ($< 20\text{ng/mL}$) 35 of 197 (17.8%), NNT 48, adjusted per study, inverted to make $\text{RR} < 1$ favor high D levels ($\geq 20\text{ng/mL}$), odds ratio converted to relative risk, sufficiency vs. deficiency, multivariable.</p>
<p><i>Davoudi</i>, 5/18/2021, retrospective, Iran, peer-reviewed, 11 authors, study period February 2020 - March 2020, excluded in exclusion analyses: excessive unadjusted differences between groups.</p>	<p>risk of death, 12.3% higher, RR 1.12, $p = 1.00$, high D levels ($\geq 30\text{ng/mL}$) 2 of 57 (3.5%), low D levels ($< 30\text{ng/mL}$) 3 of 96 (3.1%).</p>
	<p>risk of mechanical ventilation, 15.8% lower, RR 0.84, $p = 1.00$, high D levels ($\geq 30\text{ng/mL}$) 1 of 57 (1.8%), low D levels ($< 30\text{ng/mL}$) 2 of 96 (2.1%), NNT 304.</p>
	<p>risk of ICU admission, 27.8% lower, RR 0.72, $p = 0.74$, high D levels ($\geq 30\text{ng/mL}$) 3 of 57 (5.3%), low D levels ($< 30\text{ng/mL}$) 7 of 96 (7.3%), NNT 49.</p>
	<p>risk of severe case, 68.4% higher, RR 1.68, $p = 0.30$, high D levels ($\geq 30\text{ng/mL}$) 9 of 57 (15.8%), low D levels ($< 30\text{ng/mL}$) 9 of 96 (9.4%).</p>
<p><i>Davran</i>, 3/15/2023, retrospective, Turkey, peer-reviewed, mean age 53.6, 9 authors.</p>	<p>risk of death, 75.4% lower, RR 0.25, $p = 0.02$, high D levels ($\geq 10\text{ng/mL}$) 4 of 63 (6.3%), low D levels ($< 10\text{ng/mL}$) 8 of 31 (25.8%), NNT 5.1.</p>
<p><i>De Smet</i>, 11/25/2020, retrospective, Belgium, peer-reviewed, 5 authors.</p>	<p>risk of death, 70.1% lower, RR 0.30, $p = 0.02$, high D levels 7 of 77 (9.1%), low D levels 20 of 109 (18.3%), adjusted per study, odds ratio converted to relative risk, $> 20\text{ng/mL}$.</p>
<p><i>Demir</i>, 1/29/2021, retrospective, Turkey, peer-reviewed, 3 authors.</p>	<p>risk of severe case, 89.3% lower, RR 0.11, $p < 0.001$, high D levels 13, low D levels 99, ratio of the mean number of affected lung segments, $> 30\text{ng/mL}$ vs. $\leq 10\text{ng/mL}$.</p>
	<p>hospitalization time, 87.1% lower, relative time 0.13, $p < 0.001$, high D levels 13, low D levels 99, $> 30\text{ng/mL}$ vs. $\leq 10\text{ng/mL}$.</p>
	<p>risk of case, 24.2% lower, RR 0.76, $p = 0.18$, high D levels 13 of 31 (41.9%), low D levels 99 of 179 (55.3%), NNT 7.5, $> 30\text{ng/mL}$ vs. $\leq 10\text{ng/mL}$.</p>
<p><i>Derakhshanian</i>, 9/19/2021, retrospective, Iran, peer-reviewed, 11 authors.</p>	<p>risk of death, 44.8% lower, RR 0.55, $p = 0.046$, high D levels 148, low D levels 142, inverted to make $\text{RR} < 1$ favor high D levels, odds ratio converted to relative risk, control prevalence approximated with overall prevalence.</p>
	<p>risk of mechanical ventilation, 41.7% lower, RR 0.58, $p = 0.09$, high D levels 148, low D levels 142, inverted to make $\text{RR} < 1$ favor high D levels, odds ratio converted to relative risk, control prevalence approximated with overall prevalence.</p>

	<p>risk of ICU admission, 37.3% lower, RR 0.63, $p = 0.04$, high D levels 148, low D levels 142, inverted to make $RR < 1$ favor high D levels, odds ratio converted to relative risk, control prevalence approximated with overall prevalence.</p>
<p><i>di Filippo (B)</i>, 8/12/2021, retrospective, Italy, peer-reviewed, 8 authors.</p>	<p>risk of death, 10.7% lower, RR 0.89, $p = 1.00$, high D levels 5 of 28 (17.9%), low D levels 12 of 60 (20.0%), NNT 47, $>20\text{ng/mL}$.</p>
	<p>risk of ICU admission, 41.6% lower, RR 0.58, $p = 0.22$, high D levels 6 of 28 (21.4%), low D levels 22 of 60 (36.7%), NNT 6.6, $>20\text{ng/mL}$.</p>
	<p>risk of severe case, 39.6% lower, RR 0.60, $p = 0.04$, high D levels 11 of 28 (39.3%), low D levels 39 of 60 (65.0%), NNT 3.9, $>20\text{ng/mL}$.</p>
<p><i>Diaz-Curiel</i>, 6/6/2021, retrospective, Spain, peer-reviewed, 8 authors.</p>	<p>risk of ICU admission, 73.2% lower, RR 0.27, $p = 0.02$, high D levels 3 of 214 (1.4%), low D levels 91 of 1,017 (8.9%), odds ratio converted to relative risk, $>30\text{ng/mL}$ vs. $<20\text{ng/mL}$.</p>
<p><i>Doğan</i>, 8/4/2022, prospective, Turkey, peer-reviewed, 5 authors, study period 1 July, 2021 - 30 October, 2021.</p>	<p>risk of case, 63.7% lower, OR 0.36, $p = 0.003$, high D levels ($\geq 10\text{ng/mL}$) 53 of 88 (60.2%) cases, 71 of 88 (80.7%) controls, NNT 4.1, case control OR.</p>
<p><i>Dror</i>, 6/7/2021, retrospective, Israel, peer-reviewed, 18 authors.</p>	<p>risk of severe or critical case, 84.8% lower, RR 0.15, $p = 0.001$, high D levels 109 of 120 (90.8%), low D levels 76 of 133 (57.1%), adjusted per study, inverted to make $RR < 1$ favor high D levels, odds ratio converted to relative risk, $>40\text{ng/mL}$ vs. $<20\text{ng/mL}$, multivariable.</p>
<p><i>Eden</i>, 8/5/2021, retrospective, United Kingdom, peer-reviewed, 5 authors.</p>	<p>risk of death, 63.9% lower, RR 0.36, $p = 0.10$, high D levels ($\geq 25\text{nmol/L}$) 3 of 26 (11.5%), low D levels ($<25\text{nmol/L}$) 8 of 25 (32.0%), NNT 4.9.</p>
	<p>risk of death, 92.9% lower, RR 0.07, $p = 0.18$, high D levels ($\geq 50\text{nmol/L}$) 0 of 8 (0.0%), low D levels ($<50\text{nmol/L}$) 11 of 43 (25.6%), NNT 3.9, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm).</p>
<p><i>Faniyi</i>, 10/6/2020, prospective, United Kingdom, preprint, 10 authors.</p>	<p>risk of seropositive, 28.8% lower, RR 0.71, $p = 0.003$, high D levels 170 of 331 (51.4%), low D levels 44 of 61 (72.1%), NNT 4.8, $>30\text{nmol/L}$.</p>
<p><i>Fatemi</i>, 11/30/2021, prospective, Iran, peer-reviewed, 5 authors, study period 1 October, 2020 - 31 May, 2021.</p>	<p>risk of death, 42.0% lower, RR 0.58, $p = 0.07$, high D levels 18 of 139 (12.9%), low D levels 25 of 109 (22.9%), NNT 10, inverted to make $RR < 1$ favor high D levels, odds ratio converted to relative risk, vitamin D measured prior to COVID-19, multivariate.</p>
	<p>risk of death, 51.1% lower, RR 0.49, $p = 0.02$, high D levels 13 of 115 (11.3%), low D levels 30 of 133 (22.6%), NNT 8.9, inverted to make $RR < 1$ favor high D levels, odds ratio converted to relative risk, vitamin D measured on admission, multivariate.</p>
	<p>risk of severe case, 37.9% lower, RR 0.62, $p = 0.007$, high D levels 38 of 139 (27.3%), low D levels 48 of 109 (44.0%), NNT</p>

	6.0, vitamin D measured prior to COVID-19.
	risk of severe case, 34.8% lower, RR 0.65, $p = 0.02$, high D levels 31 of 115 (27.0%), low D levels 55 of 133 (41.4%), NNT 6.9, vitamin D measured on admission.
<i>Faul</i> , 6/30/2020, retrospective, Ireland, peer-reviewed, 9 authors.	risk of mechanical ventilation, 69.0% lower, RR 0.31, $p = 0.03$, high D levels 4 of 21 (19.0%), low D levels 8 of 12 (66.7%), NNT 2.1, adjusted per study, >30nmol/L.
<i>Ferrer-Sánchez</i> , 3/26/2022, retrospective, Spain, peer-reviewed, 7 authors.	risk of ICU admission, 81.8% lower, RR 0.18, $p = 1.00$, high D levels ($\geq 20\text{ng/mL}$) 0 of 9 (0.0%), low D levels ($< 20\text{ng/mL}$) 4 of 73 (5.5%), NNT 18, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm), excluded in exclusion analyses: unadjusted results with no group details.
	risk of moderate/severe case, 88.7% lower, RR 0.11, $p = 1.00$, high D levels ($\geq 20\text{ng/mL}$) 0 of 9 (0.0%), low D levels ($< 20\text{ng/mL}$) 7 of 73 (9.6%), NNT 10, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm), excluded in exclusion analyses: unadjusted results with no group details.
	risk of case, 62.7% lower, OR 0.37, $p = 0.01$, cutoff 20ng/mL, adjusted per study, inverted to make OR<1 favor high D levels ($\geq 20\text{ng/mL}$), multivariable, RR approximated with OR.
<i>Freitas</i> , 3/27/2021, retrospective, Portugal, preprint, 36 authors.	risk of death, 41.2% lower, RR 0.59, $p = 0.02$, high D levels 23 of 179 (12.8%), low D levels 68 of 311 (21.9%), NNT 11, >20ng/mL.
<i>Frish</i> , 6/15/2023, retrospective, Israel, peer-reviewed, 7 authors, study period 1 February, 2020 - 31 December, 2020.	risk of case, 35.5% lower, OR 0.65, $p = 0.001$, cutoff 20ng/mL, adjusted per study, inverted to make OR<1 favor high D levels ($\geq 20\text{ng/mL}$), multivariable, RR approximated with OR.
<i>Galaznik</i> , 5/28/2021, retrospective, USA, preprint, 6 authors.	risk of case, 35.1% lower, OR 0.65, $p = 0.01$, high D levels 13,903, low D levels 2,384, adjusted per study, inverted to make OR<1 favor high D levels, breast cancer patients, logistic regression, RR approximated with OR.
	risk of case, 32.4% lower, OR 0.68, $p = 0.045$, high D levels 13,601, low D levels 1,318, adjusted per study, inverted to make OR<1 favor high D levels, prostate cancer patients, logistic regression, RR approximated with OR.
<i>Gaudio</i> , 3/27/2021, retrospective, Italy, peer-reviewed, 6 authors.	risk of case, 79.3% lower, OR 0.21, $p < 0.001$, high D levels 27 of 50 (54.0%) cases, 85 of 100 (85.0%) controls, NNT 2.7, case control OR.
<i>Gavioli</i> , 2/19/2021, retrospective, USA, peer-reviewed, 4 authors.	risk of death, 4.7% higher, RR 1.05, $p = 0.83$, high D levels 80 of 260 (30.8%), low D levels 52 of 177 (29.4%), >20ng/ml.
	risk of death, 44.8% lower, RR 0.55, $p < 0.001$, high D levels 102 of 376 (27.1%), low D levels 30 of 61 (49.2%), NNT 4.5, >10ng/ml.

	<p>risk of oxygen therapy, 55.2% lower, RR 0.45, $p < 0.001$, high D levels 127 of 260 (48.8%), low D levels 116 of 177 (65.5%), NNT 6.0, adjusted per study, inverted to make RR<1 favor high D levels, >20ng/ml, multivariate.</p>
	<p>risk of hospitalization, 3.6% lower, RR 0.96, $p = 0.41$, high D levels 218 of 260 (83.8%), low D levels 154 of 177 (87.0%), NNT 32, >20ng/ml.</p>
<i>Ghanei</i> , 3/23/2022, prospective, Iran, peer-reviewed, 6 authors, study period 20 March, 2020 - 20 January, 2021.	<p>risk of case, 42.1% lower, OR 0.58, $p = 0.09$, high D levels (≥ 20ng/ml) 58 of 90 (64.4%) cases, 72 of 95 (75.8%) controls, NNT 7.4, case control OR.</p>
<i>Gholi</i> , 7/19/2022, prospective, Iran, peer-reviewed, 4 authors.	<p>risk of death, 74.7% lower, HR 0.25, $p < 0.001$, high D levels 157, low D levels 38, inverted to make HR<1 favor high D levels, >30ng/mL vs. <20ng/mL, model 2, day 45.</p>
	<p>risk of death, 39.8% lower, HR 0.60, $p = 0.05$, high D levels 157, low D levels 38, inverted to make HR<1 favor high D levels, >30ng/mL vs. <20ng/mL, ICU mortality, model 2.</p>
	<p>risk of mechanical ventilation, 44.9% higher, HR 1.45, $p = 0.27$, high D levels 157, low D levels 38, inverted to make HR<1 favor high D levels, >30ng/mL vs. <20ng/mL, model 2, day 45.</p>
<i>Golabi</i> , 8/26/2021, retrospective, Iran, peer-reviewed, 10 authors.	<p>odds of symptoms, 90.0% lower, OR 0.10, $p < 0.001$, high D levels 34, low D levels 10, >30ng/mL vs. <20ng/mL, GEE regression, RR approximated with OR.</p>
	<p>odds of symptoms, 81.0% lower, OR 0.19, $p = 0.006$, high D levels 34, low D levels 9, 20-30ng/mL vs. <20ng/mL, GEE regression, RR approximated with OR.</p>
	<p>risk of case, 71.7% lower, OR 0.28, $p = 0.07$, high D levels 34 of 44 (77.3%) cases, 36 of 39 (92.3%) controls, NNT 3.5, case control OR, >30ng/mL vs. <20ng/mL.</p>
<i>Gonzalez</i> , 3/13/2023, retrospective, Argentina, peer-reviewed, 10 authors.	<p>risk of death, 66.1% lower, OR 0.34, $p = 0.046$, high D levels (≥ 12ng/ml) 129, low D levels (<12ng/ml) 35, adjusted per study, inverted to make OR<1 favor high D levels (≥ 12ng/ml), multivariable, RR approximated with OR.</p>
<i>González-Estevez</i> , 7/7/2021, retrospective, Mexico, peer-reviewed, 6 authors.	<p>risk of symptomatic case, 25.0% lower, RR 0.75, $p = 0.04$, high D levels (≥ 30ng/mL) 6 of 8 (75.0%), low D levels (<30ng/mL) 32 of 32 (100.0%), NNT 4.0.</p>
<i>Green</i> , 11/7/2022, retrospective, Israel, peer-reviewed, 9 authors, study period 1 February, 2020 - 31 December, 2020.	<p>risk of case, 18.7% lower, OR 0.81, $p < 0.001$, cutoff 30ng/mL, adjusted per study, inverted to make OR<1 favor high D levels (≥ 30ng/mL), multivariable, RR approximated with OR.</p>
<i>Gönen</i> , 11/12/2021, retrospective, Turkey, peer-reviewed, 20 authors, dosage varies.	<p>risk of death, 65.8% lower, RR 0.34, $p = 0.62$, high D levels (≥ 12ng/mL) 1 of 80 (1.2%), low D levels (<12ng/mL) 3 of 82 (3.7%), NNT 42, retrospective study.</p>
	<p>risk of ICU admission, 16.9% lower, RR 0.83, $p = 1.00$, high D levels (≥ 12ng/mL) 4 of 77 (5.2%), low D levels (<12ng/mL) 5 of</p>

	80 (6.2%), NNT 95, retrospective study.
	hospital stay >8 days, 21.1% lower, RR 0.79, $p = 0.11$, high D levels ($\geq 12\text{ng/mL}$) 40 of 78 (51.3%), low D levels ($< 12\text{ng/mL}$) 52 of 80 (65.0%), NNT 7.3, retrospective study.
<i>Hafez</i> , 3/29/2022, retrospective, United Arab Emirates, peer-reviewed, mean age 43.0, 11 authors.	risk of death, 97.7% lower, RR 0.02, $p = 0.02$, high D levels ($\geq 12\text{ng/mL}$) 6 of 116 (5.2%), low D levels ($< 12\text{ng/mL}$) 3 of 10 (30.0%), NNT 4.0, adjusted per study, inverted to make $\text{RR} < 1$ favor high D levels ($\geq 12\text{ng/mL}$), odds ratio converted to relative risk, multivariable, model 2.
	risk of death, 96.3% lower, RR 0.04, $p = 0.04$, high D levels ($\geq 20\text{ng/mL}$) 4 of 64 (6.2%), low D levels ($< 20\text{ng/mL}$) 5 of 62 (8.1%), adjusted per study, inverted to make $\text{RR} < 1$ favor high D levels ($\geq 20\text{ng/mL}$), odds ratio converted to relative risk, multivariable, model 3.
<i>Hastie</i> , 8/26/2020, retrospective, population-based cohort, database analysis, United Kingdom, peer-reviewed, 14 authors.	risk of death, 17.4% lower, RR 0.83, $p = 0.31$, cutoff 25nmol/L, adjusted per study, inverted to make $\text{RR} < 1$ favor high D levels ($\geq 25\text{nmol/L}$), multivariable Cox.
	risk of hospitalization, 9.1% lower, RR 0.91, $p = 0.40$, cutoff 25nmol/L, adjusted per study, inverted to make $\text{RR} < 1$ favor high D levels ($\geq 25\text{nmol/L}$), multivariable Cox.
<i>Hermawan</i> , 3/28/2023, retrospective, Indonesia, peer-reviewed, survey, 5 authors, study period March 2022 - July 2022.	risk of symptomatic case, 70.6% lower, RR 0.29, $p < 0.001$, high D levels ($\geq 10\text{ng/ml}$) 10 of 34 (29.4%), low D levels ($< 10\text{ng/ml}$) 13 of 13 (100.0%), NNT 1.4.
	risk of symptomatic case, 45.6% lower, RR 0.54, $p = 0.42$, high D levels ($\geq 20\text{ng/ml}$) 2 of 7 (28.6%), low D levels ($< 20\text{ng/ml}$) 21 of 40 (52.5%), NNT 4.2.
<i>Hernández</i> , 10/27/2020, retrospective, Spain, peer-reviewed, mean age 60.9, 12 authors.	risk of combined death/ICU/ventilation, 83.0% lower, RR 0.17, $p < 0.001$, high D levels 35, low D levels 162, $\geq 20\text{ng/mL}$ risk of hospitalization * risk of death/ICU/ventilation hospitalization.
	risk of combined death/ICU/ventilation if hospitalized, 12.0% lower, RR 0.88, $p = 0.86$, high D levels 35, low D levels 162, $\geq 20\text{ng/mL}$ risk of death/ICU/ventilation hospitalization.
	risk of hospitalization, 80.6% lower, RR 0.19, $p < 0.001$, $\geq 20\text{ng/mL}$.
<i>Hogarth</i> , 5/3/2023, retrospective, USA, peer-reviewed, median age 56.0, 9 authors, study period 1 January, 2021 - 8 November, 2021.	risk of case, 46.5% lower, OR 0.53, $p < 0.001$, high D levels ($\geq 20\text{ng/mL}$) 96,894, low D levels ($< 20\text{ng/mL}$) 13,486, adjusted per study, inverted to make $\text{OR} < 1$ favor high D levels ($\geq 20\text{ng/mL}$), breakthrough case, multivariable, RR approximated with OR.
<i>Huang</i> , 3/24/2023, retrospective, China, peer-reviewed, 5 authors, study period 14 June, 2021 - 1 April, 2022.	recovery time, 25.0% lower, relative time 0.75, $p = 0.02$, high D levels ($\geq 20\text{ng/ml}$) 28, low D levels ($< 20\text{ng/ml}$) 18, relative time until resolution of pneumonia.

<i>Hurst</i> , 10/22/2021, prospective, United Kingdom, peer-reviewed, 23 authors.	risk of death, 68.4% lower, RR 0.32, $p = 0.005$, high D levels 68, low D levels 191, odds ratio converted to relative risk, >50nmol/l, multivariable, Supplementary Table 2, control prevalence approximated with overall prevalence.
	risk of mechanical ventilation, 66.0% lower, RR 0.34, $p = 0.004$, high D levels 6 of 68 (8.8%), low D levels 61 of 191 (31.9%), NNT 4.3, odds ratio converted to relative risk, >50nmol/l, multivariable, Supplementary Table 2.
<i>Im</i> , 8/11/2020, retrospective, South Korea, peer-reviewed, 6 authors.	risk of case, 73.1% lower, OR 0.27, $p < 0.001$, high D levels 13 of 50 (26.0%) cases, 85 of 150 (56.7%) controls, NNT 4.3, case control OR.
<i>Infante</i> , 2/18/2021, retrospective, Italy, peer-reviewed, 11 authors.	risk of death, 54.8% lower, RR 0.45, $p = 0.046$, high D levels 4 of 19 (21.1%), low D levels 55 of 118 (46.6%), NNT 3.9, >20ng/mL.
<i>Israel</i> , 9/20/2021, retrospective, population-based cohort, Israel, peer-reviewed, 9 authors, study period 1 March, 2020 - 31 October, 2020.	risk of severe case, 33.9% lower, OR 0.66, $p < 0.001$, high D levels 423 of 1,036 (40.8%) cases, 509 of 934 (54.5%) controls, NNT 7.3, adjusted per study, inverted to make OR<1 favor high D levels, case control OR, >75 nmol/L vs. <30 nmol/L, multivariable.
	risk of case, 19.7% lower, OR 0.80, $p < 0.001$, high D levels 6,152 of 15,892 (38.7%) cases, 73,810 of 159,193 (46.4%) controls, NNT 39, adjusted per study, inverted to make OR<1 favor high D levels, case control OR, >75 nmol/L vs. <30 nmol/L, among COVID+ cases, multivariable.
<i>Jain (B)</i> , 11/19/2020, prospective, India, peer-reviewed, 6 authors.	risk of death, 85.2% lower, RR 0.15, $p = 0.001$, high D levels 2 of 64 (3.1%), low D levels 19 of 90 (21.1%), NNT 5.6, >20ng/mL.
	risk of ICU admission, 95.4% lower, RR 0.05, $p < 0.001$, high D levels 2 of 64 (3.1%), low D levels 61 of 90 (67.8%), NNT 1.5, >20ng/mL.
<i>Jalavu</i> , 6/1/2023, prospective, South Africa, peer-reviewed, 16 authors, study period 29 October, 2020 - 10 February, 2021.	risk of death, 1.0% lower, HR 0.99, $p = 0.97$, high D levels (≥ 50 nmol/L) 16 of 31 (51.6%), low D levels (<50nmol/L) 38 of 55 (69.1%), NNT 5.7, Kaplan–Meier.
<i>Jimenez</i> , 7/26/2021, retrospective, Spain, peer-reviewed, 21 authors, study period 12 March, 2020 - 21 May, 2020, dosage paricalcitol 0.9µg weekly, excluded in exclusion analyses: many patients received vitamin D treatment.	risk of death, 7.7% higher, OR 1.08, $p = 0.81$, high D levels 50, low D levels 110, >30 vs. <20ng/ml, RR approximated with OR, outcome based on serum levels.
	risk of mechanical ventilation, 47.5% lower, OR 0.53, $p = 0.56$, high D levels 50, low D levels 110, >30 vs. <20ng/ml, RR approximated with OR, outcome based on serum levels.
	risk of ICU admission, 12.2% lower, OR 0.88, $p = 0.87$, high D levels 50, low D levels 110, >30 vs. <20ng/ml, RR approximated with OR, outcome based on serum levels.
	risk of hospitalization, 0.8% lower, OR 0.99, $p = 0.98$, high D levels 50, low D levels 110, >30 vs. <20ng/ml, RR approximated with OR, outcome based on serum levels.

<p><i>Jude</i>, 6/17/2021, retrospective, United Kingdom, peer-reviewed, 5 authors.</p>	<p>risk of hospitalization, 71.6% lower, RR 0.28, $p < 0.001$, adjusted per study, inverted to make $RR < 1$ favor high D levels, odds ratio converted to relative risk, > 25 nmol/L, control prevalence approximated with overall prevalence.</p>
	<p>risk of hospitalization, 57.9% lower, RR 0.42, $p < 0.001$, adjusted per study, inverted to make $RR < 1$ favor high D levels, odds ratio converted to relative risk, > 50 nmol/L, control prevalence approximated with overall prevalence.</p>
<p><i>Junior</i>, 2/17/2022, prospective, Brazil, peer-reviewed, 6 authors, dosage not specified.</p>	<p>risk of mechanical ventilation, 84.4% lower, OR 0.16, $p = 0.03$, cutoff 40ng/dl, inverted to make $OR < 1$ favor high D levels (≥ 40ng/dl), risk of mechanical ventilation for vitamin D levels > 40ng/ml, RR approximated with OR, outcome based on serum levels.</p>
<p><i>Juraj</i>, 1/22/2022, retrospective, Slovakia, peer-reviewed, 13 authors, study period 1 November, 2020 - 30 April, 2021.</p>	<p>risk of death, 19.0% lower, RR 0.81, $p = 0.05$, high D levels (≥ 12ng/mL) 127 of 283 (44.9%), low D levels (< 12ng/mL) 41 of 74 (55.4%), NNT 9.5.</p>
<p><i>Kalichuran</i>, 4/26/2022, prospective, South Africa, peer-reviewed, survey, 4 authors, study period September 2020 - February 2021.</p>	<p>risk of symptomatic case, 60.0% lower, RR 0.40, $p < 0.001$, high D levels (≥ 20ng/mL) 56, low D levels (< 20ng/mL) 44, inverted to make $RR < 1$ favor high D levels (≥ 20ng/mL).</p>
	<p>risk of symptomatic case, 58.2% lower, RR 0.42, $p = 0.004$, inverted to make $RR < 1$ favor high D levels, higher sunlight exposure vs. lower sunlight exposure.</p>
<p><i>Karahan</i>, 10/5/2020, retrospective, Turkey, peer-reviewed, 2 authors.</p>	<p>risk of death, 82.5% lower, RR 0.17, $p < 0.001$, high D levels 5 of 46 (10.9%), low D levels 64 of 103 (62.1%), NNT 2.0, > 20nmol/L.</p>
<p><i>Karonova (B)</i>, 3/2/2022, retrospective, Russia, peer-reviewed, 11 authors, study period 30 November, 2020 - 20 March, 2021.</p>	<p>risk of severe case, 22.5% lower, OR 0.78, $p = 0.01$, cutoff 11.4ng/mL, adjusted per study, inverted to make $OR < 1$ favor high D levels (≥ 11.4ng/mL), multivariable, RR approximated with OR.</p>
<p><i>Karonova (C)</i>, 8/29/2021, retrospective, Russia, peer-reviewed, 8 authors, study period April 2020 - December 2020.</p>	<p>risk of death, 77.8% lower, RR 0.22, $p = 0.006$, high D levels 8 of 96 (8.3%), low D levels 10 of 37 (27.0%), NNT 5.3, adjusted per study, inverted to make $RR < 1$ favor high D levels, odds ratio converted to relative risk, > 10ng/mL, logistic regression model 2.</p>
	<p>risk of death, 84.8% lower, RR 0.15, $p = 0.06$, high D levels 1 of 43 (2.3%), low D levels 17 of 90 (18.9%), NNT 6.0, adjusted per study, inverted to make $RR < 1$ favor high D levels, odds ratio converted to relative risk, > 20ng/mL, logistic regression model 2.</p>
	<p>risk of severe case, 67.3% lower, RR 0.33, $p = 0.005$, high D levels 12 of 96 (12.5%), low D levels 13 of 37 (35.1%), NNT 4.4, adjusted per study, inverted to make $RR < 1$ favor high D levels, odds ratio converted to relative risk, > 10ng/mL, logistic regression model 2.</p>

	<p>risk of severe case, 53.2% lower, RR 0.47, $p = 0.13$, high D levels 4 of 43 (9.3%), low D levels 21 of 90 (23.3%), NNT 7.1, adjusted per study, inverted to make $RR < 1$ favor high D levels, odds ratio converted to relative risk, >20ng/mL, logistic regression model 2.</p>
<p><i>Karonova (D)</i>, 12/31/2020, retrospective, Russia, peer-reviewed, 3 authors.</p>	<p>risk of death, 79.4% lower, RR 0.21, $p = 0.11$, high D levels 1 of 23 (4.3%), low D levels 12 of 57 (21.1%), NNT 6.0, inverted to make $RR < 1$ favor high D levels, odds ratio converted to relative risk, >20ng/ml.</p>
	<p>risk of severe case, 71.1% lower, RR 0.29, $p = 0.05$, high D levels 3 of 23 (13.0%), low D levels 22 of 57 (38.6%), NNT 3.9, inverted to make $RR < 1$ favor high D levels, odds ratio converted to relative risk, >20ng/ml.</p>
<p><i>Katz</i>, 12/4/2020, retrospective, population-based cohort, USA, peer-reviewed, 3 authors.</p>	<p>risk of case, 78.4% lower, RR 0.22, $p < 0.001$, high D levels 85 of 101,175 (0.1%), low D levels 87 of 31,950 (0.3%), NNT 531, adjusted per study, inverted to make $RR < 1$ favor high D levels.</p>
<p><i>Kaufman</i>, 9/17/2020, retrospective, population-based cohort, USA, peer-reviewed, median age 54.0, 5 authors.</p>	<p>risk of case, 53.0% lower, RR 0.47, $p < 0.001$, high D levels 12,321, low D levels 39,190, >55 ng/mL vs. <20 ng/mL.</p>
<p><i>Kaur</i>, 11/30/2021, prospective, India, peer-reviewed, 5 authors, excluded in exclusion analyses: unadjusted results with no group details.</p>	<p>risk of death, 89.8% lower, RR 0.10, $p < 0.001$, high D levels (≥ 10ng/mL) 5 of 64 (7.8%), low D levels (<10ng/mL) 13 of 17 (76.5%), NNT 1.5.</p>
	<p>risk of mechanical ventilation, 90.3% lower, RR 0.10, $p < 0.001$, high D levels (≥ 10ng/mL) 4 of 64 (6.2%), low D levels (<10ng/mL) 11 of 17 (64.7%), NNT 1.7.</p>
<p><i>Kazemi</i>, 5/7/2022, retrospective, Iran, peer-reviewed, mean age 56.0, 4 authors.</p>	<p>risk of death, 75.8% lower, RR 0.24, $p = 0.26$, high D levels (≥ 30ng/mL) 1 of 75 (1.3%), low D levels (<30ng/mL) 7 of 127 (5.5%), NNT 24.</p>
	<p>risk of severe case, 4.8% higher, RR 1.05, $p = 1.00$, high D levels (≥ 30ng/mL) 13 of 75 (17.3%), low D levels (<30ng/mL) 21 of 127 (16.5%).</p>
<p><i>Khalil</i>, 11/8/2022, retrospective, Iraq, peer-reviewed, 3 authors.</p>	<p>risk of case, 41.6% lower, OR 0.58, $p = 0.27$, high D levels (≥ 10ng/ml) 30 of 52 (57.7%) cases, 21 of 30 (70.0%) controls, NNT 8.2, case control OR.</p>
<p><i>Lau</i>, 4/28/2020, retrospective, USA, preprint, 7 authors.</p>	<p>risk of ICU admission, 45.0% lower, RR 0.55, $p = 0.29$, high D levels 2 of 5 (40.0%), low D levels 11 of 15 (73.3%), NNT 3.0, >30ng/mL.</p>
<p><i>Li</i>, 5/19/2021, retrospective, USA, peer-reviewed, 4 authors.</p>	<p>risk of case, 8.6% lower, RR 0.91, $p = 0.24$, high D levels 610 of 13,650 (4.5%), low D levels 290 of 4,498 (6.4%), adjusted per study, inverted to make $RR < 1$ favor high D levels, odds ratio converted to relative risk, >20ng/mL, Figure 2.</p>
	<p>risk of case, 12.4% lower, RR 0.88, $p = 0.07$, high D levels 289 of 7,272 (4.0%), low D levels 611 of 10,876 (5.6%), adjusted per study, inverted to make $RR < 1$ favor high D levels, odds ratio</p>

	converted to relative risk, >30ng/mL, Figure 2.
<i>Li (B)</i> , 1/11/2021, retrospective, population-based cohort, United Kingdom, peer-reviewed, 6 authors.	risk of hospitalization, 36.2% lower, RR 0.64, $p < 0.001$, NNT 932, inverted to make RR<1 favor high D levels, odds ratio converted to relative risk, >25nmol/L.
	risk of case, 29.5% lower, RR 0.71, $p < 0.001$, NNT 823, inverted to make RR<1 favor high D levels, odds ratio converted to relative risk, >25nmol/L.
<i>Livingston</i> , 4/2/2021, retrospective, United Kingdom, peer-reviewed, 7 authors.	risk of case, 50.9% lower, RR 0.49, $p = 0.02$, high D levels 16 of 52 (30.8%), low D levels 31 of 52 (59.6%), NNT 3.5, odds ratio converted to relative risk, >34.4nmol/L.
<i>Lohia</i> , 3/4/2021, retrospective, USA, peer-reviewed, 4 authors.	risk of death, 14.7% lower, RR 0.85, $p = 0.56$, high D levels 88, low D levels 95, odds ratio converted to relative risk, control prevalence approximated with overall prevalence, >30 ng/mL vs. <20 ng/mL, >30 ng/mL group size approximated.
	risk of mechanical ventilation, 18.9% lower, RR 0.81, $p = 0.48$, high D levels 88, low D levels 95, odds ratio converted to relative risk, control prevalence approximated with overall prevalence, >30 ng/mL vs. <20 ng/mL, >30 ng/mL group size approximated.
	risk of ICU admission, 28.5% lower, RR 0.72, $p = 0.17$, high D levels 88, low D levels 95, odds ratio converted to relative risk, control prevalence approximated with overall prevalence, >30 ng/mL vs. <20 ng/mL, >30 ng/mL group size approximated.
<i>Luo</i> , 11/13/2020, retrospective, China, peer-reviewed, median age 56.0, 5 authors.	risk of progression, 63.0% lower, RR 0.37, $p = 0.01$, high D levels 335, low D levels 560, >30nmol/L.
<i>Ma</i> , 12/3/2021, retrospective, USA, peer-reviewed, 16 authors, study period May 2020 - March 2021, dosage varies.	risk of hospitalization, 67.0% lower, OR 0.33, $p = 0.15$, high D levels 7,893, low D levels 7,823, adjusted per study, highest quintile vs. lowest quintile predicted vitamin D levels, model 3, supplemental table 3, multivariable, RR approximated with OR, outcome based on serum levels.
	risk of symptomatic case, 9.0% lower, OR 0.91, $p = 0.52$, high D levels 7,893, low D levels 7,823, adjusted per study, highest quintile vs. lowest quintile predicted vitamin D levels, model 3, supplemental table 3, multivariable, RR approximated with OR, outcome based on serum levels.
	risk of case, 52.0% lower, OR 0.48, $p = 0.01$, high D levels 7,893, low D levels 7,823, adjusted per study, highest quintile vs. lowest quintile predicted vitamin D levels, model 3, supplemental table 3, multivariable, RR approximated with OR, outcome based on serum levels.
<i>Macaya</i> , 10/21/2020, retrospective, Spain, peer-reviewed, 8 authors.	risk of severe case, 55.0% lower, RR 0.45, $p = 0.07$, high D levels 11 of 35 (31.4%), low D levels 20 of 45 (44.4%), NNT 7.7, inverted to make RR<1 favor high D levels, odds ratio converted to relative risk, >20ng/mL.
<i>Maghbooli (B)</i> , 9/25/2020, retrospective, Iran, peer-reviewed, 11 authors.	risk of death, 51.7% lower, RR 0.48, $p = 0.08$, high D levels 7 of 72 (9.7%), low D levels 27 of 134 (20.1%), NNT 9.6, age >40.

	<p>risk of mechanical ventilation, 31.6% lower, RR 0.68, $p = 0.49$, high D levels 6 of 77 (7.8%), low D levels 18 of 158 (11.4%), NNT 28.</p> <p>risk of ICU admission, 32.0% lower, RR 0.68, $p = 0.33$, high D levels 11 of 77 (14.3%), low D levels 33 of 158 (20.9%), NNT 15, >30nmol/L.</p>
<i>Manojlovic</i> , 6/15/2023, retrospective, Serbia, peer-reviewed, mean age 57.6, 11 authors, excluded in exclusion analyses: unadjusted differences between groups.	risk of death, 89.9% lower, RR 0.10, $p = 0.009$, high D levels ($\geq 30\text{nmol/l}$) 1 of 41 (2.4%), low D levels ($< 30\text{nmol/l}$) 8 of 33 (24.2%), NNT 4.6.
<i>Martínez-Rodríguez</i> , 3/31/2022, retrospective, Mexico, peer-reviewed, 5 authors.	risk of death, 52.2% lower, OR 0.48, $p = 0.04$, cutoff 20ng/mL, adjusted per study, multivariable, RR approximated with OR.
<i>Matin</i> , 7/30/2021, retrospective, case control, Iran, peer-reviewed, 8 authors.	risk of case, 66.1% lower, OR 0.34, $p < 0.001$, inverted to make $\text{OR} < 1$ favor high D levels, case control OR, >20ng/mL.
<i>Mayurathan</i> , 8/8/2023, retrospective, Sri Lanka, peer-reviewed, 11 authors.	risk of death, 98.2% higher, RR 1.98, $p = 0.69$, high D levels ($\geq 20\text{ng/mL}$) 8 of 113 (7.1%), low D levels ($< 20\text{ng/mL}$) 1 of 28 (3.6%).
	risk of severe case, 67.3% higher, RR 1.67, $p = 0.32$, high D levels ($\geq 20\text{ng/mL}$) 27 of 113 (23.9%), low D levels ($< 20\text{ng/mL}$) 4 of 28 (14.3%).
<i>Mazziotti</i> , 3/5/2021, retrospective, Italy, peer-reviewed, 11 authors, dosage varies.	risk of death, 2.4% lower, RR 0.98, $p = 0.91$, high D levels 187, low D levels 161, inverted to make $\text{RR} < 1$ favor high D levels, odds ratio converted to relative risk, >12ng/mL, control prevalence approximated with overall prevalence, outcome based on serum levels.
	risk of acute hypoxemic respiratory failure, 37.0% lower, RR 0.63, $p = 0.006$, high D levels 72 of 187 (38.5%), low D levels 97 of 161 (60.2%), NNT 4.6, inverted to make $\text{RR} < 1$ favor high D levels, odds ratio converted to relative risk, >12ng/mL, outcome based on serum levels.
<i>Meltzer</i> , 3/19/2021, retrospective, database analysis, USA, peer-reviewed, 6 authors.	risk of case, 34.6% lower, RR 0.65, $p = 0.11$, high D levels 61 of 1,097 (5.6%), low D levels 118 of 1,251 (9.4%), NNT 26, adjusted per study, inverted to make $\text{RR} < 1$ favor high D levels, >40ng/mL vs. <20ng/mL, Table 2, Model 2.
<i>Meltzer (B)</i> , 9/3/2020, retrospective, USA, peer-reviewed, 6 authors.	risk of case, 43.5% lower, RR 0.56, $p = 0.02$, high D levels 39 of 317 (12.3%), low D levels 32 of 172 (18.6%), NNT 16, adjusted per study, inverted to make $\text{RR} < 1$ favor high D levels, >20ng/mL.
<i>Mendy</i> , 6/27/2020, retrospective, USA, preprint, 4 authors.	risk of death, 7.0% lower, RR 0.93, $p = 0.89$, high D levels 21 of 600 (3.5%), low D levels 5 of 89 (5.6%), inverted to make $\text{RR} < 1$ favor high D levels, odds ratio converted to relative risk.
	risk of death/ICU, 16.7% lower, RR 0.83, $p < 0.001$, high D levels 68 of 600 (11.3%), low D levels 23 of 89 (25.8%), NNT 6.9, inverted to make $\text{RR} < 1$ favor high D levels, odds ratio converted to relative risk.

	<p>risk of ICU admission, 55.3% lower, RR 0.45, $p = 0.008$, high D levels 47 of 600 (7.8%), low D levels 18 of 89 (20.2%), NNT 8.1, inverted to make RR<1 favor high D levels, odds ratio converted to relative risk.</p>
	<p>risk of hospitalization, 15.1% lower, RR 0.85, $p < 0.001$, high D levels 171 of 600 (28.5%), low D levels 45 of 89 (50.6%), NNT 4.5, inverted to make RR<1 favor high D levels, odds ratio converted to relative risk.</p>
<p><i>Merzon</i>, 7/23/2020, retrospective, Israel, peer-reviewed, 3 authors.</p>	<p>risk of hospitalization, 46.4% lower, RR 0.54, $p = 0.06$, high D levels 79, low D levels 703, odds ratio converted to relative risk, >30ng/mL.</p>
	<p>risk of case, 28.4% lower, RR 0.72, $p < 0.001$, high D levels 1,139, low D levels 6,668, odds ratio converted to relative risk, >30ng/mL.</p>
<p><i>Mingiano</i>, 7/30/2023, retrospective, Italy, peer-reviewed, 11 authors, study period November 2020 - February 2021, dosage calcifediol 450µg days 1-2, patients with deficiency only.</p>	<p>risk of death, 49.8% lower, RR 0.50, $p = 0.005$, cutoff 10ng/mL, outcome based on serum levels.</p>
	<p>risk of death, 35.9% lower, RR 0.64, $p = 0.04$, cutoff 20ng/mL, outcome based on serum levels.</p>
<p><i>Mostafa</i>, 11/30/2022, retrospective, Egypt, peer-reviewed, 10 authors, study period November 2020 - December 2021, excluded in exclusion analyses: categorical results are unadjusted with significant differences between groups.</p>	<p>risk of death, 92.8% lower, RR 0.07, $p < 0.001$, high D levels (≥ 20ng/mL) 4 of 135 (3.0%), low D levels (<20ng/mL) 21 of 51 (41.2%), NNT 2.6, unadjusted, normal vs. deficiency.</p>
	<p>risk of mechanical ventilation, 95.0% lower, RR 0.05, $p < 0.001$, high D levels (≥ 20ng/mL) 4 of 135 (3.0%), low D levels (<20ng/mL) 30 of 51 (58.8%), NNT 1.8, unadjusted, normal vs. deficiency.</p>
	<p>risk of ICU admission, 90.6% lower, RR 0.09, $p < 0.001$, high D levels (≥ 20ng/mL) 9 of 135 (6.7%), low D levels (<20ng/mL) 36 of 51 (70.6%), NNT 1.6, unadjusted, normal vs. deficiency.</p>
<p><i>Nasiri</i>, 6/30/2021, retrospective, Iran, peer-reviewed, 3 authors.</p>	<p>risk of death, 8.9% higher, OR 1.09, $p = 0.89$, high D levels (≥ 30ng/mL) 238, low D levels (<20ng/mL) 43, inverted to make OR<1 favor high D levels (≥ 30ng/mL), RR approximated with OR.</p>
<p><i>Neves</i>, 6/14/2022, retrospective, Brazil, peer-reviewed, mean age 62.1, 7 authors, study period July 2020 - December 2020, excluded in exclusion analyses: excessive unadjusted differences between groups.</p>	<p>risk of death, 57.1% lower, RR 0.43, $p = 0.046$, high D levels (≥ 50nmol/L) 12 of 87 (13.8%), low D levels (<50nmol/L) 9 of 28 (32.1%), NNT 5.4.</p>
	<p>risk of ICU admission, 19.5% higher, RR 1.20, $p = 0.81$, high D levels (≥ 50nmol/L) 26 of 87 (29.9%), low D levels (<50nmol/L) 7 of 28 (25.0%).</p>
<p><i>Nguyen</i>, 5/3/2022, retrospective, USA, peer-reviewed, 11 authors, study period 15 July, 2020 - 15 October, 2020.</p>	<p>risk of death, 81.1% lower, OR 0.19, $p = 0.008$, cutoff 20ng/mL, adjusted per study, inverted to make OR<1 favor high D levels (≥ 20ng/mL), 25-OH-D3, multivariable, RR approximated with OR.</p>
	<p>risk of mechanical ventilation, 52.8% lower, OR 0.47, $p = 0.13$, cutoff 20ng/mL, adjusted per study, inverted to make OR<1 favor high D levels (≥ 20ng/mL), 25-OH-D3, multivariable, RR</p>

	approximated with OR.
	risk of no hospital discharge, 74.0% lower, HR 0.26, $p < 0.001$, cutoff 20ng/mL, 25-OH-D3, Cox proportional hazards.
<i>Nimavat</i> , 8/5/2021, retrospective, India, peer-reviewed, 5 authors.	risk of death, 50.4% lower, RR 0.50, $p = 0.17$, high D levels 13 of 131 (9.9%), low D levels 5 of 25 (20.0%), NNT 9.9, >10ng/mL, within cases.
	risk of severe case, 67.6% lower, RR 0.32, $p = 0.003$, high D levels 17 of 131 (13.0%), low D levels 10 of 25 (40.0%), NNT 3.7, >10ng/mL, within cases.
<i>Orchard</i> , 1/19/2021, retrospective, United Kingdom, peer-reviewed, 7 authors.	risk of ICU admission, 58.8% lower, RR 0.41, $p = 0.001$, high D levels 9 of 40 (22.5%), low D levels 41 of 75 (54.7%), NNT 3.1, all hospitalized patients, >50 nmol/L.
	risk of death, 24.1% lower, RR 0.76, $p = 1.00$, high D levels 1 of 9 (11.1%), low D levels 6 of 41 (14.6%), NNT 28, ICU patients only, >50 nmol/L.
	risk of mechanical ventilation, 8.9% lower, RR 0.91, $p = 0.70$, high D levels 6 of 9 (66.7%), low D levels 30 of 41 (73.2%), NNT 15, ICU patients only, >50 nmol/L.
<i>Ortatatli</i> , 2/14/2023, prospective, Turkey, peer-reviewed, 9 authors, excluded in exclusion analyses: unadjusted results with no group details.	risk of death, 82.1% lower, RR 0.18, $p = 0.09$, cutoff 20ng/mL, inverted to make $RR < 1$ favor high D levels (≥ 20 ng/mL), 25(OH)D.
	risk of death, 73.7% lower, RR 0.26, $p = 0.04$, cutoff 1ng/mL, inverted to make $RR < 1$ favor high D levels (≥ 1 ng/mL), 1,25(OH)2D.
<i>Ozturk</i> , 5/16/2022, retrospective, Turkey, peer-reviewed, 6 authors, excluded in exclusion analyses: unadjusted results with no group details.	risk of severe case, 46.4% lower, RR 0.54, $p = 0.10$, high D levels (≥ 20 ng/mL) 9 of 110 (8.2%), low D levels (<20ng/mL) 29 of 190 (15.3%), NNT 14.
<i>Panagiotou</i> , 6/30/2020, retrospective, United Kingdom, preprint, 12 authors.	risk of ICU admission, 52.0% lower, RR 0.48, $p = 0.02$, high D levels 8 of 44 (18.2%), low D levels 34 of 90 (37.8%), NNT 5.1, >50nmol/L.
<i>Pande</i> , 3/16/2022, retrospective, India, peer-reviewed, 7 authors, study period October 2020 - October 2021, excluded in exclusion analyses: unadjusted results with no group details.	risk of severe case, 93.4% lower, RR 0.07, $p < 0.001$, high D levels (≥ 20 ng/ml) 7 of 116 (6.0%), low D levels (<20ng/ml) 85 of 93 (91.4%), NNT 1.2.
<i>Parra-Ortega</i> , 8/24/2021, prospective, Mexico, peer-reviewed, 9 authors, excluded in exclusion analyses: unadjusted results with no group details.	risk of death, 98.7% lower, RR 0.01, $p < 0.001$, high D levels (≥ 20 ng/dL) 0 of 15 (0.0%), low D levels (<20ng/dL) 63 of 79 (79.7%), NNT 1.3, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm), unadjusted.
<i>Pecina</i> , 8/27/2021, retrospective, USA, peer-reviewed, 4 authors, dosage not specified.	risk of death, 35.9% lower, RR 0.64, $p = 0.74$, high D levels (≥ 20 ng/mL) 6 of 77 (7.8%), low D levels (<20ng/mL) 1 of 15 (6.7%), inverted to make $RR < 1$ favor high D levels (≥ 20 ng/mL), odds ratio converted to relative risk, multivariable logistic regression, outcome based on serum levels.

	<p>risk of mechanical ventilation, 56.9% lower, RR 0.43, $p = 0.22$, high D levels ($\geq 20\text{ng/mL}$) 8 of 15 (53.3%), low D levels ($< 20\text{ng/mL}$) 4 of 15 (26.7%), inverted to make $\text{RR} < 1$ favor high D levels ($\geq 20\text{ng/mL}$), odds ratio converted to relative risk, multivariable logistic regression, outcome based on serum levels.</p>
	<p>risk of ICU admission, 13.1% higher, RR 1.13, $p = 0.57$, high D levels ($\geq 20\text{ng/mL}$) 54 of 77 (70.1%), low D levels ($< 20\text{ng/mL}$) 9 of 15 (60.0%), inverted to make $\text{RR} < 1$ favor high D levels ($\geq 20\text{ng/mL}$), odds ratio converted to relative risk, multivariable logistic regression, outcome based on serum levels.</p>
<i>Pepkowitz</i> , 9/29/2020, retrospective, USA, preprint, 7 authors.	<p>risk of ICU admission, 55.8% lower, RR 0.44, $p = 0.01$, high D levels ($\geq 20\text{ng/mL}$) 9 of 24 (37.5%), low D levels ($< 20\text{ng/mL}$) 11 of 13 (84.6%), NNT 2.1, inverted to make $\text{RR} < 1$ favor high D levels ($\geq 20\text{ng/mL}$).</p>
<i>Pimental</i> , 5/31/2021, retrospective, Brazil, peer-reviewed, 3 authors.	<p>risk of death, 29.4% lower, RR 0.71, $p = 1.00$, high D levels 3 of 17 (17.6%), low D levels 2 of 8 (25.0%), NNT 14, $> 20\text{ng/mL}$.</p>
<i>Protas</i> , 4/6/2023, retrospective, Kazakhstan, peer-reviewed, survey, 6 authors, study period October 2022 - November 2022.	<p>risk of case, 76.6% lower, OR 0.23, $p = 0.06$, high D levels ($\geq 10\text{ng/mL}$) 68 of 88 (77.3%) cases, 29 of 31 (93.5%) controls, NNT 4.8, case control OR.</p>
	<p>risk of case, 46.2% lower, OR 0.54, $p = 0.17$, high D levels ($\geq 20\text{ng/mL}$) 50 of 88 (56.8%) cases, 22 of 31 (71.0%) controls, NNT 8.8, case control OR.</p>
<i>Putra</i> , 12/10/2021, retrospective, Indonesia, peer-reviewed, 3 authors, study period February 2020 - September 2020.	<p>risk of hospitalization, 25.6% lower, OR 0.74, $p = 0.59$, high D levels 9 of 31 (29.0%) cases, 11 of 31 (35.5%) controls, NNT 14, case control OR.</p>
<i>Rachman</i> , 4/13/2023, prospective, Indonesia, peer-reviewed, 4 authors, study period October 2021 - February 2022.	<p>risk of death, 94.8% lower, RR 0.05, $p = 0.04$, high D levels ($\geq 20\text{ng/mL}$) 0 of 45 (0.0%), low D levels ($< 20\text{ng/mL}$) 14 of 146 (9.6%), NNT 10, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm).</p>
	<p>risk of severe case, 77.6% lower, RR 0.22, $p = 0.01$, high D levels ($\geq 20\text{ng/mL}$) 2 of 45 (4.4%), low D levels ($< 20\text{ng/mL}$) 29 of 146 (19.9%), NNT 6.5.</p>
<i>Radujkovic</i> , 9/10/2020, prospective, Germany, peer-reviewed, 6 authors.	<p>risk of death, 93.2% lower, HR 0.07, $p = 0.001$, high D levels 144, low D levels 12, $> 30\text{nmol/L}$.</p>
	<p>risk of death/intubation, 84.0% lower, HR 0.16, $p < 0.001$, high D levels 144, low D levels 12, $> 30\text{nmol/L}$.</p>
<i>Ramirez-Sandoval</i> , 10/15/2021, retrospective, Mexico, peer-reviewed, 7 authors.	<p>risk of death, 31.5% lower, HR 0.68, $p < 0.001$, high D levels 2,337, low D levels 571, adjusted per study, inverted to make $\text{HR} < 1$ favor high D levels, $> 12.5\text{ng/mL}$, 30 day in-hospital mortality.</p>
	<p>hospitalization time, 22.2% lower, relative time 0.78, $p < 0.001$, high D levels 2,337, low D levels 571.</p>

<i>Ramos</i> , 11/15/2021, retrospective, Brazil, peer-reviewed, 11 authors.	risk of case, 45.7% lower, RR 0.54, $p = 0.16$, high D levels ($\geq 20\text{ng/mL}$) 4 of 9 (44.4%), low D levels ($< 20\text{ng/mL}$) 9 of 11 (81.8%), NNT 2.7.
<i>Ranjbar</i> , 11/29/2021, retrospective, Iran, peer-reviewed, 27 authors, study period 16 February, 2020 - 21 March, 2020.	risk of death, 41.9% lower, RR 0.58, $p = 0.07$, high D levels ($\geq 20\text{ng/mL}$) 16 of 163 (9.8%), low D levels ($< 20\text{ng/mL}$) 26 of 154 (16.9%), NNT 14.
<i>Reis</i> , 5/21/2021, prospective, Brazil, peer-reviewed, 19 authors.	risk of death, 23.0% lower, HR 0.77, $p = 0.82$, high D levels ($\geq 10\text{ng/mL}$) 198, low D levels ($< 10\text{ng/mL}$) 16, model 2, Cox proportional hazards.
	risk of mechanical ventilation, 45.0% higher, HR 1.45, $p = 0.77$, high D levels ($\geq 10\text{ng/mL}$) 198, low D levels ($< 10\text{ng/mL}$) 16, adjusted per study, model 2, multivariable, Cox proportional hazards.
	risk of no hospital discharge, 33.3% lower, HR 0.67, $p = 0.18$, high D levels ($\geq 10\text{ng/mL}$) 198, low D levels ($< 10\text{ng/mL}$) 16, adjusted per study, inverted to make $\text{HR} < 1$ favor high D levels ($\geq 10\text{ng/mL}$), model 2, multivariable, Cox proportional hazards.
	hospitalization time, 22.2% lower, relative time 0.78, $p = 0.06$, high D levels ($\geq 10\text{ng/mL}$) 191, low D levels ($< 10\text{ng/mL}$) 15, model 2.
<i>Reyes Pérez</i> , 4/30/2020, retrospective, Mexico, peer-reviewed, 5 authors, excluded in exclusion analyses: unadjusted results with no group details.	risk of death, 61.7% lower, RR 0.38, $p = 0.006$, high D levels ($\geq 8\text{ng/mL}$) 21 of 137 (15.3%), low D levels ($< 8\text{ng/mL}$) 14 of 35 (40.0%), NNT 4.1, inverted to make $\text{RR} < 1$ favor high D levels ($\geq 8\text{ng/mL}$), odds ratio converted to relative risk.
<i>Ribeiro</i> , 8/5/2021, retrospective, Brazil, peer-reviewed, 8 authors.	risk of case, 50.5% lower, OR 0.50, $p = 0.01$, inverted to make $\text{OR} < 1$ favor high D levels, $> 30\text{ng/mL}$, multivariate, RR approximated with OR.
<i>Ricci</i> , 3/3/2021, retrospective, Italy, peer-reviewed, 15 authors.	risk of death, 87.6% lower, RR 0.12, $p = 0.07$, high D levels 0 of 30 (0.0%), low D levels 3 of 22 (13.6%), NNT 7.3, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm), $> 10\text{ ng/mL}$.
<i>Ritsinger</i> , 4/28/2023, retrospective, Sweden, peer-reviewed, mean age 79.8, 8 authors, study period 1 January, 2020 - 9 September, 2021.	risk of death, 9.1% lower, HR 0.91, $p < 0.001$, high D levels ($\geq 50\text{nmol/L}$) 37,972, low D levels ($< 50\text{nmol/L}$) 6,894, inverted to make $\text{HR} < 1$ favor high D levels ($\geq 50\text{nmol/L}$).
<i>Rodríguez-Vidales</i> , 2/24/2022, retrospective, Mexico, peer-reviewed, 8 authors, study period March 2020 - September 2020.	risk of severe case, 38.9% lower, RR 0.61, $p = 0.05$, high D levels ($\geq 10\text{ng/mL}$) 89 of 265 (33.6%), low D levels ($< 10\text{ng/mL}$) 27 of 32 (84.4%), NNT 2.0, adjusted per study, inverted to make $\text{RR} < 1$ favor high D levels ($\geq 10\text{ng/mL}$), odds ratio converted to relative risk, multivariable.
<i>Sanamandra</i> , 4/30/2023, prospective, India, peer-reviewed, 6 authors, study period August 2020 - March 2021.	risk of death, 20.9% lower, OR 0.79, $p = 0.67$, high D levels ($\geq 10\text{ng/mL}$) 155, low D levels ($< 10\text{ng/mL}$) 45, inverted to make $\text{OR} < 1$ favor high D levels ($\geq 10\text{ng/mL}$), RR approximated with OR.
	risk of mechanical ventilation, 15.3% lower, OR 0.85, $p = 0.73$, high D levels ($\geq 10\text{ng/mL}$) 155, low D levels ($< 10\text{ng/mL}$) 45, inverted to make $\text{OR} < 1$ favor high D levels ($\geq 10\text{ng/mL}$), RR

	approximated with OR.
	risk of severe case, 434.8% higher, OR 5.35, $p = 0.12$, high D levels ($\geq 10\text{ng/mL}$) 155, low D levels ($< 10\text{ng/mL}$) 45, inverted to make $\text{OR} < 1$ favor high D levels ($\geq 10\text{ng/mL}$), RR approximated with OR.
<i>Sanson</i> , 2/19/2022, prospective, Italy, peer-reviewed, 13 authors, study period March 2020 - September 2020, excluded in exclusion analyses: unadjusted results with no group details.	NIV/IMV/death, 64.0% lower, RR 0.36, $p = 0.03$, high D levels ($\geq 30\text{ng/mL}$) 2 of 9 (22.2%), low D levels ($< 30\text{ng/mL}$) 37 of 60 (61.7%), NNT 2.5.
<i>Saponaro</i> , 1/24/2022, retrospective, Italy, peer-reviewed, 13 authors, study period March 2020 - May 2020.	risk of ARDS, 36.5% lower, RR 0.64, $p = 0.43$, high D levels ($\geq 20\text{ng/mL}$) 5 of 32 (15.6%), low D levels ($< 20\text{ng/mL}$) 15 of 61 (24.6%), NNT 11, severe ARDS.
<i>Savitri</i> , 5/8/2021, retrospective, Indonesia, peer-reviewed, 5 authors.	risk of symptomatic case, 88.0% lower, RR 0.12, $p < 0.001$, high D levels 3 of 25 (12.0%), low D levels 17 of 17 (100.0%), NNT 1.1, $> 20\text{ng/mL}$.
<i>Schmidt</i> , 3/22/2023, prospective, Poland, peer-reviewed, 4 authors, study period 4 February, 2021 - 31 December, 2021.	risk of death, 85.5% lower, OR 0.14, $p = 0.003$, cutoff 27ng/mL, inverted to make $\text{OR} < 1$ favor high D levels ($\geq 27\text{ng/mL}$), RR approximated with OR.
<i>Seal</i> , 1/1/2022, retrospective, USA, peer-reviewed, 6 authors.	risk of death, 45.1% lower, RR 0.55, $p = 0.001$, adjusted per study, inverted to make $\text{RR} < 1$ favor high D levels, 60ng/mL vs. 15 ng/mL.
	risk of death, 40.5% lower, RR 0.60, $p = 0.001$, adjusted per study, inverted to make $\text{RR} < 1$ favor high D levels, 50ng/mL vs. 15 ng/mL.
	risk of death, 34.6% lower, RR 0.65, $p = 0.001$, adjusted per study, inverted to make $\text{RR} < 1$ favor high D levels, 40ng/mL vs. 15 ng/mL.
	risk of death, 25.9% lower, RR 0.74, $p = 0.001$, adjusted per study, inverted to make $\text{RR} < 1$ favor high D levels, 30ng/mL vs. 15 ng/mL.
	risk of death, 20.0% lower, RR 0.80, $p = 0.001$, adjusted per study, inverted to make $\text{RR} < 1$ favor high D levels, 25ng/mL vs. 15 ng/mL.
	risk of death, 11.5% lower, RR 0.88, $p = 0.001$, adjusted per study, inverted to make $\text{RR} < 1$ favor high D levels, 20ng/mL vs. 15 ng/mL.
	risk of hospitalization, 22.5% lower, RR 0.78, $p = 0.01$, adjusted per study, inverted to make $\text{RR} < 1$ favor high D levels, 60ng/mL vs. 15 ng/mL.
	risk of hospitalization, 20.0% lower, RR 0.80, $p = 0.009$, adjusted per study, inverted to make $\text{RR} < 1$ favor high D levels, 50ng/mL vs. 15 ng/mL.

	risk of hospitalization, 16.7% lower, RR 0.83, $p = 0.007$, adjusted per study, inverted to make $RR < 1$ favor high D levels, 40ng/mL vs. 15 ng/mL.
	risk of hospitalization, 12.3% lower, RR 0.88, $p = 0.008$, adjusted per study, inverted to make $RR < 1$ favor high D levels, 30ng/mL vs. 15 ng/mL.
	risk of hospitalization, 9.1% lower, RR 0.91, $p = 0.01$, adjusted per study, inverted to make $RR < 1$ favor high D levels, 25ng/mL vs. 15 ng/mL.
	risk of hospitalization, 4.8% lower, RR 0.95, $p = 0.02$, adjusted per study, inverted to make $RR < 1$ favor high D levels, 20ng/mL vs. 15 ng/mL.
<i>Seven</i> , 11/23/2021, prospective, Turkey, peer-reviewed, 6 authors, study period September 2020 - November 2020.	risk of severe disease or poor prognostic factor, 46.5% lower, RR 0.53, $p = 0.006$, cutoff 14.5ng/ml, inverted to make $RR < 1$ favor high D levels (≥ 14.5 ng/ml).
<i>Sinaci</i> , 8/11/2021, retrospective, Turkey, peer-reviewed, 10 authors, dosage not specified.	risk of moderate/severe case, 79.5% lower, RR 0.21, $p < 0.001$, high D levels (≥ 10 ng/mL) 8 of 100 (8.0%), low D levels (< 10 ng/mL) 23 of 59 (39.0%), NNT 3.2, outcome based on serum levels.
	risk of case, 59.9% lower, RR 0.40, $p < 0.001$, high D levels (≥ 10 ng/mL) 100 of 397 (25.2%), low D levels (< 10 ng/mL) 59 of 94 (62.8%), NNT 2.7, outcome based on serum levels.
<i>Siuka</i> , 3/9/2023, prospective, Slovenia, peer-reviewed, 7 authors, study period December 2020 - December 2021.	risk of death, 55.9% lower, RR 0.44, $p = 0.24$, high D levels (≥ 30 nmol/L) 10 of 255 (3.9%), low D levels (< 30 nmol/L) 4 of 45 (8.9%), NNT 20.
	risk of ICU admission, 58.8% higher, RR 1.59, $p = 0.59$, high D levels (≥ 30 nmol/L) 27 of 255 (10.6%), low D levels (< 30 nmol/L) 3 of 45 (6.7%).
	risk of severe case, 61.0% higher, RR 1.61, $p = 0.009$, high D levels (≥ 30 nmol/L) 146 of 255 (57.3%), low D levels (< 30 nmol/L) 16 of 45 (35.6%).
<i>Subramanian</i> , 1/31/2022, prospective, United Kingdom, peer-reviewed, 16 authors, dosage not specified.	risk of death, 49.7% lower, RR 0.50, $p = 0.02$, high D levels 16 of 115 (13.9%), low D levels 33 of 118 (28.0%), NNT 7.1, adjusted per study, inverted to make $RR < 1$ favor high D levels, odds ratio converted to relative risk, 50-74 nmol/L vs. < 25 nmol/L, multivariable, outcome based on serum levels.
	risk of death, 39.7% lower, RR 0.60, $p = 0.07$, high D levels 16 of 115 (13.9%), low D levels 38 of 157 (24.2%), NNT 9.7, adjusted per study, inverted to make $RR < 1$ favor high D levels, odds ratio converted to relative risk, 50-74 nmol/L vs. 25-49nmol/L, multivariable, outcome based on serum levels.
<i>Sulli</i> , 2/24/2021, retrospective, Italy, peer-reviewed, 10 authors, dosage not specified.	risk of case, 79.2% lower, OR 0.21, $p < 0.001$, high D levels 28 of 65 (43.1%) cases, 51 of 65 (78.5%) controls, NNT 2.7, case control OR, > 10 ng/mL.

<i>Susianti</i> , 2/12/2021, retrospective, Indonesia, peer-reviewed, 8 authors.	risk of death, 91.5% lower, RR 0.09, $p = 0.32$, high D levels 0 of 8 (0.0%), low D levels 9 of 42 (21.4%), NNT 4.7, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm), >49.92 nmol/L.
	risk of ICU admission, 90.5% lower, RR 0.10, $p = 0.32$, high D levels 0 of 8 (0.0%), low D levels 8 of 42 (19.0%), NNT 5.2, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm), >49.92 nmol/L.
	risk of progression, 81.5% lower, OR 0.19, $p = 0.04$, high D levels 8, low D levels 42, inverted to make OR<1 favor high D levels, ISTH DIC>=5, >49.92 nmol/L, bivariate, RR approximated with OR.
	risk of progression, 44.4% lower, OR 0.56, $p = 0.03$, high D levels 8, low D levels 42, inverted to make OR<1 favor high D levels, increased D-dimer >2 mg/L, >49.92 nmol/L, multivariate, RR approximated with OR.
<i>Szeto</i> , 12/30/2020, retrospective, USA, peer-reviewed, 7 authors.	risk of death, 5.6% higher, RR 1.06, $p = 1.00$, high D levels 14 of 58 (24.1%), low D levels 8 of 35 (22.9%).
	risk of mechanical ventilation, 39.7% lower, RR 0.60, $p = 0.21$, high D levels 10 of 58 (17.2%), low D levels 10 of 35 (28.6%), NNT 8.8.
	risk of no hospital discharge, 26.7% higher, RR 1.27, $p = 0.50$, high D levels 21 of 58 (36.2%), low D levels 10 of 35 (28.6%).
<i>Sánchez-Zuno (B)</i> , 5/28/2021, prospective, Mexico, peer-reviewed, 12 authors, dosage 10,000IU days 1-14.	risk of severe case, 5.6% lower, RR 0.94, $p = 1.00$, high D levels 4 of 8 (50.0%), low D levels 18 of 34 (52.9%), NNT 34, >30ng/mL, >4 symptoms.
<i>Tallon</i> , 11/15/2022, retrospective, USA, peer-reviewed, 17 authors.	risk of hospitalization, 41.5% lower, OR 0.58, $p < 0.001$, high D levels (≥ 30 ng/mL) 113,143, low D levels (<30ng/mL) 3,227, adjusted per study, inverted to make OR<1 favor high D levels (≥ 30 ng/mL), RR approximated with OR.
<i>Tan</i> , 2/27/2023, retrospective, Philippines, peer-reviewed, 3 authors.	risk of progression, 71.5% lower, RR 0.29, $p = 0.04$, high D levels (≥ 30 ng/mL) 7 of 38 (18.4%), low D levels (<20ng/mL) 18 of 34 (52.9%), NNT 2.9, adjusted per study, inverted to make RR<1 favor high D levels (≥ 30 ng/mL), odds ratio converted to relative risk, combined mortality and morbidity, multivariable.
	risk of death, 91.1% lower, RR 0.09, $p = 0.002$, high D levels (≥ 30 ng/mL) 1 of 38 (2.6%), low D levels (<20ng/mL) 10 of 34 (29.4%), NNT 3.7, unadjusted.
	risk of ICU admission, 82.1% lower, RR 0.18, $p = 0.010$, high D levels (≥ 30 ng/mL) 2 of 38 (5.3%), low D levels (<20ng/mL) 10 of 34 (29.4%), NNT 4.1, unadjusted.
<i>Tehrani</i> , 1/25/2021, retrospective, Iran, peer-reviewed, 5 authors.	risk of death, 47.5% lower, RR 0.52, $p = 0.07$, high D levels 34 of 180 (18.9%), low D levels 9 of 25 (36.0%), NNT 5.8, >10ng/mL.

<p><i>Tomasa-Irriguable</i>, 10/26/2020, retrospective, Spain, peer-reviewed, 7 authors, study period March 2020 - May 2020.</p>	<p>risk of mechanical ventilation, 35.0% lower, RR 0.65, $p = 0.21$, high D levels 15 of 27 (55.6%), low D levels 18 of 78 (23.1%), adjusted per study, inverted to make $RR < 1$ favor high D levels, odds ratio converted to relative risk, ≥ 20 ng/mL, bivariate logistic regression.</p>
	<p>risk of ICU admission, 16.9% lower, RR 0.83, $p = 0.58$, high D levels 11 of 27 (40.7%), low D levels 17 of 78 (21.8%), adjusted per study, inverted to make $RR < 1$ favor high D levels, odds ratio converted to relative risk, ≥ 20 ng/mL, bivariate logistic regression.</p>
<p><i>Topan</i>, 2/28/2023, retrospective, Romania, peer-reviewed, survey, 6 authors, study period April 2020 - May 2022.</p>	<p>risk of death, 30.6% lower, RR 0.69, $p = 0.02$, high D levels (≥ 20ng/mL) 61 of 1,148 (5.3%), low D levels (< 20ng/mL) 118 of 1,194 (9.9%), adjusted per study, inverted to make $RR < 1$ favor high D levels (≥ 20ng/mL), odds ratio converted to relative risk, multivariable.</p>
	<p>risk of severe case, 10.9% lower, RR 0.89, $p = 0.02$, high D levels (≥ 20ng/mL) 432 of 1,148 (37.6%), low D levels (< 20ng/mL) 560 of 1,194 (46.9%), NNT 11, adjusted per study, inverted to make $RR < 1$ favor high D levels (≥ 20ng/mL), odds ratio converted to relative risk, severe/critical case, multivariable.</p>
<p><i>Umay</i>, 7/26/2023, retrospective, Turkey, peer-reviewed, 4 authors, study period 1 March, 2020 - 31 January, 2021.</p>	<p>hospitalization time, 13.5% lower, relative time 0.87, $p = 0.33$, high D levels 374, low D levels 39.</p>
<p><i>Vanegas-Cedillo</i>, 3/14/2021, retrospective, Mexico, peer-reviewed, 15 authors.</p>	<p>risk of death, 52.6% lower, RR 0.47, $p = 0.006$, high D levels (≥ 12ng/mL) 95 of 494 (19.2%), low D levels (< 12ng/mL) 21 of 57 (36.8%), NNT 5.7, adjusted per study, inverted to make $RR < 1$ favor high D levels (≥ 12ng/mL).</p>
<p><i>Vasheghani</i>, 1/18/2021, retrospective, Iran, preprint, 6 authors, dosage not specified.</p>	<p>risk of ICU admission, 63.8% lower, RR 0.36, $p = 0.009$, high D levels 13 of 185 (7.0%), low D levels 53 of 323 (16.4%), NNT 11, adjusted per study, inverted to make $RR < 1$ favor high D levels, vitamin D levels > 30ng/mL.</p>
<p><i>Vassiliou (B)</i>, 12/9/2020, prospective, Greece, peer-reviewed, 6 authors.</p>	<p>risk of death, 90.9% lower, RR 0.09, $p = 0.04$, high D levels 0 of 15 (0.0%), low D levels 5 of 15 (33.3%), NNT 3.0, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm), > 15.2ng/mL.</p>
<p><i>Voelkle</i>, 4/30/2022, prospective, Switzerland, peer-reviewed, median age 67.0, 9 authors, study period 17 March, 2020 - 30 April, 2020.</p>	<p>risk of death/ICU, 23.4% lower, RR 0.77, $p = 0.55$, high D levels 8 of 34 (23.5%), low D levels 7 of 23 (30.4%), NNT 14, adjusted per study, inverted to make $RR < 1$ favor high D levels, odds ratio converted to relative risk.</p>
<p><i>Vásquez-Procopio</i>, 12/2/2022, retrospective, Mexico, peer-reviewed, 12 authors.</p>	<p>risk of severe case, 82.8% lower, OR 0.17, $p = 0.04$, high D levels (≥ 20ng/mL) 111, low D levels (< 20ng/mL) 54, adjusted per study, inverted to make $OR < 1$ favor high D levels (≥ 20ng/mL), multivariable, RR approximated with OR.</p>
<p><i>Walk</i>, 11/9/2020, retrospective, Netherlands, preprint, 5 authors.</p>	<p>risk of death/intubation, 0.4% higher, RR 1.00, $p = 1.00$, high D levels 48 of 110 (43.6%), low D levels 10 of 23 (43.5%), > 25nmol/L.</p>

<i>Wang</i> , 3/29/2023, prospective, China, preprint, median age 36.5, 23 authors, study period 18 December, 2022 - 20 February, 2023, dosage 200,000IU days 1, 14, trial NCT05673980 (history).	risk of case, 22.7% lower, RR 0.77, $p = 0.19$, high D levels ($\geq 30\text{ng/ml}$) 20 of 44 (45.5%), low D levels ($< 20\text{ng/ml}$) 50 of 85 (58.8%), NNT 7.5, outcome based on serum levels.
<i>Wani</i> , 6/1/2023, retrospective, India, peer-reviewed, 5 authors, excluded in exclusion analyses: unadjusted results with no group details.	risk of severe case, 72.2% lower, OR 0.28, $p = 0.007$, high D levels ($\geq 28\text{ng/mL}$) 66, low D levels ($< 28\text{ng/mL}$) 170, inverted to make $\text{OR} < 1$ favor high D levels ($\geq 28\text{ng/mL}$), RR approximated with OR.
<i>Ye</i> , 10/13/2020, retrospective, China, peer-reviewed, 18 authors.	risk of severe/critical COVID-19, 93.4% lower, RR 0.07, $p = 0.03$, high D levels 2 of 36 (5.6%), low D levels 8 of 26 (30.8%), NNT 4.0, adjusted per study, inverted to make $\text{RR} < 1$ favor high D levels, $> 50\text{nmol/L}$.
<i>Yilmaz</i> , 10/5/2020, retrospective, Turkey, peer-reviewed, 2 authors.	risk of severe case, 73.4% lower, RR 0.27, $p = 1.00$, high D levels 0 of 11 (0.0%), low D levels 2 of 29 (6.9%), NNT 14, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm), $> 20\text{ng/ml}$.
	risk of moderate or severe case, 41.4% lower, RR 0.59, $p = 0.69$, high D levels 2 of 11 (18.2%), low D levels 9 of 29 (31.0%), NNT 7.8, $> 20\text{ng/ml}$.
<i>Zafar</i> , 9/6/2021, retrospective, United Kingdom, peer-reviewed, median age 68.0, 37 authors.	risk of death, 42.9% higher, RR 1.43, $p = 0.71$, high D levels ($\geq 25\text{nmol/L}$) 12 of 42 (28.6%), low D levels ($< 25\text{nmol/L}$) 2 of 10 (20.0%), COVID+ patients.
	risk of death, 6.0% lower, OR 0.94, $p = 0.68$, high D levels 42, low D levels 10, COVID+ patients, RR approximated with OR.
<i>Zeidan</i> , 9/9/2022, prospective, Egypt, peer-reviewed, median age 11.4, 38 authors.	risk of hospitalization, 61.5% lower, OR 0.38, $p = 0.002$, cutoff 20ng/mL, adjusted per study, inverted to make $\text{OR} < 1$ favor high D levels ($\geq 20\text{ng/mL}$), case control OR, multivariable.
<i>Zelzer</i> , 6/22/2021, retrospective, Austria, peer-reviewed, 7 authors.	risk of death, 46.4% lower, RR 0.54, $p = 0.08$, high D levels 24 of 121 (19.8%), low D levels 10 of 27 (37.0%), NNT 5.8, $> 30\text{nmol/L}$.
<i>Zidrou</i> , 2/19/2022, retrospective, Greece, peer-reviewed, 6 authors, study period August 2020 - October 2020.	risk of death, 26.4% lower, RR 0.74, $p = 1.00$, high D levels ($\geq 20\text{ng/ml}$) 2 of 25 (8.0%), low D levels ($< 20\text{ng/ml}$) 5 of 46 (10.9%), NNT 35.
	radiographic changes, 18.2% lower, RR 0.82, $p = 0.26$, high D levels ($\geq 20\text{ng/ml}$) 16 of 25 (64.0%), low D levels ($< 20\text{ng/ml}$) 36 of 46 (78.3%), NNT 7.0.
	hospitalization time, 37.7% lower, relative time 0.62, $p = 0.16$, high D levels ($\geq 20\text{ng/ml}$) 25, low D levels ($< 20\text{ng/ml}$) 46.
<i>Álvarez</i> , 10/28/2022, retrospective, Spain, preprint, 1 author, study period March 2020 - March 2021, excluded in exclusion analyses: unadjusted results with no group details.	risk of death, 38.8% lower, RR 0.61, $p < 0.001$, high D levels 4,871 of 33,673 (14.5%), low D levels 611 of 2,588 (23.6%), NNT 11, inverted to make $\text{RR} < 1$ favor high D levels, odds ratio converted to relative risk.

	risk of ICU admission, 54.7% lower, RR 0.45, $p < 0.001$, high D levels 289 of 33,673 (0.9%), low D levels 49 of 2,588 (1.9%), NNT 97, inverted to make $RR < 1$ favor high D levels, odds ratio converted to relative risk.
	risk of hospitalization, 43.0% lower, RR 0.57, $p < 0.001$, high D levels 8,905 of 33,673 (26.4%), low D levels 1,202 of 2,588 (46.4%), NNT 5.0, inverted to make $RR < 1$ favor high D levels, odds ratio converted to relative risk.
<i>Ünsal</i> , 4/5/2021, retrospective, Turkey, peer-reviewed, 10 authors.	risk of death, 80.6% lower, RR 0.19, $p = 0.23$, high D levels 0 of 29 (0.0%), low D levels 2 of 27 (7.4%), NNT 14, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm), $\geq 20\text{ng/mL}$.
	risk of oxygen therapy, 73.4% lower, RR 0.27, $p = 0.07$, high D levels 2 of 29 (6.9%), low D levels 7 of 27 (25.9%), NNT 5.3, $\geq 20\text{ng/mL}$.
<i>Şengül</i> , 12/31/2022, retrospective, Turkey, peer-reviewed, 4 authors, study period March 2020 - December 2021, dosage not specified.	risk of case, 75.6% lower, OR 0.24, $p < 0.001$, high D levels ($\geq 20\text{ng/mL}$) 7 of 54 (13.0%) cases, 100 of 264 (37.9%) controls, NNT 6.4, case control OR, outcome based on serum levels.

Early treatment

Effect extraction follows pre-specified rules as detailed above and gives priority to more serious outcomes. Only the first (most serious) outcome is used in pooled analysis, which may differ from the effect a paper focuses on. Other outcomes are used in outcome specific analyses.

<i>Annweiler</i> , 11/2/2020, retrospective, France, peer-reviewed, 7 authors, dosage 80,000IU single dose.	risk of death, 63.0% lower, RR 0.37, $p = 0.28$, treatment 3 of 16 (18.8%), control 10 of 32 (31.2%), NNT 8.0, adjusted per study, supplementation after diagnosis.
<i>Annweiler (B)</i> , 10/13/2020, retrospective, France, peer-reviewed, mean age 87.7, 6 authors, dosage 80,000IU single dose, 80,000IU either in the week following the suspicion or diagnosis of COVID-19, or during the previous month.	risk of death, 89.0% lower, RR 0.11, $p = 0.002$, treatment 10 of 57 (17.5%), control 5 of 9 (55.6%), NNT 2.6, adjusted per study.
<i>Asimi</i> , 5/22/2021, retrospective, Bosnia and Herzegovina, preprint, 3 authors, dosage 2,000IU daily, this trial uses multiple treatments in the treatment arm (combined with zinc and selenium) - results of individual treatments may vary, excluded in exclusion analyses: excessive unadjusted differences between groups.	risk of mechanical ventilation, 97.4% lower, RR 0.03, $p < 0.001$, treatment 0 of 270 (0.0%), control 9 of 86 (10.5%), NNT 9.6, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm), unadjusted.
	risk of hospitalization, 99.0% lower, RR 0.010, $p < 0.001$, treatment 0 of 270 (0.0%), control 24 of 86 (27.9%), NNT 3.6, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm), unadjusted.
	risk of severe case, 99.5% lower, RR 0.005, $p < 0.001$, treatment 0 of 270 (0.0%), control 51 of 86 (59.3%), NNT 1.7, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm), unadjusted.

<i>Boukef</i> , 2/28/2023, Double Blind Randomized Controlled Trial, placebo-controlled, Tunisia, trial NCT05670444 (history).	150 patient RCT with results missing over 7 months.
<i>Burahee</i> , 2/17/2021, retrospective, United Kingdom, peer-reviewed, 4 authors, dosage 100,000IU days 1-4, additional 200000IU over four weeks if serum level insufficient.	risk of death, 93.3% lower, RR 0.07, $p = 0.01$, treatment 0 of 12 (0.0%), control 2 of 2 (100.0%), NNT 1.0, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm).
<i>Din Ujjan</i> , 1/18/2023, Randomized Controlled Trial, Pakistan, peer-reviewed, 6 authors, study period 21 September, 2021 - 21 January, 2022, dosage 360IU days 1-14, this trial uses multiple treatments in the treatment arm (combined with curcumin and quercetin) - results of individual treatments may vary, trial NCT04603690 (history), excluded in exclusion analyses: based on dosages and previous research, combined treatments may contribute more to the effect seen.	risk of no recovery, 28.6% lower, RR 0.71, $p = 0.11$, treatment 15 of 25 (60.0%), control 21 of 25 (84.0%), NNT 4.2, no symptoms, day 7.
	risk of no recovery, 71.4% lower, RR 0.29, $p < 0.001$, treatment 6 of 25 (24.0%), control 21 of 25 (84.0%), NNT 1.7, ≤ 1 symptom, day 7.
	risk of no recovery, 76.9% lower, RR 0.23, $p = 0.005$, treatment 3 of 25 (12.0%), control 13 of 25 (52.0%), NNT 2.5, ≤ 2 symptoms, day 7.
	risk of no recovery, 85.7% lower, RR 0.14, $p = 0.23$, treatment 0 of 25 (0.0%), control 3 of 25 (12.0%), NNT 8.3, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm), ≤ 3 symptoms, day 7.
	risk of no viral clearance, 90.9% lower, RR 0.09, $p = 0.05$, treatment 0 of 25 (0.0%), control 5 of 25 (20.0%), NNT 5.0, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm), day 14.
	risk of no viral clearance, 73.7% lower, RR 0.26, $p < 0.001$, treatment 5 of 25 (20.0%), control 19 of 25 (76.0%), NNT 1.8, day 7.
<i>Efird</i> , 12/31/2021, retrospective, USA, peer-reviewed, 10 authors, study period 1 March, 2020 - 10 September, 2020, dosage varies.	risk of death, 48.9% lower, RR 0.51, $p = 0.10$, treatment 11 of 544 (2.0%), control 413 of 15,794 (2.6%), adjusted per study, non-hospitalized patients, vitamin D + no corticosteroids vs. no vitamin D + no corticosteroids.
	risk of death, 54.5% lower, RR 0.45, $p = 0.02$, treatment 11 of 192 (5.7%), control 553 of 4,340 (12.7%), NNT 14, adjusted per study, hospitalized patients, vitamin D + no corticosteroids vs. no vitamin D + no corticosteroids.
<i>Hunt</i> , 6/29/2022, retrospective, USA, peer-reviewed, 8 authors, study period 1 March, 2020 - 10 September, 2020, dosage not specified.	risk of death, 47.0% lower, RR 0.53, $p < 0.001$, treatment 43 of 1,019 (4.2%), control 1,569 of 25,489 (6.2%), adjusted per study, day 30.
<i>Khan</i> , 5/1/2022, Randomized Controlled Trial, Pakistan, peer-reviewed, 7 authors, study period 2 September, 2021 - 28 November, 2021, dosage 360IU days 1-14, this trial uses multiple treatments in the treatment arm (combined with curcumin and quercetin) - results of individual treatments may vary, trial NCT05130671 (history), excluded in	risk of no recovery, 33.3% lower, RR 0.67, $p = 0.15$, treatment 10 of 25 (40.0%), control 15 of 25 (60.0%), NNT 5.0.
	relative CRP reduction, 39.1% better, RR 0.61, $p = 0.006$, treatment 25, control 25.

exclusion analyses: based on dosages and previous research, combined treatments may contribute more to the effect seen.	risk of no viral clearance, 50.0% lower, RR 0.50, $p = 0.009$, treatment 10 of 25 (40.0%), control 20 of 25 (80.0%), NNT 2.5.
<p><i>Said</i>, 11/8/2022, Randomized Controlled Trial, Egypt, peer-reviewed, 5 authors, study period 21 July, 2021 - 30 December, 2021, dosage 2,000IU daily, trial NCT04981743 (history).</p>	risk of no recovery, 42.0% lower, OR 0.58, $p = 0.57$, treatment 30, control 30, adjusted per study, multivariable, dyspnea, RR approximated with OR.
	risk of no recovery, 89.0% lower, OR 0.11, $p = 0.01$, treatment 30, control 30, adjusted per study, vitamin D and nigella sativa, multivariable, dyspnea, RR approximated with OR.
	risk of no recovery, 52.0% lower, OR 0.48, $p = 0.16$, treatment 30, control 30, adjusted per study, multivariable, cough, RR approximated with OR.
	risk of no recovery, 77.0% lower, OR 0.23, $p = 0.01$, treatment 30, control 30, adjusted per study, vitamin D and nigella sativa, multivariable, cough, RR approximated with OR.
	risk of no recovery, 56.0% lower, OR 0.44, $p = 0.20$, treatment 30, control 30, adjusted per study, multivariable, fatigue, RR approximated with OR.
	risk of no recovery, 90.0% lower, OR 0.10, $p < 0.001$, treatment 30, control 30, adjusted per study, vitamin D and nigella sativa, multivariable, fatigue, RR approximated with OR.
	risk of no recovery, 33.0% lower, OR 0.67, $p = 0.67$, treatment 30, control 30, adjusted per study, multivariable, smell, RR approximated with OR.
	risk of no recovery, 67.0% lower, OR 0.33, $p = 0.23$, treatment 30, control 30, adjusted per study, vitamin D and nigella sativa, multivariable, smell, RR approximated with OR.
	risk of no recovery, 25.0% higher, OR 1.25, $p = 0.79$, treatment 30, control 30, adjusted per study, multivariable, taste, RR approximated with OR.
	risk of no recovery, 58.0% lower, OR 0.42, $p = 0.28$, treatment 30, control 30, adjusted per study, vitamin D and nigella sativa, multivariable, taste, RR approximated with OR.
	risk of no recovery, 56.0% lower, OR 0.44, $p = 0.36$, treatment 30, control 30, adjusted per study, multivariable, sore throat, RR approximated with OR.
	risk of no recovery, 86.0% lower, OR 0.14, $p = 0.03$, treatment 30, control 30, adjusted per study, vitamin D and nigella sativa, multivariable, sore throat, RR approximated with OR.
	risk of no recovery, 175.0% higher, OR 2.75, $p = 0.13$, treatment 30, control 30, adjusted per study, multivariable, headache, RR approximated with OR.

	risk of no recovery, 56.0% lower, OR 0.44, $p = 0.21$, treatment 30, control 30, adjusted per study, vitamin D and nigella sativa, multivariable, headache, RR approximated with OR.
	risk of no recovery, 87.0% lower, OR 0.13, $p = 0.05$, treatment 30, control 30, adjusted per study, multivariable, diarrhea, RR approximated with OR.
	risk of no recovery, 90.0% lower, OR 0.10, $p = 0.03$, treatment 30, control 30, adjusted per study, vitamin D and nigella sativa, multivariable, diarrhea, RR approximated with OR.
	risk of no viral clearance, 49.0% lower, OR 0.51, $p = 0.20$, treatment 30, control 30, day 14, RR approximated with OR.
	risk of no viral clearance, 23.0% lower, OR 0.77, $p = 0.74$, treatment 30, control 30, day 7, RR approximated with OR.
	risk of no viral clearance, 91.0% lower, OR 0.09, $p < 0.001$, treatment 30, control 30, vitamin D and nigella sativa, day 14, RR approximated with OR.
	risk of no viral clearance, 87.0% lower, OR 0.13, $p = 0.003$, treatment 30, control 30, vitamin D and nigella sativa, day 7, RR approximated with OR.
<i>Sánchez-Zuno</i> , 5/28/2021, Randomized Controlled Trial, Mexico, peer-reviewed, 12 authors, dosage 10,000IU days 1-14.	risk of severe case, 89.4% lower, RR 0.11, $p = 0.04$, treatment 0 of 22 (0.0%), control 4 of 20 (20.0%), NNT 5.0, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm), risk of >3 symptoms at day 14.
	risk of no recovery, 80.8% lower, RR 0.19, $p = 0.22$, treatment 0 of 22 (0.0%), control 2 of 20 (10.0%), NNT 10.0, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm), risk of fever at day 14, Table S1.
<i>Valecha</i> , 4/26/2022, prospective, India, peer-reviewed, 1 author, average treatment delay 3.7 days, this trial uses multiple treatments in the treatment arm (combined with magnesium and vitamin B12) - results of individual treatments may vary.	risk of ICU admission, 86.8% lower, RR 0.13, $p = 0.09$, treatment 0 of 30 (0.0%), control 3 of 25 (12.0%), NNT 8.3, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm).
	hospitalization time, 38.5% lower, relative time 0.62, $p < 0.001$, treatment mean 11.2 (± 2.8) $n=30$, control mean 18.2 (± 1.21) $n=25$.

Late treatment

Effect extraction follows pre-specified rules as detailed above and gives priority to more serious outcomes. Only the first (most serious) outcome is used in pooled analysis, which may differ from the effect a paper focuses on. Other outcomes are used in outcome specific analyses.

<p><i>Al Sulaiman</i>, 8/14/2023, retrospective, Saudi Arabia, peer-reviewed, 25 authors, study period March 2020 - July 2021, dosage not specified, excluded in exclusion analyses: very late stage study using cholecalciferol instead of calcifediol or calcitriol.</p>	<p>risk of death, 22.0% higher, HR 1.22, $p = 0.25$, treatment 72 of 144 (50.0%), control 62 of 144 (43.1%).</p>
	<p>risk of mechanical ventilation, 27.0% higher, OR 1.27, $p = 0.046$, treatment 144, control 144, RR approximated with OR.</p>
	<p>risk of ICU admission, 17.0% higher, OR 1.17, $p = 0.07$, treatment 144, control 144, RR approximated with OR.</p>
	<p>risk of hospitalization, no change, OR 1.00, $p = 1.00$, treatment 144, control 144, RR approximated with OR.</p>
<p><i>Alcala-Diaz</i>, 5/21/2021, retrospective, Spain, peer-reviewed, 17 authors, dosage calcifediol 0.5mg day 1, 0.27mg day 3, 0.27mg day 7, 0.27mg day 14, 0.27mg day 21, 0.27mg day 28.</p>	<p>risk of death, 80.8% lower, RR 0.19, $p = 0.04$, treatment 4 of 79 (5.1%), control 90 of 458 (19.7%), NNT 6.9, adjusted per study, odds ratio converted to relative risk, day 30, multivariate logistic regression.</p>
<p><i>Assiri</i>, 8/28/2021, retrospective, Saudi Arabia, peer-reviewed, 8 authors, dosage not specified, excluded in exclusion analyses: unadjusted results with no group details.</p>	<p>risk of death, 66.5% higher, RR 1.66, $p = 0.60$, treatment 12 of 90 (13.3%), control 2 of 28 (7.1%), inverted to make $RR < 1$ favor treatment, odds ratio converted to relative risk.</p>
<p><i>Baguma</i>, 12/28/2021, retrospective, Uganda, preprint, 16 authors, study period March 2020 - October 2021, dosage not specified.</p>	<p>risk of death, 96.7% lower, RR 0.03, $p = 0.02$, treatment 23, control 458, adjusted per study, inverted to make $RR < 1$ favor treatment, odds ratio converted to relative risk, multivariable, control prevalence approximated with overall prevalence.</p>
<p><i>Baykal</i>, 5/30/2022, retrospective, Turkey, peer-reviewed, 2 authors, study period 1 April, 2020 - 1 March, 2021, dosage 300,000IU single dose, excluded in exclusion analyses: unadjusted results with no group details; significant confounding by time possible due to separation of groups in different time periods.</p>	<p>risk of death, 22.2% lower, RR 0.78, $p = 0.43$, treatment 7 of 18 (38.9%), control 28 of 56 (50.0%), NNT 9.0.</p>
	<p>risk of ICU admission, 59.4% lower, RR 0.41, $p = 0.005$, treatment 5 of 18 (27.8%), control 39 of 57 (68.4%), NNT 2.5.</p>
<p><i>Beigmohammadi</i>, 11/14/2021, Single Blind Randomized Controlled Trial, Iran, peer-reviewed, 6 authors, study period April 2020 - July 2020, dosage 600,000IU single dose, this trial uses multiple treatments in the treatment arm (combined with vitamins A, B, C, E) - results of individual treatments may vary, trial IRCT20200319046819N1, excluded in exclusion analyses: very late stage study using cholecalciferol instead of calcifediol or calcitriol.</p>	<p>risk of death, 88.9% lower, RR 0.11, $p = 0.11$, treatment 0 of 30 (0.0%), control 4 of 30 (13.3%), NNT 7.5, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm).</p>
	<p>risk of hospitalization >7 days, 41.0% lower, RR 0.59, $p = 0.25$, treatment 4 of 30 (13.3%), control 16 of 30 (53.3%), NNT 2.5, adjusted per study, odds ratio converted to relative risk.</p>
	<p>relative SOFA score @day 7, 45.5% better, RR 0.55, $p < 0.001$, treatment 30, control 30.</p>
<p><i>Bishop</i>, 2/5/2022, Double Blind Randomized Controlled Trial, placebo-controlled, USA, peer-reviewed, survey, 11 authors, study period 2 November, 2020 - 8 October, 2021, dosage calcifediol 300µg days 1-3, 60µg days 4-27, trial NCT04551911 (history) (REsCue).</p>	<p>risk of no recovery, 33.7% lower, RR 0.66, $p = 0.56$, treatment 5 of 65 (7.7%), control 8 of 69 (11.6%), NNT 26, day 21, mid-trial.</p>
	<p>risk of no recovery, 73.5% lower, RR 0.27, $p = 0.37$, treatment 1 of 65 (1.5%), control 4 of 69 (5.8%), NNT 23, day 35.</p>
	<p>risk of no recovery, 57.5% lower, RR 0.42, $p = 0.44$, treatment 2 of 65 (3.1%), control 5 of 69 (7.2%), NNT 24, day 28.</p>

	risk of no recovery, 6.2% higher, RR 1.06, $p = 0.85$, treatment 17 of 65 (26.2%), control 17 of 69 (24.6%), day 14.
	risk of no recovery, 3.0% higher, RR 1.03, $p = 1.00$, treatment 33 of 65 (50.8%), control 34 of 69 (49.3%), day 7.
<p><i>Bychinin</i>, 11/3/2022, Double Blind Randomized Controlled Trial, placebo-controlled, Russia, peer-reviewed, 7 authors, study period 1 May, 2020 - 31 January, 2022, average treatment delay 9.0 days, dosage 60,000IU day 1, 5,000IU days 2-7, 8, 5,000IU days 9-14, 15, 5,000IU days 16-21, 22, 5,000IU days 23-28, trial NCT05092698 (history) (COVID-VIT), excluded in exclusion analyses: very late stage study using cholecalciferol instead of calcifediol or calcitriol.</p>	risk of death, 26.9% lower, RR 0.73, $p = 0.18$, treatment 19 of 52 (36.5%), control 27 of 54 (50.0%), NNT 7.4.
	risk of mechanical ventilation, 7.4% lower, RR 0.93, $p = 0.68$, treatment 33 of 52 (63.5%), control 37 of 54 (68.5%), NNT 20.
<p><i>Cannata-Andía</i>, 2/18/2022, Randomized Controlled Trial, multiple countries, peer-reviewed, median age 59.0, 22 authors, study period 4 April, 2020 - 22 April, 2021, dosage 100,000IU single dose, trial NCT04552951 (history) (COVID-VIT-D), excluded in exclusion analyses: very late stage study using cholecalciferol instead of calcifediol or calcitriol.</p>	risk of death, 44.0% higher, RR 1.44, $p = 0.31$, treatment 22 of 274 (8.0%), control 15 of 269 (5.6%).
	risk of ICU admission, 4.9% higher, RR 1.05, $p = 0.82$, treatment 47 of 274 (17.2%), control 44 of 269 (16.4%).
<p><i>Castillo</i>, 8/29/2020, Randomized Controlled Trial, Spain, peer-reviewed, 7 authors, study period May 2020 - June 2020, dosage calcifediol 0.5mg day 1, 0.27mg day 3, 0.27mg day 7, and then weekly until discharge or ICU admission, trial NCT04366908 (history) (COVIDIOL).</p>	risk of death, 85.4% lower, RR 0.15, $p = 0.11$, treatment 0 of 50 (0.0%), control 2 of 26 (7.7%), NNT 13, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm).
	risk of ICU admission, 94.2% lower, RR 0.06, $p = 0.008$, treatment 1 of 50 (2.0%), control 13 of 26 (50.0%), NNT 2.1, odds ratio converted to relative risk.
<p><i>De Niet</i>, 7/26/2022, Double Blind Randomized Controlled Trial, placebo-controlled, Belgium, peer-reviewed, 16 authors, study period August 2020 - August 2021, dosage 25,000IU days 1-4, 11, 18, 25, trial NCT04636086 (history).</p>	risk of death, 65.1% lower, RR 0.35, $p = 0.61$, treatment 1 of 21 (4.8%), control 3 of 22 (13.6%), NNT 11, COVID-19 mortality.
	risk of death, 39.7% higher, RR 1.40, $p = 0.70$, treatment 4 of 21 (19.0%), control 3 of 22 (13.6%), all cause including after discharge and non-COVID-19.
	risk of ICU admission, 58.1% lower, RR 0.42, $p = 0.41$, treatment 2 of 21 (9.5%), control 5 of 22 (22.7%), NNT 7.6.
	ICU time, 67.7% lower, relative time 0.32, $p = 0.47$, treatment 21, control 22.
	risk of no hospital discharge, 79.6% lower, RR 0.20, $p = 0.49$, treatment 0 of 21 (0.0%), control 2 of 22 (9.1%), NNT 11, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm), day 36.
	risk of no hospital discharge, 85.4% lower, RR 0.15, $p = 0.23$, treatment 0 of 21 (0.0%), control 3 of 22 (13.6%), NNT 7.3, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm), day 28.

	risk of no hospital discharge, 85.4% lower, RR 0.15, $p = 0.23$, treatment 0 of 21 (0.0%), control 3 of 22 (13.6%), NNT 7.3, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm), day 21.
	risk of no hospital discharge, 65.1% lower, RR 0.35, $p = 0.61$, treatment 1 of 21 (4.8%), control 3 of 22 (13.6%), NNT 11, day 14.
	risk of no hospital discharge, 65.1% lower, RR 0.35, $p = 0.03$, treatment 4 of 21 (19.0%), control 12 of 22 (54.5%), NNT 2.8, day 7.
	recovery time, 45.4% lower, relative time 0.55, $p = 0.06$, treatment 21, control 22, fever.
	hospitalization time, 50.0% lower, relative time 0.50, $p = 0.003$, treatment 21, control 22.
<i>Domazet Bugarin</i> , 2/28/2023, Randomized Controlled Trial, Croatia, peer-reviewed, 9 authors, study period November 2021 - May 2022, dosage 10,000IU days 1-14, trial NCT05384574 (history), excluded in exclusion analyses: very late stage study using cholecalciferol instead of calcifediol or calcitriol.	risk of death, 21.0% lower, RR 0.79, $p = 0.20$, treatment 30 of 75 (40.0%), control 39 of 77 (50.6%), NNT 9.4, day 60.
	risk of death, 12.5% lower, RR 0.87, $p = 0.61$, treatment 23 of 75 (30.7%), control 27 of 77 (35.1%), NNT 23, day 28.
	risk of death, 28.9% lower, RR 0.71, $p = 0.49$, treatment 9 of 75 (12.0%), control 13 of 77 (16.9%), NNT 20, day 14.
<i>Elamir</i> , 9/8/2021, Randomized Controlled Trial, USA, peer-reviewed, 9 authors, study period September 2020 - December 2020, dosage calcitriol 0.5µg days 1-14.	risk of death, 85.7% lower, RR 0.14, $p = 0.23$, treatment 0 of 25 (0.0%), control 3 of 25 (12.0%), NNT 8.3, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm).
	risk of mechanical ventilation, 80.0% lower, RR 0.20, $p = 0.48$, treatment 0 of 25 (0.0%), control 2 of 25 (8.0%), NNT 12, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm).
	risk of ICU admission, 37.5% lower, RR 0.62, $p = 0.33$, treatment 5 of 25 (20.0%), control 8 of 25 (32.0%), NNT 8.3.
	hospitalization time, 40.5% lower, relative time 0.60, $p = 0.14$, treatment 25, control 25.
	relative $\Delta \text{SaO}_2/\text{FiO}_2$, RR 0.14, $p = 0.03$, treatment 25, control 25, primary outcome.
<i>Elhadi</i> , 4/30/2021, prospective, Libya, peer-reviewed, 21 authors, study period 29 May, 2020 - 30 December, 2020, dosage not specified, excluded in exclusion analyses: unadjusted results with no group details.	risk of death, 23.4% lower, RR 0.77, $p = 0.29$, treatment 7 of 15 (46.7%), control 274 of 450 (60.9%), NNT 7.0.
<i>Fairfield</i> , 7/26/2022, retrospective, USA, peer-reviewed, 10 authors, study period 1 January, 2020 - 31 July, 2021, dosage not specified, excluded in	risk of death, 8.9% higher, RR 1.09, $p < 0.001$, treatment 3,653 of 28,993 (12.6%), control 13,185 of 129,842 (10.2%), odds ratio converted to relative risk.

exclusion analyses: substantial unadjusted confounding by indication likely.	risk of mechanical ventilation, 40.8% higher, RR 1.41, $p < 0.001$, treatment 4,897 of 28,993 (16.9%), control 15,520 of 129,842 (12.0%), odds ratio converted to relative risk.
<i>Fiore</i> , 5/22/2022, retrospective, matched cohort, Italy, peer-reviewed, mean age 62.5, 10 authors, dosage 100,000IU days 1-2.	risk of death, 92.7% lower, RR 0.07, $p = 0.01$, treatment 3 of 58 (5.2%), control 11 of 58 (19.0%), NNT 7.2, adjusted per study, odds ratio converted to relative risk, multivariable.
	risk of mechanical ventilation, 50.0% lower, RR 0.50, $p = 0.36$, treatment 4 of 58 (6.9%), control 8 of 58 (13.8%), NNT 14.
	risk of ICU admission, 50.0% lower, RR 0.50, $p = 0.36$, treatment 4 of 58 (6.9%), control 8 of 58 (13.8%), NNT 14.
	NIV, 47.8% lower, RR 0.52, $p = 0.04$, treatment 12 of 58 (20.7%), control 23 of 58 (39.7%), NNT 5.3.
<i>Giannini</i> , 1/14/2021, retrospective, Italy, peer-reviewed, 21 authors, dosage 200,000IU days 1-2.	risk of death/ICU, 36.6% lower, RR 0.63, $p = 0.13$, treatment 14 of 36 (38.9%), control 29 of 55 (52.7%), NNT 7.2, odds ratio converted to relative risk.
<i>Güven</i> , 7/23/2021, retrospective, Turkey, peer-reviewed, 2 authors, dosage 300,000IU single dose, excluded in exclusion analyses: very late stage, ICU patients.	risk of death, 24.8% lower, RR 0.75, $p = 0.32$, treatment 43 of 113 (38.1%), control 30 of 62 (48.4%), NNT 9.7, odds ratio converted to relative risk.
<i>Hafez (B)</i> , 8/9/2022, retrospective, Egypt, peer-reviewed, 2 authors, study period April 2020 - June 2020, dosage 50,000IU days 1, 3, 5, 7, 9, 11, 13, 50,000IU every other day for two weeks or one intramuscular shot of 300,000IU.	risk of death, 93.7% lower, RR 0.06, $p = 0.07$, treatment 0 of 7 (0.0%), control 12 of 30 (40.0%), NNT 2.5, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm), high dose, 50,000IU every other day for two weeks or one intramuscular shot of 300,000IU.
	risk of death, 58.3% lower, RR 0.42, $p = 0.28$, treatment 2 of 12 (16.7%), control 12 of 30 (40.0%), NNT 4.3, low dose, $\leq 10,000$ IU/day.
<i>Hafezi</i> , 10/22/2022, retrospective, United Arab Emirates, peer-reviewed, 8 authors, study period September 2020 - January 2021, dosage 50,000IU days 1, 8, 15, excluded in exclusion analyses: very late stage study using cholecalciferol instead of calcifediol or calcitriol.	risk of death, 63.0% lower, HR 0.37, $p = 0.04$, treatment 8 of 43 (18.6%), control 12 of 37 (32.4%), NNT 7.2, Cox proportional hazards, day 29.
<i>Jevalikar</i> , 12/28/2020, prospective, India, peer-reviewed, 8 authors, dosage 60,000IU single dose, median total dose.	risk of death, 82.0% lower, RR 0.18, $p = 0.12$, treatment 1 of 128 (0.8%), control 3 of 69 (4.3%), NNT 28.
	risk of ICU admission, 33.7% lower, RR 0.66, $p = 0.29$, treatment 16 of 128 (12.5%), control 13 of 69 (18.8%), NNT 16.
	risk of oxygen therapy, 31.7% lower, RR 0.68, $p = 0.06$, treatment 38 of 128 (29.7%), control 30 of 69 (43.5%), NNT 7.3.
<i>Karimpour-Razkenari</i> , 10/3/2022, retrospective, Iran, peer-reviewed, median age 58.5, 9 authors, study period 23 February, 2020 - 23 May, 2020, dosage not specified.	risk of death, 79.0% lower, RR 0.21, $p < 0.001$, treatment 10 of 124 (8.1%), control 93 of 329 (28.3%), NNT 4.9, adjusted per study, inverted to make $RR < 1$ favor treatment, odds ratio converted to relative risk, multivariable.

<p><i>Karonova</i>, 6/23/2022, Randomized Controlled Trial, Russia, peer-reviewed, 12 authors, study period 30 November, 2020 - 20 March, 2021, dosage 50,000IU days 1, 8, trial NCT05166005 (history).</p>	<p>risk of ICU admission, 85.9% lower, RR 0.14, $p = 0.11$, treatment 0 of 56 (0.0%), control 3 of 54 (5.6%), NNT 18, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm), day 9.</p>
	<p>risk of oxygen therapy, 7.0% lower, RR 0.93, $p = 0.85$, treatment 27 of 56 (48.2%), control 28 of 54 (51.9%), NNT 27, baseline oxygen supplementation was higher in the treatment group, 38 vs. 32, day 9.</p>
<p><i>Krishnan</i>, 7/20/2020, retrospective, USA, peer-reviewed, 13 authors, dosage not specified, excluded in exclusion analyses: unadjusted results with no group details.</p>	<p>risk of death, 19.0% lower, RR 0.81, $p = 0.42$, treatment 8 of 16 (50.0%), control 84 of 136 (61.8%), NNT 8.5.</p>
<p><i>Lakkireddy</i>, 7/27/2022, Randomized Controlled Trial, India, peer-reviewed, mean age 45.5, 9 authors, dosage 60,000IU days 1-8, 8 or 10 days depending on BMI.</p>	<p>risk of death, 60.9% lower, RR 0.39, $p = 0.27$, treatment 2 of 44 (4.5%), control 5 of 43 (11.6%), NNT 14.</p>
	<p>risk of ICU admission, 21.8% lower, RR 0.78, $p = 0.74$, treatment 4 of 44 (9.1%), control 5 of 43 (11.6%), NNT 39.</p>
	<p>hospitalization time, 7.1% lower, relative time 0.93, $p = 0.90$, treatment 44, control 43.</p>
<p><i>Leal-Martínez</i>, 10/25/2021, Randomized Controlled Trial, Mexico, peer-reviewed, 7 authors, study period 1 September, 2020 - 28 February, 2021, dosage 4,000IU days 1-21, this trial uses multiple treatments in the treatment arm (combined with comprehensive nutritional support) - results of individual treatments may vary, trial NCT04507867 (history), excluded in exclusion analyses: combined treatments may contribute more to the effect seen.</p>	<p>risk of death, 85.7% lower, RR 0.14, $p = 0.03$, treatment 1 of 40 (2.5%), control 7 of 40 (17.5%), NNT 6.7.</p>
	<p>risk of mechanical ventilation, 57.1% lower, RR 0.43, $p = 0.31$, treatment 3 of 40 (7.5%), control 7 of 40 (17.5%), NNT 10.0.</p>
<p><i>Ling</i>, 12/11/2020, retrospective, United Kingdom, peer-reviewed, 7 authors, dosage 40,000IU weekly, regimen varied with 77% receiving a total of 40,000IU/week.</p>	<p>risk of death, 79.8% lower, RR 0.20, $p < 0.001$, treatment 73, control 253, odds ratio converted to relative risk, primary cohort.</p>
	<p>risk of death, 55.5% lower, RR 0.44, $p = 0.02$, treatment 80, control 443, odds ratio converted to relative risk, validation cohort.</p>
<p><i>Lohia (B)</i>, 3/4/2021, retrospective, USA, peer-reviewed, 4 authors, dosage not specified.</p>	<p>risk of death, 10.7% lower, RR 0.89, $p = 0.80$, treatment 26, control 69, odds ratio converted to relative risk, <20 ng/mL, control prevalence approximated with overall prevalence.</p>
	<p>risk of mechanical ventilation, 26.9% lower, RR 0.73, $p = 0.51$, treatment 26, control 69, odds ratio converted to relative risk, <20 ng/mL, control prevalence approximated with overall prevalence.</p>
	<p>risk of ICU admission, 2.7% lower, RR 0.97, $p = 0.93$, treatment 26, control 69, odds ratio converted to relative risk, <20 ng/mL, control prevalence approximated with overall prevalence.</p>
<p><i>Maghbooli</i>, 10/13/2021, Double Blind Randomized Controlled Trial, Iran, peer-reviewed, 12 authors, dosage calcifediol 25µg daily, mean daily dose.</p>	<p>risk of death, 40.0% lower, RR 0.60, $p = 0.72$, treatment 3 of 53 (5.7%), control 5 of 53 (9.4%), NNT 26.</p>

	<p>risk of mechanical ventilation, 60.0% lower, RR 0.40, $p = 0.44$, treatment 2 of 53 (3.8%), control 5 of 53 (9.4%), NNT 18.</p> <p>risk of ICU admission, 40.0% lower, RR 0.60, $p = 0.42$, treatment 6 of 53 (11.3%), control 10 of 53 (18.9%), NNT 13.</p> <p>ICU time, 36.4% lower, relative time 0.64, $p = 0.20$, treatment 53, control 53.</p> <p>hospitalization time, 16.7% lower, relative time 0.83, $p = 0.10$, treatment 53, control 53.</p>
<p><i>Mahmood</i>, 12/29/2021, retrospective, United Kingdom, peer-reviewed, 4 authors, study period 23 March, 2020 - 31 December, 2020, dosage varies, excluded in exclusion analyses: unadjusted results with no group details; substantial unadjusted confounding by indication likely.</p>	<p>risk of death, 30.5% lower, RR 0.70, $p = 0.10$, treatment 45 of 238 (18.9%), control 31 of 114 (27.2%), NNT 12, started after admission, late treatment result.</p>
<p><i>Mariani</i>, 5/27/2022, Double Blind Randomized Controlled Trial, placebo-controlled, Argentina, peer-reviewed, mean age 59.1, 33 authors, study period 14 August, 2020 - 22 June, 2021, average treatment delay 7.0 days, dosage 500,000IU single dose, trial NCT04411446 (history) (CARED).</p>	<p>risk of death, 124.0% higher, RR 2.24, $p = 0.45$, treatment 5 of 115 (4.3%), control 2 of 103 (1.9%).</p> <p>risk of mechanical ventilation, 25.0% lower, RR 0.75, $p = 0.85$, treatment 5 of 115 (4.3%), control 6 of 103 (5.8%), NNT 68.</p> <p>risk of ICU admission, 27.0% lower, RR 0.73, $p = 0.62$, treatment 9 of 115 (7.8%), control 11 of 103 (10.7%), NNT 35.</p> <p>risk of progression, 3.0% lower, OR 0.97, $p = 0.82$, treatment 115, control 103, Wilcoxon-Mann-Whitney, primary outcome, RR approximated with OR.</p> <p>risk of progression, 32.8% lower, RR 0.67, $p = 0.71$, treatment 3 of 115 (2.6%), control 4 of 103 (3.9%), NNT 78, Δ rSOFA 4.</p> <p>risk of progression, 79.1% higher, RR 1.79, $p = 0.30$, treatment 10 of 115 (8.7%), control 5 of 103 (4.9%), Δ rSOFA 3.</p> <p>risk of progression, 25.4% lower, RR 0.75, $p = 0.76$, treatment 5 of 115 (4.3%), control 6 of 103 (5.8%), NNT 68, Δ rSOFA 2.</p> <p>risk of progression, 16.0% lower, RR 0.84, $p = 0.70$, treatment 15 of 115 (13.0%), control 16 of 103 (15.5%), NNT 40, Δ rSOFA 1.</p>
<p><i>Mazziotti</i>, 3/5/2021, retrospective, Italy, peer-reviewed, 11 authors, dosage varies.</p>	<p>risk of death, 19.0% lower, OR 0.81, $p = 0.49$, treatment 116, control 232, supplementation, RR approximated with OR.</p> <p>risk of mechanical ventilation, 67.0% higher, OR 1.67, $p = 0.08$, treatment 116, control 232, supplementation, RR approximated with OR.</p>
<p><i>Mingiano</i>, 7/30/2023, retrospective, Italy, peer-reviewed, 11 authors, study period November 2020 - February 2021, dosage calcifediol 450μg days 1-2, patients with deficiency only.</p>	<p>risk of death, 38.8% lower, RR 0.61, $p = 0.04$, treatment 13 of 56 (23.2%), control 88 of 232 (37.9%), NNT 6.8.</p>

	risk of oxygen therapy, 23.1% lower, RR 0.77, $p = 0.22$, treatment 18 of 56 (32.1%), control 97 of 232 (41.8%), NNT 10.
	hospitalization time, 34.6% lower, relative time 0.65, $p = 0.01$, treatment 56, control 232.
<p><i>Murai</i>, 11/17/2020, Double Blind Randomized Controlled Trial, Brazil, peer-reviewed, 17 authors, study period 2 June, 2020 - 27 August, 2020, average treatment delay 10.2 days, dosage 200,000IU single dose, trial NCT04449718 (history), excluded in exclusion analyses: very late stage, >50% on oxygen/ventilation at baseline; very late stage study using cholecalciferol instead of calcifediol or calcitriol.</p>	risk of death, 48.7% higher, RR 1.49, $p = 0.43$, treatment 9 of 119 (7.6%), control 6 of 118 (5.1%).
	risk of mechanical ventilation, 47.5% lower, RR 0.52, $p = 0.09$, treatment 9 of 119 (7.6%), control 17 of 118 (14.4%), NNT 15.
	risk of ICU admission, 24.6% lower, RR 0.75, $p = 0.30$, treatment 19 of 119 (16.0%), control 25 of 118 (21.2%), NNT 19.
	risk of no hospital discharge, 6.5% lower, HR 0.93, $p = 0.63$, treatment 119, control 118, inverted to make HR<1 favor treatment.
<p><i>Nogués</i>, 1/22/2021, prospective quasi-randomized (ward), Spain, peer-reviewed, 16 authors, dosage calcifediol 0.5mg day 1, 0.27mg day 3, 0.27mg day 7, 0.27mg day 15, 0.27mg day 30.</p>	risk of death, 79.0% lower, RR 0.21, $p = 0.001$, treatment 21 of 447 (4.7%), control 62 of 391 (15.9%), NNT 9.0, adjusted per study, ITT.
	risk of death, 48.0% lower, RR 0.52, $p = 0.001$, treatment 500, control 338, adjusted per study, including patients treated later.
	risk of ICU admission, 87.0% lower, RR 0.13, $p < 0.001$, treatment 20 of 447 (4.5%), control 82 of 391 (21.0%), NNT 6.1, adjusted per study, ITT.
<p><i>Ogasawara</i>, 9/1/2023, retrospective, Japan, peer-reviewed, 10 authors, study period April 2021 - September 2022, dosage alfacalcidol 1µg days 1-8, median duration, alfacalcidol and eldecacitol used.</p>	risk of death, 66.7% lower, RR 0.33, $p = 1.00$, treatment 0 of 54 (0.0%), control 1 of 54 (1.9%), NNT 54, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm).
	risk of progression, 77.8% lower, RR 0.22, $p = 0.05$, treatment 2 of 54 (3.7%), control 9 of 54 (16.7%), NNT 7.7, high-flow oxygen, mechanical ventilation, or mortality, primary outcome.
	risk of oxygen therapy, 75.0% lower, RR 0.25, $p = 0.09$, treatment 2 of 54 (3.7%), control 8 of 54 (14.8%), NNT 9.0.
<p><i>Rastogi</i>, 11/12/2020, Randomized Controlled Trial, India, peer-reviewed, 8 authors, dosage 60,000IU days 1-7, trial NCT04459247 (history) (SHADE).</p>	risk of no viral clearance, 52.6% lower, RR 0.47, $p = 0.02$, treatment 6 of 16 (37.5%), control 19 of 24 (79.2%), NNT 2.4.
<p><i>Saheb Sharif-Askari</i>, 8/24/2022, retrospective, USA, peer-reviewed, 10 authors, dosage 50,000IU days 1, 8, 15, excluded in exclusion analyses: very late stage study using cholecalciferol instead of calcifediol or calcitriol.</p>	ICU time, 35.7% lower, relative time 0.64, $p = 0.01$, treatment 20, control 25.
<p><i>Seely</i>, 9/22/2023, Double Blind Randomized Controlled Trial, placebo-controlled, Canada, peer-reviewed, mean age 39.9, 10 authors, study period September 2021 - April 2022, dosage 51,000IU day 1, 1,000IU days 2-21, this trial uses multiple</p>	ER visit, 47.6% lower, RR 0.52, $p = 0.68$, treatment 2 of 42 (4.8%), control 4 of 44 (9.1%), NNT 23.

treatments in the treatment arm (combined with vitamin C, D, K2, and zinc) - results of individual treatments may vary, trial NCT04780061 (history).	relative mean cumulative symptom score, 13.8% better, RR 0.86, $p = 0.41$, treatment mean 166.3 (± 92.3) $n=34$, control mean 192.9 (± 153.6) $n=24$.
	EQ-VAS average score <80, 29.4% lower, RR 0.71, $p = 0.54$, treatment 7 of 34 (20.6%), control 7 of 24 (29.2%), NNT 12, average daily EQ-VAS score <80.
	relative EQ5D improvement, 28.6% better, RR 0.71, $p = 0.44$, treatment 32, control 31, relative improvement in EQ5D, week 1.
	relative EQ5D improvement, 14.3% better, RR 0.86, $p = 0.73$, treatment 33, control 30, relative improvement in EQ5D, week 2.
	relative EQ5D improvement, 50.0% better, RR 0.50, $p = 0.17$, treatment 32, control 33, relative improvement in EQ5D, week 3.
	relative EQ5D improvement, 12.5% worse, RR 1.12, $p = 0.47$, treatment 30, control 25, relative improvement in EQ5D, week 4.
	recovery time, 4.0% higher, relative time 1.04, $p = 0.81$, treatment 34, control 24.
	risk of PASC, 12.1% lower, RR 0.88, $p = 1.00$, treatment 3 of 33 (9.1%), control 3 of 29 (10.3%), NNT 80, 12 weeks.
	risk of PASC, 35.7% lower, RR 0.64, $p = 0.69$, treatment 3 of 35 (8.6%), control 4 of 30 (13.3%), NNT 21, 8 weeks.
	risk of PASC, 0.6% lower, RR 0.99, $p = 1.00$, treatment 6 of 35 (17.1%), control 5 of 29 (17.2%), NNT 1015, 4 weeks.
<i>Shahid</i> , 6/17/2022, retrospective, USA, peer-reviewed, 2 authors, dosage not specified, excluded in exclusion analyses: minimal details provided.	risk of death, 38.0% lower, RR 0.62, $p < 0.001$, treatment 705, control 773.
<i>Shamsi</i> , 7/17/2023, retrospective, Iran, peer-reviewed, 4 authors, study period 1 March, 2020 - 1 August, 2021, dosage not specified, excluded in exclusion analyses: unadjusted results with no group details.	risk of death, 57.5% lower, RR 0.42, $p = 0.70$, treatment 1 of 17 (5.9%), control 23 of 166 (13.9%), NNT 13.
<i>Singh</i> , 6/1/2022, Double Blind Randomized Controlled Trial, placebo-controlled, India, peer-reviewed, 10 authors, study period 1 August, 2021 - 10 December, 2021, dosage 600,000IU single dose, trial NCT04952857 (history) (Shade-S), excluded in exclusion analyses: minimal details provided.	risk of death, 45.0% lower, RR 0.55, $p = 0.046$, treatment 11 of 45 (24.4%), control 20 of 45 (44.4%), NNT 5.0.
<i>Soliman</i> , 9/1/2021, Randomized Controlled Trial, placebo-controlled, Egypt, peer-reviewed, 3 authors, dosage 200,000IU single dose.	risk of death, 63.4% lower, RR 0.37, $p = 0.21$, treatment 7 of 40 (17.5%), control 3 of 16 (18.8%), adjusted per study, odds ratio converted to relative risk, logistic regression.
	risk of mechanical ventilation, 20.0% lower, RR 0.80, $p = 0.56$, treatment 14 of 40 (35.0%), control 7 of 16 (43.8%), NNT 11, unadjusted.

	risk of no recovery, 20.0% lower, RR 0.80, $p = 0.56$, treatment 14 of 40 (35.0%), control 7 of 16 (43.8%), NNT 11, unadjusted.
<i>Tan (B)</i> , 6/10/2020, retrospective, Singapore, peer-reviewed, 14 authors, dosage 1,000IU daily, this trial uses multiple treatments in the treatment arm (combined with magnesium and vitamin B12) - results of individual treatments may vary.	risk of oxygen therapy, 80.5% lower, RR 0.20, $p = 0.04$, treatment 3 of 17 (17.6%), control 16 of 26 (61.5%), NNT 2.3, adjusted per study, multivariate.
	risk of ICU admission, 80.9% lower, RR 0.19, $p = 0.07$, treatment 1 of 17 (5.9%), control 8 of 26 (30.8%), NNT 4.0, no adjusted result available.
<i>Yildiz</i> , 9/27/2021, retrospective, Turkey, peer-reviewed, 5 authors, dosage 300,000IU single dose.	risk of death, 80.9% lower, RR 0.19, $p = 0.04$, treatment 1 of 37 (2.7%), control 24 of 170 (14.1%), NNT 8.8.
	risk of ICU admission, 94.5% lower, RR 0.06, $p = 0.13$, treatment 0 of 37 (0.0%), control 14 of 170 (8.2%), NNT 12, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm).
	hospitalization time, 9.6% lower, relative time 0.90, $p = 0.32$, treatment 37, control 170.
<i>Zangeneh</i> , 5/13/2022, retrospective, Iran, peer-reviewed, 3 authors, dosage not specified, excluded in exclusion analyses: very late stage study using cholecalciferol instead of calcifediol or calcitriol.	risk of death, 26.0% higher, HR 1.26, $p = 0.40$, Cox proportional hazards.
<i>Zurita-Cruz</i> , 7/25/2022, Single Blind Randomized Controlled Trial, Mexico, peer-reviewed, median age 12.0, 7 authors, study period 24 March, 2020 - 31 March, 2021, dosage 2,000IU daily, daily, 1,000IU for children <1 year, trial NCT04502667 (history), excluded in exclusion analyses: randomization resulted in significant baseline differences that were not adjusted for.	risk of death, 79.2% lower, RR 0.21, $p = 0.11$, treatment 1 of 20 (5.0%), control 6 of 25 (24.0%), NNT 5.3.
	risk of mechanical ventilation, 72.2% lower, RR 0.28, $p = 0.08$, treatment 2 of 20 (10.0%), control 9 of 25 (36.0%), NNT 3.8.
	risk of ICU admission, 73.2% lower, RR 0.27, $p = 0.006$, treatment 3 of 20 (15.0%), control 14 of 25 (56.0%), NNT 2.4.

Prophylaxis

Effect extraction follows pre-specified rules as detailed above and gives priority to more serious outcomes. Only the first (most serious) outcome is used in pooled analysis, which may differ from the effect a paper focuses on. Other outcomes are used in outcome specific analyses.

<i>Abdulateef</i> , 4/8/2021, retrospective, Iraq, peer-reviewed, 7 authors, study period July 2020 - August 2020, dosage varies, excluded in exclusion analyses: unadjusted results with no group details.	risk of hospitalization, 40.9% lower, RR 0.59, $p = 0.30$, treatment 6 of 127 (4.7%), control 24 of 300 (8.0%), NNT 31, unadjusted.
<i>Ahmed</i> , 11/21/2021, retrospective, USA, preprint, 5 authors, dosage not specified.	risk of death, 10.5% lower, RR 0.90, $p = 0.28$.
<i>Aldwihi</i> , 5/11/2021, retrospective, Saudi Arabia, peer-reviewed, survey, mean age 36.5, 8 authors, study period August 2020 - October 2020, dosage not specified.	risk of hospitalization, 49.3% higher, RR 1.49, $p = 0.002$, treatment 94 of 259 (36.3%), control 143 of 479 (29.9%), adjusted per study, odds ratio converted to relative risk, multivariable.

<i>Annweiler (C)</i> , 11/2/2020, retrospective, France, peer-reviewed, mean age 88.0, 7 authors, dosage 50,000IU monthly, dose varies - 50,000 IU/month, or 80,000IU/100,000IU every 2–3 months.	risk of death, 93.0% lower, RR 0.07, $p = 0.02$, treatment 2 of 29 (6.9%), control 10 of 32 (31.2%), NNT 4.1, adjusted per study, regular bolus supplementation.
<i>Arroyo-Díaz</i> , 9/24/2021, retrospective, Spain, peer-reviewed, 11 authors, dosage not specified.	risk of death, 12.4% higher, RR 1.12, $p = 0.59$, treatment 50 of 189 (26.5%), control 167 of 1,078 (15.5%), adjusted per study, odds ratio converted to relative risk.
	risk of mechanical ventilation, 43.3% lower, RR 0.57, $p = 0.22$, treatment 11 of 189 (5.8%), control 113 of 1,078 (10.5%), NNT 21, adjusted per study, odds ratio converted to relative risk.
	risk of ICU admission, 44.2% lower, RR 0.56, $p = 0.03$, treatment 13 of 189 (6.9%), control 133 of 1,078 (12.3%), NNT 18, unadjusted.
	hospitalization time, 11.8% lower, relative time 0.88, $p = 0.20$, treatment 189, control 1,078, unadjusted.
<i>Aweimer</i> , 3/29/2023, retrospective, Germany, peer-reviewed, median age 67.0, 19 authors, study period 1 March, 2020 - 31 August, 2021, excluded in exclusion analyses: unadjusted results with no group details.	risk of death, 20.9% lower, RR 0.79, $p = 0.31$, treatment 7 of 12 (58.3%), control 101 of 137 (73.7%), NNT 6.5.
<i>Bagheri</i> , 9/1/2021, retrospective, Iran, peer-reviewed, 6 authors, dosage not specified.	risk of severe case, 70.9% lower, OR 0.29, $p = 0.02$, treatment 131, control 379, adjusted per study, multinomial logistic regression, RR approximated with OR.
	risk of hospitalization, 37.9% lower, RR 0.62, $p = 0.11$, treatment 28 of 131 (21.4%), control 143 of 379 (37.7%), NNT 6.1, adjusted per study, inverted to make $RR < 1$ favor treatment, odds ratio converted to relative risk, binary logistic regression.
<i>Baralić</i> , 4/24/2023, prospective, France, peer-reviewed, 15 authors, study period March 2020 - September 2022, dosage not specified.	risk of death, 66.8% lower, HR 0.33, $p = 0.02$, treatment 7 of 31 (22.6%), control 11 of 21 (52.4%), NNT 3.4, Cox proportional hazards.
<i>Bhat</i> , 3/6/2023, prospective, placebo-controlled, India, peer-reviewed, 13 authors, dosage calcifediol 50µg days 1-180, trial CTRI/2021/08/035709.	risk of symptomatic case, 34.2% lower, RR 0.66, $p = 0.01$, treatment 59 of 262 (22.5%), control 52 of 152 (34.2%), NNT 8.6.
<i>Blanch-Rubió</i> , 10/20/2020, retrospective, Spain, peer-reviewed, mean age 66.4, 10 authors, dosage not specified.	risk of case, 8.0% lower, RR 0.92, $p = 0.68$, treatment 62 of 1,303 (4.8%), control 47 of 799 (5.9%), adjusted per study.
<i>Brunvoll</i> , 9/7/2022, Double Blind Randomized Controlled Trial, placebo-controlled, Norway, peer-reviewed, mean age 44.9, 15 authors, study period 10 November, 2020 - 2 June, 2021, dosage 400IU daily, this trial uses multiple treatments in the treatment arm (combined with cod liver oil) - results of individual treatments may vary, trial NCT04609423 (history).	risk of ICU admission, 0.3% higher, RR 1.00, $p = 1.00$, treatment 4 of 17,278 (0.0%), control 4 of 17,323 (0.0%).
	risk of hospitalization, 10.9% lower, RR 0.89, $p = 1.00$, treatment 8 of 17,278 (0.0%), control 9 of 17,323 (0.1%), NNT 17692.
	risk of severe case, 20.0% higher, RR 1.20, $p = 0.17$, treatment 121 of 17,278 (0.7%), control 101 of 17,323 (0.6%).

	risk of case, no change, RR 1.00, $p = 0.98$, treatment 227 of 17,278 (1.3%), control 228 of 17,323 (1.3%), NNT 42377.
<i>Campi</i> , 6/14/2021, prospective, Italy, peer-reviewed, 21 authors, dosage not specified, excluded in exclusion analyses: significant unadjusted differences between groups.	risk of severe case, 88.4% lower, OR 0.12, $p < 0.001$, treatment 31 of 103 (30.1%) cases, 41 of 52 (78.8%) controls, NNT 2.3, case control OR, vitamin D supplementation, hospitalized patients vs. controls.
<i>Cangiano</i> , 12/22/2020, retrospective, Italy, peer-reviewed, 14 authors, dosage 25,000IU 2x per month.	risk of death, 70.0% lower, RR 0.30, $p = 0.04$, treatment 3 of 20 (15.0%), control 39 of 78 (50.0%), NNT 2.9.
<i>Cereda (B)</i> , 11/11/2020, retrospective, Italy, peer-reviewed, mean age 68.8, 7 authors, dosage varies.	risk of death, 73.0% higher, RR 1.73, $p = 0.14$, treatment 7 of 18 (38.9%), control 40 of 152 (26.3%), odds ratio converted to relative risk, $\geq 25,000\text{IU/month}$ for at least 3 months.
	risk of hospitalization, 17.3% higher, RR 1.17, $p = 0.68$, treatment 7 of 27 (25.9%), control 36 of 170 (21.2%), odds ratio converted to relative risk.
<i>De Nicolò</i> , 12/29/2022, prospective, Italy, peer-reviewed, 11 authors, study period January 2021 - April 2021, dosage not specified.	risk of IgG positive, 88.4% lower, OR 0.12, $p = 0.002$, treatment 43, control 63, adjusted per study, multivariable, RR approximated with OR.
<i>Dudley</i> , 5/18/2021, retrospective, United Kingdom, peer-reviewed, 5 authors, dosage 800IU daily.	risk of symptomatic case, 22.4% lower, RR 0.78, $p = 0.65$, treatment 15 of 58 (25.9%), control 2 of 6 (33.3%), NNT 13, positive test.
<i>Fasano</i> , 6/2/2021, retrospective, Italy, peer-reviewed, 7 authors, dosage not specified.	risk of case, 42.0% lower, RR 0.58, $p = 0.048$, treatment 13 of 329 (4.0%), control 92 of 1,157 (8.0%), NNT 25, odds ratio converted to relative risk.
<i>Gibbons</i> , 11/12/2022, retrospective, USA, peer-reviewed, 7 authors, dosage varies.	risk of death, 33.3% lower, HR 0.67, $p < 0.001$, treatment 5,315 of 199,498 (2.7%), control 6,591 of 199,498 (3.3%), D3, propensity score matching, Cox proportional hazards.
	risk of death, 23.5% lower, HR 0.77, $p = 0.10$, treatment 716 of 33,216 (2.2%), control 987 of 33,216 (3.0%), NNT 123, D2, propensity score matching, Cox proportional hazards.
	risk of case, 20.3% lower, HR 0.80, $p < 0.001$, treatment 462 of 199,498 (0.2%), control 689 of 199,498 (0.3%), D3, propensity score matching, Cox proportional hazards.
	risk of case, 28.0% lower, HR 0.72, $p < 0.001$, treatment 65 of 33,216 (0.2%), control 86 of 33,216 (0.3%), NNT 1582, D2, propensity score matching, Cox proportional hazards.
<i>Golabi (B)</i> , 8/26/2021, retrospective, Iran, peer-reviewed, 10 authors, dosage not specified.	risk of case, 25.4% higher, OR 1.25, $p = 0.56$, treatment 28 of 53 (52.8%) cases, 25 of 53 (47.2%) controls, case control OR.
<i>Guldemir</i> , 11/16/2022, retrospective, Turkey, peer-reviewed, 3 authors, study period 30 March, 2020 - 23 September, 2020, dosage not specified, excluded in exclusion analyses: unadjusted results with no group details.	risk of hospitalization, 5.2% lower, RR 0.95, $p = 0.89$ (Fisher's exact test), treatment 19 of 81 (23.5%), control 98 of 396 (24.7%), NNT 77.

<p><i>Hernández (B)</i>, 10/27/2020, retrospective, Spain, peer-reviewed, mean age 60.9, 12 authors, dosage varies.</p>	<p>risk of death, 3.7% higher, RR 1.04, $p = 1.00$, treatment 2 of 19 (10.5%), control 20 of 197 (10.2%).</p>
	<p>risk of mechanical ventilation, 75.9% lower, RR 0.24, $p = 0.13$, treatment 1 of 19 (5.3%), control 43 of 197 (21.8%), NNT 6.0.</p>
	<p>risk of ICU admission, 79.3% lower, RR 0.21, $p = 0.05$, treatment 1 of 19 (5.3%), control 50 of 197 (25.4%), NNT 5.0.</p>
	<p>hospitalization time, 33.3% lower, relative time 0.67, $p = 0.11$, treatment 19, control 197.</p>
<p><i>Holt</i>, 3/30/2021, prospective, United Kingdom, peer-reviewed, 34 authors, study period 1 May, 2020 - 5 February, 2021, dosage not specified, trial NCT04330599 (history) (COVIDENCE UK), excluded in exclusion analyses: significant unadjusted confounding possible.</p>	<p>risk of case, 6.8% lower, RR 0.93, $p = 0.53$, treatment 141 of 5,640 (2.5%), control 305 of 9,587 (3.2%), adjusted per study, odds ratio converted to relative risk, fully adjusted, group sizes approximated.</p>
<p><i>Hosseini (C)</i>, 7/19/2022, Double Blind Randomized Controlled Trial, placebo-controlled, Canada, preprint, mean age 39.5, 9 authors, study period 8 February, 2021 - 4 May, 2021, dosage 100,000IU day 1, 10,000IU day 7, 10,000IU day 14, 10,000IU day 21, 10,000IU day 28, 100,000IU cholecalciferol at baseline, 10,000IU weekly for 16 weeks, trial NCT04483635 (history) (PROTECT).</p>	<p>risk of case, 81.9% lower, RR 0.18, $p = 0.19$, treatment 0 of 19 (0.0%), control 2 of 15 (13.3%), NNT 7.5, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm).</p>
<p><i>Israel (B)</i>, 7/27/2021, retrospective, Israel, peer-reviewed, 10 authors, dosage not specified.</p>	<p>risk of hospitalization, 13.1% lower, OR 0.87, $p = 0.003$, treatment 737 of 6,953 (10.6%) cases, 1,669 of 13,906 (12.0%) controls, NNT 33, case control OR, PCR+, cohort 2.</p>
<p><i>Jabeen</i>, 5/11/2022, prospective, Pakistan, peer-reviewed, 7 authors, dosage 200,000IU single dose.</p>	<p>risk of symptomatic case, 88.9% lower, RR 0.11, $p = 0.11$, treatment 0 of 20 (0.0%), control 4 of 20 (20.0%), NNT 5.0, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm).</p>
<p><i>Jimenez</i>, 7/26/2021, retrospective, Spain, peer-reviewed, 21 authors, study period 12 March, 2020 - 21 May, 2020, dosage paricalcitol 0.9µg weekly.</p>	<p>risk of death, 50.1% lower, HR 0.50, $p = 0.02$, treatment 16 of 94 (17.0%), control 65 of 191 (34.0%), NNT 5.9, adjusted per study, paricalcitol treatment, multivariate Cox regression.</p>
	<p>risk of death, 50.7% lower, HR 0.49, $p = 0.003$, all vitamin D derivatives, univariate.</p>
<p><i>Jolliffe</i>, 3/23/2022, Randomized Controlled Trial, United Kingdom, peer-reviewed, median age 60.2, 24 authors, study period December 2020 - June 2021, dosage 3,200IU daily, daily, trial NCT04579640 (history) (CORONAVIT).</p>	<p>risk of mechanical ventilation, 94.7% higher, RR 1.95, $p = 1.00$, treatment 1 of 1,515 (0.1%), control 1 of 2,949 (0.0%), 3200IU/day.</p>
	<p>risk of mechanical ventilation, 94.7% higher, RR 1.95, $p = 1.00$, treatment 1 of 1,515 (0.1%), control 1 of 2,949 (0.0%), 800IU/day.</p>
	<p>risk of hospitalization, 41.1% higher, RR 1.41, $p = 0.16$, treatment 29 of 1,515 (1.9%), control 40 of 2,949 (1.4%), 3200IU/day.</p>

	risk of hospitalization, 16.8% higher, RR 1.17, $p = 0.60$, treatment 24 of 1,515 (1.6%), control 40 of 2,949 (1.4%), 800IU/day.
	risk of case, 8.8% higher, RR 1.09, $p = 0.55$, treatment 76 of 1,515 (5.0%), control 136 of 2,949 (4.6%), 3200IU/day.
	risk of case, 24.5% higher, RR 1.25, $p = 0.11$, treatment 87 of 1,515 (5.7%), control 136 of 2,949 (4.6%), 800IU/day.
	risk of case, 12.3% higher, RR 1.12, $p = 0.56$, treatment 45 of 1,515 (3.0%), control 78 of 2,949 (2.6%), confirmed, 3200IU/day.
	risk of case, 37.3% higher, RR 1.37, $p = 0.08$, treatment 55 of 1,515 (3.6%), control 78 of 2,949 (2.6%), confirmed, 800IU/day.
<i>Junior</i> , 2/17/2022, prospective, Brazil, peer-reviewed, 6 authors, dosage not specified, excluded in exclusion analyses: unadjusted results with no group details.	risk of death, 22.1% lower, RR 0.78, $p = 0.61$, treatment 8 of 113 (7.1%), control 8 of 88 (9.1%), NNT 50.
	risk of progression, 30.8% lower, RR 0.69, $p = 0.26$, treatment 16 of 113 (14.2%), control 18 of 88 (20.5%), NNT 16, respiratory failure.
<i>Levitus</i> , 5/3/2021, retrospective, USA, peer-reviewed, 9 authors, dosage varies.	risk of severe case, 30.8% lower, RR 0.69, $p = 0.25$, treatment 65, control 64, odds ratio converted to relative risk, $\geq 1,000$ IU, control prevalence approximated with overall prevalence.
	risk of severe case, 40.0% lower, RR 0.60, $p = 0.15$, treatment 65, control 64, odds ratio converted to relative risk, $\geq 5,000$ IU, control prevalence approximated with overall prevalence.
	risk of severe case, no change, RR 1.00, $p = 0.92$, treatment 65, control 64, odds ratio converted to relative risk, $\geq 50,000$ IU, control prevalence approximated with overall prevalence.
<i>Levy</i> , 1/31/2022, retrospective, Israel, peer-reviewed, 10 authors, dosage not specified.	risk of death/hospitalization, 30.0% lower, HR 0.70, $p = 0.05$, treatment 39 of 208 (18.8%), control 168 of 641 (26.2%), NNT 13, adjusted per study, multivariable, Cox proportional hazards, day 40.
<i>Louca</i> , 11/30/2020, retrospective, population-based cohort, United Kingdom, peer-reviewed, mean age 49.6, 26 authors, dosage not specified.	risk of case, 7.5% lower, RR 0.92, $p < 0.001$, odds ratio converted to relative risk, United Kingdom, all adjustment model.
<i>Loucera</i> , 4/29/2021, retrospective, propensity score matching, Spain, peer-reviewed, 11 authors, dosage varies (calcifediol).	risk of death, 33.0% lower, HR 0.67, $p = 0.009$, treatment 374, control 374, calcifediol, <15 days before hospitalization, Cox model with inverse propensity weighting.
	risk of death, 27.0% lower, HR 0.73, $p = 0.02$, treatment 439, control 439, calcifediol, <30 days before hospitalization, Cox model with inverse propensity weighting.
	risk of death, 25.0% lower, HR 0.75, $p = 0.005$, treatment 570, control 570, cholecalciferol, <15 days before hospitalization, Cox model with inverse propensity weighting.

	risk of death, 12.0% lower, HR 0.88, $p = 0.11$, treatment 802, control 802, cholecalciferol, <30 days before hospitalization, Cox model with inverse propensity weighting.
<i>Lázaro</i> , 9/5/2021, retrospective, Spain, preprint, 9 authors, dosage not specified, excluded in exclusion analyses: very few events; unadjusted results with no group details; minimal details provided.	risk of case, 26.8% lower, RR 0.73, $p = 1.00$, treatment 1 of 97 (1.0%), control 2 of 142 (1.4%), NNT 265.
<i>Ma</i> , 12/3/2021, retrospective, USA, peer-reviewed, 16 authors, study period May 2020 - March 2021, dosage varies.	risk of hospitalization, 49.0% lower, OR 0.51, $p = 0.04$, treatment 26,605, control 12,710, adjusted per study, supplementation ≥ 400 IU/day, model 3, supplemental table 3, multivariable, RR approximated with OR.
	risk of symptomatic case, 7.0% higher, OR 1.07, $p = 0.25$, treatment 7,895, control 31,420, adjusted per study, supplementation ≥ 2000 IU/day vs. <400 IU/day, model 3, supplemental table 3, multivariable, RR approximated with OR.
	risk of case, 17.0% lower, OR 0.83, $p = 0.07$, treatment 7,895, control 31,420, adjusted per study, supplementation ≥ 2000 IU/day vs. <400 IU/day, model 3, supplemental table 3, multivariable, RR approximated with OR.
<i>Ma (B)</i> , 1/29/2021, retrospective, United Kingdom, peer-reviewed, 4 authors, dosage not specified.	risk of case, 30.0% lower, RR 0.70, $p = 0.03$, treatment 49 of 363 (13.5%), control 1,329 of 7,934 (16.8%), adjusted per study, odds ratio converted to relative risk.
<i>Mahmood</i> , 12/29/2021, retrospective, United Kingdom, peer-reviewed, 4 authors, study period 23 March, 2020 - 31 December, 2020, dosage varies, excluded in exclusion analyses: unadjusted results with no group details; substantial unadjusted confounding by indication likely.	risk of death, 9.4% lower, RR 0.91, $p = 0.67$, treatment 34 of 138 (24.6%), control 31 of 114 (27.2%), NNT 39, prescribed by GP.
<i>Meltzer (C)</i> , 3/19/2021, retrospective, database analysis, USA, peer-reviewed, 6 authors, dosage not specified.	risk of case, 36.0% lower, RR 0.64, $p = 0.38$, treatment 6 of 131 (4.6%), control 239 of 3,338 (7.2%), NNT 39, $\geq 2,000$ IU/d.
	risk of case, 31.1% lower, RR 0.69, $p = 0.16$, treatment 15 of 304 (4.9%), control 239 of 3,338 (7.2%), NNT 45, $\geq 1,001$ IU/d.
	risk of case, 8.9% lower, RR 0.91, $p = 0.56$, treatment 60 of 920 (6.5%), control 239 of 3,338 (7.2%), NNT 157, ≥ 1 IU/d.
<i>Mohseni</i> , 8/4/2021, retrospective, Iran, peer-reviewed, 4 authors, dosage not specified, excluded in exclusion analyses: unadjusted results with no group details.	risk of case, 12.4% lower, RR 0.88, $p = 0.09$, treatment 99 of 192 (51.6%), control 242 of 411 (58.9%), NNT 14.
<i>Nimer</i> , 2/28/2022, retrospective, Jordan, peer-reviewed, survey, 4 authors, study period March 2021 - July 2021, dosage not specified.	risk of hospitalization, 33.3% lower, RR 0.67, $p = 0.001$, treatment 66 of 796 (8.3%), control 153 of 1,352 (11.3%), NNT 33, adjusted per study, odds ratio converted to relative risk, multivariable.

	<p>risk of severe case, 29.0% lower, RR 0.71, $p = 0.01$, treatment 81 of 796 (10.2%), control 179 of 1,352 (13.2%), NNT 33, adjusted per study, odds ratio converted to relative risk, multivariable.</p>
<p><i>Oristrell</i>, 7/17/2021, retrospective, population-based cohort, Spain, peer-reviewed, 8 authors, dosage varies (calcifediol).</p>	<p>risk of death, 1.0% higher, RR 1.01, $p = 0.91$, calcifediol, univariate.</p>
	<p>risk of death, 4.0% lower, RR 0.96, $p = 0.37$, cholecalciferol, univariate.</p>
	<p>risk of case, 1.0% lower, RR 0.99, $p = 0.65$, NNT 3499, calcifediol, univariate.</p>
	<p>risk of case, 5.0% lower, RR 0.95, $p = 0.004$, cholecalciferol, multivariate.</p>
<p><i>Oristrell (B)</i>, 4/6/2021, retrospective, Spain, peer-reviewed, 10 authors, dosage calcitriol 0.3µg daily, mean daily dose.</p>	<p>risk of death, 43.0% lower, HR 0.57, $p = 0.001$, treatment 2,296, control 3,407, multivariate, patients with CKD stages 4-5.</p>
	<p>risk of severe case, 43.0% lower, HR 0.57, $p < 0.001$, treatment 2,296, control 3,407, multivariate, patients with CKD stages 4-5.</p>
	<p>risk of case, 22.0% lower, HR 0.78, $p = 0.01$, treatment 163 of 2,296 (7.1%), control 326 of 3,407 (9.6%), NNT 40, multivariate, patients with CKD stages 4-5.</p>
<p><i>Parant</i>, 4/14/2022, retrospective, France, peer-reviewed, median age 78.0, 12 authors, study period 1 March, 2020 - 30 June, 2020, dosage varies, trial NCT04877509 (history).</p>	<p>risk of death, 50.5% lower, RR 0.50, $p = 0.11$, treatment 7 of 66 (10.6%), control 28 of 162 (17.3%), adjusted per study, odds ratio converted to relative risk, multivariable.</p>
	<p>risk of ICU admission, 51.2% lower, RR 0.49, $p = 0.008$, treatment 10 of 66 (15.2%), control 74 of 162 (45.7%), NNT 3.3, adjusted per study, odds ratio converted to relative risk, multivariable.</p>
	<p>risk of severe case, 38.7% lower, RR 0.61, $p = 0.01$, treatment 19 of 66 (28.8%), control 86 of 162 (53.1%), NNT 4.1, adjusted per study, odds ratio converted to relative risk, multivariable.</p>
<p><i>Pecina</i>, 8/27/2021, retrospective, USA, peer-reviewed, 4 authors, dosage not specified, excluded in exclusion analyses: unadjusted results with no group details.</p>	<p>risk of death, 70.0% higher, OR 1.70, $p = 0.52$, treatment 29, control 63, supplementation, unadjusted, RR approximated with OR.</p>
	<p>risk of mechanical ventilation, 10.0% higher, OR 1.10, $p = 0.89$, treatment 29, control 63, supplementation, unadjusted, RR approximated with OR.</p>
	<p>risk of ICU admission, 30.0% higher, OR 1.30, $p = 0.61$, treatment 29, control 63, supplementation, unadjusted, RR approximated with OR.</p>
<p><i>Regalia</i>, 1/13/2022, retrospective, Italy, peer-reviewed, 10 authors, dosage varies.</p>	<p>risk of case, 33.0% lower, OR 0.67, $p = 0.21$, treatment 32 of 60 (53.3%) cases, 75 of 119 (63.0%) controls, NNT 11, case control OR, vitamin D supplementation for ≥ 3 months in the last year.</p>

<p><i>Sainz-Amo</i>, 10/24/2020, retrospective, Spain, peer-reviewed, mean age 74.5, 13 authors, dosage not specified.</p>	<p>risk of severe case, 32.7% lower, OR 0.67, $p = 0.45$, treatment 5 of 29 (17.2%) cases, 43 of 182 (23.6%) controls, NNT 23, case control OR.</p>
	<p>risk of case, 43.7% lower, OR 0.56, $p = 0.23$, treatment 6 of 39 (15.4%) cases, 42 of 172 (24.4%) controls, NNT 13, case control OR.</p>
<p><i>Sharif</i>, 11/26/2022, retrospective, Bangladesh, peer-reviewed, 14 authors, study period 13 December, 2020 - 4 February, 2021, dosage 2,000IU daily.</p>	<p>risk of severe case, 28.0% lower, OR 0.72, $p = 0.001$, adjusted per study, multivariable, RR approximated with OR.</p>
	<p>risk of severe case, 97.0% lower, OR 0.03, $p = 0.005$, adjusted per study, combined use of vitamin C, vitamin D, and zinc, multivariable, RR approximated with OR.</p>
<p><i>Shehab</i>, 2/28/2022, retrospective, multiple countries, peer-reviewed, survey, 7 authors, study period September 2020 - March 2021, dosage not specified, excluded in exclusion analyses: unadjusted results with no group details.</p>	<p>risk of severe case, 45.7% lower, RR 0.54, $p = 0.20$, treatment 6 of 90 (6.7%), control 20 of 163 (12.3%), NNT 18, unadjusted, severe vs. mild cases.</p>
<p><i>Sinaci</i>, 8/11/2021, retrospective, Turkey, peer-reviewed, 10 authors, dosage not specified.</p>	<p>risk of severe case, 90.0% lower, RR 0.10, $p = 0.35$, treatment 0 of 36 (0.0%), control 7 of 123 (5.7%), NNT 18, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm), supplementation.</p>
	<p>risk of moderate/severe case, 18.8% higher, RR 1.19, $p = 0.64$, treatment 8 of 36 (22.2%), control 23 of 123 (18.7%), supplementation.</p>
<p><i>Subramanian</i>, 1/31/2022, prospective, United Kingdom, peer-reviewed, 16 authors, dosage not specified.</p>	<p>risk of death, 27.3% lower, RR 0.73, $p = 0.12$, treatment 31 of 131 (23.7%), control 80 of 336 (23.8%), adjusted per study, odds ratio converted to relative risk, prescribed supplement use, multivariable.</p>
<p><i>Sulli (B)</i>, 2/24/2021, retrospective, Italy, peer-reviewed, 10 authors, dosage not specified.</p>	<p>risk of case, 75.6% lower, OR 0.24, $p < 0.001$, treatment 22 of 65 (33.8%) cases, 44 of 65 (67.7%) controls, NNT 3.0, case control OR, vitamin D supplementation.</p>
<p><i>Tylicki</i>, 1/6/2022, retrospective, Poland, peer-reviewed, 10 authors, study period 6 October, 2020 - 28 February, 2021, dosage not specified.</p>	<p>risk of death, 14.4% lower, RR 0.86, $p = 0.61$, treatment 28 of 85 (32.9%), control 25 of 48 (52.1%), NNT 5.2, adjusted per study, odds ratio converted to relative risk, multivariable.</p>
<p><i>Ullah</i>, 3/4/2021, retrospective, United Kingdom, peer-reviewed, 3 authors, dosage not specified, excluded in exclusion analyses: significant unadjusted confounding possible.</p>	<p>risk of death, 42.1% higher, RR 1.42, $p = 0.34$, treatment 21 of 64 (32.8%), control 26 of 135 (19.3%), adjusted per study, odds ratio converted to relative risk.</p>
	<p>risk of case, 146.0% higher, RR 2.46, $p < 0.001$, treatment 69 of 2,168 (3.2%), control 139 of 12,681 (1.1%), adjusted per study, odds ratio converted to relative risk.</p>
<p><i>van Helmond</i>, 9/17/2022, prospective, USA, peer-reviewed, 14 authors, study period 27 October, 2020 - 31 January, 2021, dosage 5,000IU daily, trial NCT04596657 (history).</p>	<p>risk of case, 97.5% lower, RR 0.02, $p = 0.07$, treatment 0 of 255 (0.0%), control 36 of 2,827 (1.3%), NNT 79, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm).</p>

<p><i>Vasheghani (B)</i>, 1/18/2021, retrospective, Iran, preprint, 6 authors, dosage not specified.</p>	<p>risk of death, 30.4% lower, RR 0.70, $p = 0.45$, treatment 7 of 88 (8.0%), control 48 of 420 (11.4%), NNT 29, vitamin D supplementation.</p>
	<p>risk of ICU admission, 63.8% lower, RR 0.36, $p = 0.009$, treatment 13 of 185 (7.0%), control 53 of 323 (16.4%), NNT 11, adjusted per study, inverted to make $RR < 1$ favor treatment, vitamin D levels $> 30\text{ng/mL}$.</p>
<p><i>Villasis-Keever</i>, 4/18/2022, Double Blind Randomized Controlled Trial, placebo-controlled, Mexico, peer-reviewed, 16 authors, study period 15 July, 2020 - 30 December, 2020, dosage 4,000IU daily.</p>	<p>risk of hospitalization, 66.5% lower, RR 0.33, $p = 1.00$, treatment 0 of 150 (0.0%), control 1 of 152 (0.7%), NNT 152, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm), ITT.</p>
	<p>risk of case, 78.0% lower, RR 0.22, $p = 0.001$, treatment 7 of 150 (4.7%), control 26 of 152 (17.1%), NNT 8.0, adjusted per study, multivariable, Table 3.</p>
<p><i>Wang</i>, 3/29/2023, Randomized Controlled Trial, China, preprint, median age 36.5, 23 authors, study period 18 December, 2022 - 20 February, 2023, dosage 200,000IU days 1, 14, trial NCT05673980 (history).</p>	<p>risk of progression, 22.6% lower, RR 0.77, $p = 0.20$, treatment 99, control 103, all severe symptoms below combined via meta analysis.</p>
	<p>risk of case, 9.0% lower, RR 0.91, $p = 0.57$, treatment 49 of 99 (49.5%), control 56 of 103 (54.4%), NNT 21.</p>
	<p>risk of case, 12.3% higher, RR 1.12, $p = 0.56$, treatment 41 of 99 (41.4%), control 38 of 103 (36.9%), first two weeks.</p>
	<p>risk of case, 53.8% lower, RR 0.46, $p = 0.06$, treatment 8 of 99 (8.1%), control 18 of 103 (17.5%), NNT 11, last two weeks.</p>
	<p>risk of progression, 4.0% higher, RR 1.04, $p = 1.00$, treatment 5 of 99 (5.1%), control 5 of 103 (4.9%), risk of severe case, fever.</p>
	<p>risk of progression, 7.5% lower, RR 0.92, $p = 1.00$, treatment 8 of 99 (8.1%), control 9 of 103 (8.7%), NNT 152, risk of severe case, sore throat.</p>
	<p>risk of progression, 42.2% lower, RR 0.58, $p = 0.41$, treatment 5 of 99 (5.1%), control 9 of 103 (8.7%), NNT 27, risk of severe case, rhinorrhea or congestion.</p>
	<p>risk of progression, 66.2% lower, RR 0.34, $p = 1.00$, treatment 0 of 99 (0.0%), control 1 of 103 (1.0%), NNT 103, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm), risk of severe case, diarrhea.</p>
	<p>risk of progression, 204.0% higher, RR 3.04, $p = 0.49$, treatment 1 of 99 (1.0%), control 0 of 103 (0.0%), continuity correction due to zero event (with reciprocal of the contrasting arm), risk of severe case, vomiting.</p>
	<p>risk of progression, 13.3% lower, RR 0.87, $p = 0.82$, treatment 10 of 99 (10.1%), control 12 of 103 (11.7%), NNT 65, risk of severe case, cough.</p>

	<p>risk of progression, 48.0% lower, RR 0.52, $p = 0.13$, treatment 8 of 99 (8.1%), control 16 of 103 (15.5%), NNT 13, risk of severe case, muscle/joint aches.</p> <p>risk of progression, 56.1% higher, RR 1.56, $p = 0.68$, treatment 3 of 99 (3.0%), control 2 of 103 (1.9%), risk of severe case, taste/smell.</p>
Ünsal (B), 4/5/2021, retrospective, Turkey, peer-reviewed, 10 authors, dosage varies.	risk of pneumonia, 71.4% lower, RR 0.29, $p = 0.009$, treatment 4 of 28 (14.3%), control 14 of 28 (50.0%), NNT 2.8, average 800-1000IU/day cholecalciferol.
Şengül, 12/31/2022, retrospective, Turkey, peer-reviewed, 4 authors, study period March 2020 - December 2021, dosage not specified.	risk of case, 68.5% lower, OR 0.31, $p = 0.004$, treatment 8 of 54 (14.8%) cases, 94 of 264 (35.6%) controls, NNT 7.4, case control OR.

References

1. **Abdollahi** et al., *The Association Between the Level of Serum 25(OH) Vitamin D, Obesity, and underlying Diseases with the risk of Developing COVID-19 Infection: A case-control study of hospitalized patients in Tehran, Iran*, Journal of Medical Virology, doi:10.1002/jmv.26726.
2. **Abdrabbo AlYafei** et al., *Association of Serum Vitamin D level and COVID-19 infection: A Case-control Study*, Qatar Medical Journal, doi:10.5339/qmj.2022.48.
3. **Abdulameer** et al., *The vitamin D binding protein gene polymorphism association with Covid-19-infected Iraqi patients*, Cellular and Molecular Biology, doi:10.14715/cmb/2023.69.5.5.
4. **Abdulateef** et al., *COVID-19 severity in relation to sociodemographics and vitamin D use*, Open Medicine, doi:10.1515/med-2021-0273.
5. **Abdulrahman** et al., *Correlates of poor clinical outcomes related to COVID-19 among older people with psychiatric illness - a mixed methods study*, The International Journal of Psychiatry in Medicine, doi:10.1177/00912174231171220.
6. **Abrishami** et al., *Possible association of vitamin D status with lung involvement and outcome in patients with COVID-19: a retrospective study*, European Journal of Nutrition, doi:10.1007/s00394-020-02411-0.
7. **Aci** et al., *Effect of vitamin D receptor gene BsmI polymorphism on hospitalization of SARS-CoV-2 positive patients*, Nucleosides, Nucleotides & Nucleic Acids, doi:10.1080/15257770.2023.2253281.
8. **Afaghi** et al., *Prevalence and Clinical Outcomes of Vitamin D Deficiency in COVID-19 Hospitalized Patients: A Retrospective Single-Center Analysis*, The Tohoku Journal of Experimental Medicine, doi:10.1620/tjem.255.127.
9. **Ahmed** et al., *Causal Inference and COVID-19 Nursing Home Patients: Identifying Factors That Reduced Mortality Risk*, medRxiv, doi:10.1101/2021.11.18.21266489.
10. **Al Sulaiman** et al., *Survival implications vs. complications: unraveling the impact of vitamin D adjunctive use in critically ill patients with COVID-19—A multicenter cohort study*, Frontiers in Medicine, doi:10.3389/fmed.2023.1237903.
11. **Al-Daghri** et al., *Vitamin D status of Arab Gulf residents screened for SARS-CoV-2 and its association with COVID-19 infection: a multi-centre case-control study*, Journal of Translational Medicine, doi:10.1186/s12967-021-02838-x.
12. **Al-Jarallah** et al., *In-hospital mortality in SARS-CoV-2 stratified by serum 25-hydroxy-vitamin D levels: A retrospective study*, Journal of Medical Virology, doi:10.1002/jmv.27133.
13. **Al-Mazaideh** et al., *Vitamin D is a New Promising Inhibitor to the Main Protease (Mpro) of COVID-19 by Molecular Docking*, Journal of Pharmaceutical Research International, doi:10.9734/jpri/2021/v33i29B31603.

14. **Al-Salman** et al., *In COVID-19 patients, low 25-hydroxyvitamin D level in serum is associated with longer viral clearance time and higher risk of intensive care unit admission*, Nutrition & Food Science, doi:10.1108/NFS-05-2021-0143.
15. **Alarslan** et al., *Vitamin D levels and disease severity in COVID-19*, Medical Journal of İzmir Hospital, 26:3, bozyakaeah.saglik.gov.tr/Eklenti/306811/0/tip-2022—3—91-98pdf.pdf.
16. **Alcala-Diaz** et al., *Calcifediol Treatment and Hospital Mortality Due to COVID-19: A Cohort Study*, Nutrients, doi:10.3390/nu13061760.
17. **Aldwihi** et al., *Patients' Behavior Regarding Dietary or Herbal Supplements before and during COVID-19 in Saudi Arabia*, International Journal of Environmental Research and Public Health, doi:10.3390/ijerph18105086.
18. **Alguwaihes** et al., *Diabetes and Covid-19 among hospitalized patients in Saudi Arabia: a single-centre retrospective study*, Cardiovascular Diabetology, doi:10.1186/s12933-020-01184-4.
19. **AlKhafaji** et al., *The Impact of Vitamin D Level on the Severity and Outcome of Hospitalized Patients with COVID-19 Disease*, International Journal of General Medicine, doi:10.2147/ijgm.s346169.
20. **Allami** et al., *The risk of up normal values of two parameters obesity and vitamin D in incidence of coronavirus disease-19 among Iraqi patients*, 1st Samarra International Conference for Pure and Applied Sciences (SICPS2021), doi:10.1063/5.0121166.
21. **Alpcan** et al., *Vitamin D levels in children with COVID-19: a report from Turkey*, Epidemiology & Infection, doi:10.1017/S0950268821001825.
22. **AlSafar** et al., *COVID-19 Disease Severity and Death in Relation to Vitamin D Status among SARS-CoV-2-Positive UAE Residents*, Nutrients, doi:10.3390/nu13051714.
23. **Alsaidi** et al., *Griffithsin and Carrageenan Combination Results in Antiviral Synergy against SARS-CoV-1 and 2 in a Pseudoviral Model*, Marine Drugs, doi:10.3390/md19080418.
24. **Altman**, D., *How to obtain the P value from a confidence interval*, BMJ, doi:10.1136/bmj.d2304.
25. **Altman (B)** et al., *How to obtain the confidence interval from a P value*, BMJ, doi:10.1136/bmj.d2090.
26. **Álvarez** et al., *Vitamin D deficiency and SARS-CoV-2 infection: Big-data analysis from March 2020 to March 2021. D-COVID study*, bioRxiv, doi:10.1101/2022.10.27.514012.
27. **Alzahrani** et al., *The Association Between Vitamin D Serum Level and COVID-19 Patients' Outcomes in a Tertiary Center in Saudi Arabia: A Retrospective Cohort Study*, Cureus, doi:10.7759/cureus.26266.
28. **Amin** et al., *No evidence that vitamin D is able to prevent or affect the severity of COVID-19 in individuals with European ancestry: a Mendelian randomisation study of open data*, BMJ Nutrition, Prevention & Health, doi:10.1136/bmjnp-2020-000151.
29. **Andreani** et al., *In vitro testing of combined hydroxychloroquine and azithromycin on SARS-CoV-2 shows synergistic effect*, Microbial Pathogenesis, doi:10.1016/j.micpath.2020.104228.
30. **Angelidi** et al., *Vitamin D Status is Associated With In-hospital Mortality and Mechanical Ventilation: A Cohort of COVID-19 Hospitalized Patients*, Mayo Clinic Proceedings, doi:10.1016/j.mayocp.2021.01.001.
31. **Anglemeyer** et al., *Healthcare outcomes assessed with observational study designs compared with those assessed in randomized trials*, Cochrane Database of Systematic Reviews 2014, Issue 4, doi:10.1002/14651858.MR000034.pub2.
32. **Anjum** et al., *Examine the association between severe vitamin D deficiency and mortality in patients with Covid-19*, Pakistan J. Med. Heal. Sci., 14:3, pjmhsonline.com/2020/july-sep/1184.pdf.
33. **Annweiler** et al., *Vitamin D Supplementation Associated to Better Survival in Hospitalized Frail Elderly COVID-19 Patients: The GERIA-COVID Quasi-Experimental Study*, Nutrients, doi:10.3390/nu12113377.
34. **Annweiler (B)** et al., *Vitamin D and survival in COVID-19 patients: A quasi-experimental study*, The Journal of Steroid Biochemistry and Molecular Biology, doi:10.1016/j.jsbmb.2020.105771.
35. **Annweiler (C)** et al., *Vitamin D Supplementation Associated to Better Survival in Hospitalized Frail Elderly COVID-19 Patients: The GERIA-COVID Quasi-Experimental Study*, Nutrients, doi:10.3390/nu12113377.

36. **Ansari** et al., *Frequency of Severe Vitamin D Deficiency and its Association with Mortality in Patients with Corona virus Disease*, Pakistan J. Med. Heal. Sci., 14:4, pjmhsonline.com/2020/oct_dec/1206.pdf.
37. **Arabadzhiyska** et al., *Serum vitamin D levels and inflammatory status in COVID-19 patients*, Bratislava Medical Journal, doi:10.4149/bll_2023_069.
38. **Arabi** et al., *The association between vitamin D3 deficiency and acute kidney injury in COVID-19 patients*, Journal of Renal Injury Prevention, doi:10.34172/jrip.2022.32126.
39. **Argano** et al., *Protective Effect of Vitamin D Supplementation on COVID-19-Related Intensive Care Hospitalization and Mortality: Definitive Evidence from Meta-Analysis and Trial Sequential Analysis*, Pharmaceuticals, doi:10.3390/ph16010130.
40. **Arroyo-Díaz** et al., *Previous Vitamin D Supplementation and Morbidity and Mortality Outcomes in People Hospitalised for COVID19: A Cross-Sectional Study*, Frontiers in Public Health, doi:10.3389/fpubh.2021.758347.
41. **Asgari** et al., *Vitamin D Insufficiency in Disease Severity and Prognosis of the Patients With SARS Corona Virus-2 Infection*, Acta Medica Iranica, doi:10.18502/acta.v59i11.7779.
42. **Asghar** et al., *Evaluation of Vitamin-D Status and Its Association with Clinical Outcomes Among COVID-19 Patients in Pakistan*, Am. J. Trop. Med. Hyg., doi:10.4269/ajtmh.21-0577.
43. **Asimi** et al., *Selenium, zinc, and vitamin D supplementation affect the clinical course of COVID-19 infection in Hashimoto's thyroiditis*, Endocrine Abstracts, doi:10.1530/endoabs.73.PEP14.2.
44. **Assiri** et al., *COVID-19 related treatment and outcomes among COVID-19 ICU patients: A retrospective cohort study*, Journal of Infection and Public Health, doi:10.1016/j.jiph.2021.08.030.
45. **Atanasovska** et al., *Vitamin D levels and oxidative stress markers in patients hospitalized with COVID-19*, Redox Report , doi:10.1080/13510002.2021.1999126.
46. **Aweimer** et al., *Mortality rates of severe COVID-19-related respiratory failure with and without extracorporeal membrane oxygenation in the Middle Ruhr Region of Germany*, Scientific Reports, doi:10.1038/s41598-023-31944-7.
47. **Azadeh** et al., *Serum Vitamin D Concentrations in CoVID19 Patients*, J. Mazandaran Univ. Med. Sci. 31:195, jmums.mazums.ac.ir/article-1-16104-en.html.
48. **Bader** et al., *The Effect of Weekly 50,000 IU Vitamin D3 Supplements on the Serum Levels of Selected Cytokines Involved in Cytokine Storm: A Randomized Clinical Trial in Adults with Vitamin D Deficiency*, Nutrients, doi:10.3390/nu15051188.
49. **Bagheri** et al., *Supplement Usage Pattern in a Group of COVID-19 Patients in Tehran*, Journal of Family & Reproductive Health, doi:10.18502/jfrh.v14i3.4668.
50. **Baguma** et al., *Characteristics of the COVID-19 patients treated at Gulu Regional Referral Hospital, Northern Uganda: A cross-sectional study*, Research Square, doi:10.21203/rs.3.rs-1193578/v1.
51. **Baguma (B)** et al., *Characteristics of the COVID-19 patients treated at Gulu Regional Referral Hospital, Northern Uganda: A cross-sectional study*, Research Square, doi:10.21203/rs.3.rs-1193578/v1.
52. **Bakaloudi** et al., *A critical update on the role of mild and serious vitamin D deficiency prevalence and the COVID-19 epidemic in Europe*, Nutrition, doi:10.1016/j.nut.2021.111441.
53. **Baktash** et al., *Vitamin D status and outcomes for hospitalised older patients with COVID-19*, Postgraduate Medical Journal, doi:10.1136/postgradmedj-2020-138712.
54. **Baralić** et al., *Significance of 1,25-Dihydroxyvitamin D3 on Overall Mortality in Peritoneal Dialysis Patients with COVID-19*, Nutrients, doi:10.3390/nu15092050.
55. **Barassi** et al., *Vitamin D in severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) patients with non-invasive ventilation support*, Panminerva Med., doi:10.23736/S0031-0808.21.04277-4.
56. **Barrett** et al., *Vitamin D Status and Mortality from SARS CoV-2: A Prospective Study of Unvaccinated Caucasian Adults*, Nutrients, doi:10.3390/nu14163252.
57. **Basaran** et al., *The relationship between vitamin D and the severity of COVID-19*, Bratislava Medical Journal, doi:10.4149/bll_2021_034.

58. **Basińska-Lewandowska** et al., *Frequency of COVID-19 Infection as a Function of Vitamin D Levels*, *Nutrients*, doi:10.3390/nu15071581.
59. **Batur** et al., *Association between Vitamin D Status and Secondary Infections in Patients with Severe COVID-19 Admitted in the Intensive Care Unit of a Tertiary-Level Hospital in Turkey*, *Diagnostics*, doi:10.3390/diagnostics13010059.
60. **Baykal** et al., *Correlation of vitamin D level with the clinical-radiological severity of COVID-19 in geriatric patients*, *Journal of Health Sciences and Medicine*, doi:10.32322/jhsm.1063405.
61. **Bayrak** et al., *Association Between Vitamin D Levels and COVID-19 Infection in Children: A Case-Control Study*, *Turkish Archives of Pediatrics*, doi:10.5152/turkarchpediatr.2023.22217.
62. **Bayramoğlu** et al., *The association between vitamin D levels and the clinical severity and inflammation markers in pediatric COVID-19 patients: single-center experience from a pandemic hospital*, *European Journal of Pediatrics*, doi:10.1007/s00431-021-04030-1.
63. **Beheshti** et al., *Correlation of vitamin D levels with serum parameters in Covid-19 patients*, *Clinical Nutrition ESPEN*, doi:10.1016/j.clnesp.2023.04.012.
64. **Beigmohammadi** et al., *The effect of supplementation with vitamins A, B, C, D, and E on disease severity and inflammatory responses in patients with COVID-19: a randomized clinical trial*, *Trials*, doi:10.1186/s13063-021-05795-4.
65. **Benbouar** et al., *Vitamin D Deficiency and Low Serum Calcium as Predictors of Poor Prognosis in Patients with Severe COVID-19*, *Journal of the American College of Nutrition*, doi:10.1080/07315724.2020.1856013.
66. **Bhat** et al., *Effect of calcifediol supplementation as add-on therapy on the immune repertoire in recipients of the ChAdOx1 nCoV-19 vaccine: A prospective open-label, placebo-controlled, clinical trial*, *Journal of Infection*, doi:10.1016/j.jinf.2023.03.004.
67. **Biancatelli** et al., *Quercetin and Vitamin C: An Experimental, Synergistic Therapy for the Prevention and Treatment of SARS-CoV-2 Related Disease (COVID-19)*, *Frontiers in Immunology*, doi:10.3389/fimmu.2020.01451.
68. **Bianconi** et al., *Prevalence of vitamin D deficiency and its prognostic impact on patients hospitalized with COVID-19*, *Nutrition*, doi:10.1016/j.nut.2021.111408.
69. **Bishop** et al., *REScue Trial: Randomized Controlled Clinical Trial with Extended-Release Calcifediol in Symptomatic COVID-19 Outpatients*, *Nutrition*, doi:10.1016/j.nut.2022.111899.
70. **Blanch-Rubió** et al., *Influence of anti-osteoporosis treatments on the incidence of COVID-19 in patients with non-inflammatory rheumatic conditions*, *Aging*, doi:10.18632/aging.104117.
71. **Bogliolo** et al., *Vitamin D 25OH Deficiency and Mortality in Moderate to Severe COVID-19: A Multi-Center Prospective Observational Study*, *Frontiers in Nutrition*, doi:10.3389/fnut.2022.934258.
72. **Bogomaz** et al., *Vitamin D as a predictor of negative outcomes in hospitalized COVID-19 patients: An observational study*, *Canadian Journal of Respiratory Therapy*, doi:10.29390/001c.87408.
73. **Boukef** et al., *Melatonin, Vitamins and Minerals Supplements for the Treatment of Covid-19 and Covid-like Illness: Results of a Prospective, Randomised, Double-blinded Multicentre Study*, NCT05670444, *clinicaltrials.gov/study/NCT05670444*.
74. **Boulware**, D., *Comments regarding paper rejection*, *twitter.com/boulware_dr/status/1311331372884205570*.
75. **Breslin** et al., *The relationship between vitamin D, biomarkers and clinical outcome in hospitalised Covid-19 patients*, *Proceedings of the Nutrition Society*, doi:10.1017/S0029665121002214.
76. **Brunvoll** et al., *Prevention of covid-19 and other acute respiratory infections with cod liver oil supplementation, a low dose vitamin D supplement: quadruple blinded, randomised placebo controlled trial*, *BMJ*, doi:10.1136/bmj-2022-071245.
77. **Bucurica** et al., *Association of Vitamin D Deficiency and Insufficiency with Pathology in Hospitalized Patients*, *Diagnostics*, doi:10.3390/diagnostics13050998.
78. **Burahee** et al., *Older patients with proximal femur fractures and SARS-CoV-2 infection – An observational study*, *SICOT-J*, doi:10.1051/sicotj/2021001.

79. **Bushnaq** et al., *The Impact of Vitamin D Status on COVID-19 Severity among Hospitalized Patients in the Western Region of Saudi Arabia: A Retrospective Cross-Sectional Study*, International Journal of Environmental Research and Public Health, doi:10.3390/ijerph19031901.
80. **Butler-Laporte** et al., *Vitamin D and COVID-19 susceptibility and severity in the COVID-19 Host Genetics Initiative: A Mendelian randomization study*, PLOS Medicine, doi:10.1371/journal.pmed.1003605.
81. **Bychinin** et al., *Effect of vitamin D3 supplementation on cellular immunity and inflammatory markers in COVID-19 patients admitted to the ICU*, Scientific Reports, doi:10.1038/s41598-022-22045-y.
82. **Bychinin (B)** et al., *Prevalence of hypovitaminosis D in COVID-19 patients in the intensive care unit*, Journal of Clinical Practice, doi:10.17816/clinpract64976.
83. **c19early.org**, c19early.org/jmeta.html.
84. **c19early.org (B)**, c19early.org/exmeta.html.
85. **c19early.org (C)**, c19early.org/files/lakkireddy-response.zip.
86. **Campi** et al., *Vitamin D and COVID-19 severity and related mortality: a prospective study in Italy*, BMC Infectious Diseases, doi:10.1186/s12879-021-06281-7.
87. **Cangiano** et al., *Mortality in an Italian nursing home during COVID-19 pandemic: correlation with gender, age, ADL, vitamin D supplementation, and limitations of the diagnostic tests*, Aging, doi:10.18632/aging.202307.
88. **Cannata-Andía** et al., *A single-oral bolus of 100,000 IU of cholecalciferol at hospital admission did not improve outcomes in the COVID-19 disease: the COVID-VIT-D — a randomised multicentre international clinical trial*, BMC Medicine, doi:10.1186/s12916-022-02290-8.
89. **Carlberg** et al., *In vivo response of the human epigenome to vitamin D: A Proof-of-principle study*, The Journal of Steroid Biochemistry and Molecular Biology, doi:10.1016/j.jsbmb.2018.01.002.
90. **Carpagnano** et al., *Vitamin D deficiency as a predictor of poor prognosis in patients with acute respiratory failure due to COVID-19*, J. Endocrinol. Invest., 2020, Aug 9, 1-7, doi:10.1007/s40618-020-01370-x.
91. **Castillo** et al., *Effect of calcifediol treatment and best available therapy versus best available therapy on intensive care unit admission and mortality among patients hospitalized for COVID-19: A pilot randomized clinical study*, Journal of Steroid Biochemistry and Molecular Biology, 203, October 2020, doi:10.1016/j.jsbmb.2020.105751.
92. **Cereda** et al., *Vitamin D 25OH deficiency in COVID-19 patients admitted to a tertiary referral hospital*, Clinical Nutrition (Edinburgh, Scotland), doi:10.1016/j.clnu.2020.10.055.
93. **Cereda (B)** et al., *Vitamin D supplementation and outcomes in coronavirus disease 2019 (COVID-19) patients from the outbreak area of Lombardy, Italy*, Nutrition, doi:10.1016/j.nut.2020.111055.
94. **Cetin Ozbek** et al., *Does the Level of Vitamin D in COVID-19 Patients Affect the Survival and Duration of Hospital Stay?*, Clinical Science of Nutrition, doi:10.5152/ClinSciNutr.2023.22059.
95. **Charkowick** et al., *Vitamin D Deficiency and Thrombosis in Hospitalized SARS-CoV-2 Patients with Suspected Pulmonary Embolism*, AJRCCM Conference, www.atsjournals.org/doi/pdf/10.1164/ajrccm-conference.2022.205.1_MeetingAbstracts.A4571.
96. **Charla** et al., *Is suboptimal circulating level of vitamin D a risk factor for the poor prognosis of COVID-19? – A comparison of first and second waves in India*, Research Square, doi:10.21203/rs.3.rs-1826271/v1.
97. **Charoenngam** et al., *Association of vitamin D status with hospital morbidity and mortality in adult hospitalized COVID-19 patients*, Endocrine Practice, doi:10.1016/j.eprac.2021.02.013.
98. **Chellasamy** et al., *Docking and molecular dynamics studies of human ezrin protein with a modelled SARS-CoV-2 endodomain and their interaction with potential invasion inhibitors*, Journal of King Saud University - Science, doi:10.1016/j.jksus.2022.102277.
99. **Chen** et al., *Plasma 25(OH)D Level is Associated with the Nucleic Acid Negative Conversion Time of COVID-19 Patients: An Exploratory Study*, Infection and Drug Resistance, doi:10.2147/idr.s400561.

100. **Chiodini** et al., *Vitamin D Status and SARS-CoV-2 Infection and COVID-19 Clinical Outcomes*, *Frontiers in Public Health*, doi:10.3389/fpubh.2021.736665.
101. **Chodick** et al., *Angiotensin-converting enzyme inhibitors and angiotensin-receptor blockers are not associated with increased risk of SARS-CoV-2 infection*, *Journal of Travel Medicine*, doi:10.1093/jtm/taaa069.
102. **Concato** et al., *NEJM*, 342:1887-1892, doi:10.1056/NEJM200006223422507.
103. **Connolly** et al., *An observational study of the association of vitamin D status and other patient characteristics with COVID-19 severity and mortality*, *Proceedings of the Nutrition Society*, doi:10.1017/S0029665121002482.
104. **covid19treatmentguidelines.nih.gov**, www.covid19treatmentguidelines.nih.gov/therapies/supplements/vitamin-d/.
105. **Cozier** et al., *Lower serum 25(OH)D levels associated with higher risk of COVID-19 infection in U.S. Black women*, *PLoS ONE*, doi:10.1371/journal.pone.0255132.
106. **Crawford** et al., *Analysis of Select Dietary Supplement Products Marketed to Support or Boost the Immune System*, *JAMA Network Open*, doi:10.1001/jamanetworkopen.2022.26040.
107. **Crighon** et al., *Toxicological screening and DNA sequencing detects contamination and adulteration in regulated herbal medicines and supplements for diet, weight loss and cardiovascular health*, *Journal of Pharmaceutical and Biomedical Analysis*, doi:10.1016/j.jpba.2019.112834.
108. **D'Avolio** et al., *25-hydroxyvitamin D concentrations are lower in patients with positive PCR for SARS-CoV-2*, *Nutrients*, 12:5, 1–7, doi:10.3390/nu12051359.
109. **D'Ecclesiis** et al., *Vitamin D and SARS-CoV2 infection, severity and mortality: A systematic review and meta-analysis*, *PLOS ONE*, doi:10.1371/journal.pone.0268396.
110. **Dana** et al., *Vitamin D Level in Laboratory Confirmed COVID-19 and Disease Progression*, *The Eurasian Journal of Medicine*, doi:10.5152/eurasianjmed.2022.21088.
111. **Davoudi** et al., *Lack of association between vitamin D insufficiency and clinical outcomes of patients with COVID-19 infection*, *BMC Infectious Diseases*, doi:10.1186/s12879-021-06168-7.
112. **Davran** et al., *Relationship between vitamin D level and clinical status in COVID-19 patients*, *Konuralp Tıp Dergisi*, doi:10.18521/ktd.1134319.
113. **De Forni** et al., *Synergistic drug combinations designed to fully suppress SARS-CoV-2 in the lung of COVID-19 patients*, *PLoS ONE*, doi:10.1371/journal.pone.0276751.
114. **De Nicolò** et al., *Possible Impact of Vitamin D Status and Supplementation on SARS-CoV-2 Infection Risk and COVID-19 Symptoms in a Cohort of Patients with Inflammatory Bowel Disease*, *Nutrients*, doi:10.3390/nu15010169.
115. **De Niet** et al., *Positive Effects of Vitamin D Supplementation in Patients Hospitalized for COVID-19: A Randomized, Double-Blind, Placebo-Controlled Trial*, *Nutrients*, doi:10.3390/nu14153048.
116. **De Smet** et al., *Serum 25(OH)D Level on Hospital Admission Associated With COVID-19 Stage and Mortality*, *American Journal of Clinical Pathology*, doi:10.1093/ajcp/aqaa252.
117. **Deaton** et al., *Understanding and misunderstanding randomized controlled trials*, *Social Science & Medicine*, 210, doi:10.1016/j.socscimed.2017.12.005.
118. **Demir** et al., *Vitamin D deficiency is associated with COVID-19 positivity and the severity of the disease*, *Journal of Medical Virology*, doi:10.1002/jmv.26832.
119. **Deng**, H., *PyMeta*, *Python module for meta-analysis*, www.pymeta.com/.
120. **Derakhshanian** et al., *The predictive power of serum vitamin D for poor outcomes in COVID-19 patients*, *Food Science & Nutrition*, doi:10.1002/fsn3.2591.
121. **Desai** et al., *Vitamin K & D Deficiencies Are Independently Associated With COVID-19 Disease Severity*, *Open Forum Infectious Diseases*, doi:10.1093/ofid/ofab408.

122. **di Filippo** et al., *Low vitamin D levels are associated with Long COVID syndrome in COVID-19 survivors*, The Journal of Clinical Endocrinology & Metabolism, doi:10.1210/clinem/dgad207.
123. **di Filippo (B)** et al., *Vitamin D levels associate with blood glucose and BMI in COVID-19 patients predicting disease severity*, The Journal of Clinical Endocrinology & Metabolism, doi:10.1210/clinem/dgab599.
124. **Diaz-Curiel** et al., *The relationship between 25(OH) vitamin D levels and COVID-19 onset and disease course in Spanish patients*, Journal of Steroid Biochemistry and Molecular Biology, doi:10.1016/j.jsbmb.2021.105928.
125. **DiGuilio** et al., *The multiphasic TNF- α -induced compromise of Calu-3 airway epithelial barrier function*, Experimental Lung Research, doi:10.1080/01902148.2023.2193637.
126. **Din Ujjan** et al., *The possible therapeutic role of curcumin and quercetin in the early-stage of COVID-19—Results from a pragmatic randomized clinical trial*, Frontiers in Nutrition, doi:10.3389/fnut.2022.1023997.
127. **Doğan** et al., *The Clinical Significance of Vitamin D and Zinc Levels with Respect to Immune Response in COVID-19 Positive Children*, Journal of Tropical Pediatrics, doi:10.1093/tropej/fmac072.
128. **Domazet Bugarin** et al., *Vitamin D Supplementation and Clinical Outcomes in Severe COVID-19 Patients—Randomized Controlled Trial*, Nutrients, doi:10.3390/nu15051234.
129. **Dror** et al., *Pre-infection 25-hydroxyvitamin D3 levels and association with severity of COVID-19 illness*, PLOS ONE, doi:10.1371/journal.pone.0263069.
130. **Dudley** et al., *Revisiting vitamin D status and supplementation for in-patients with intellectual and developmental disability in the North of England, UK*, BJPsych Bulletin, doi:10.1192/bjb.2021.55.
131. **Eden** et al., *Nutritional parameters and outcomes in patients admitted to intensive care with COVID-19: a retrospective single-centre service evaluation*, BMJ Nutrition, Prevention & Health, doi:10.1136/bmjnp-2021-000270.
132. **Efird** et al., *The Interaction of Vitamin D and Corticosteroids: A Mortality Analysis of 26,508 Veterans Who Tested Positive for SARS-CoV-2*, International Journal of Environmental Research and Public Health, doi:10.3390/ijerph19010447.
133. **Egger** et al., *Bias in meta-analysis detected by a simple, graphical test*, BMJ, doi:10.1136/bmj.315.7109.629.
134. **El Hajj** et al., *Effect of Vitamin D Supplementation on Inflammatory Markers in Non-Obese Lebanese Patients with Type 2 Diabetes: A Randomized Controlled Trial*<https://www.mdpi.com/2072-6643/12/7/2033>, Nutrients, doi:10.3390/nu12072033.
135. **Elamir** et al., *A Randomized Pilot Study Using Calcitriol in Hospitalized Patients*, Bone, doi:10.1016/j.bone.2021.116175.
136. **Elhadi** et al., *Epidemiology, outcomes, and utilization of intensive care unit resources for critically ill COVID-19 patients in Libya: A prospective multi-center cohort study*, PLOS ONE, doi:10.1371/journal.pone.0251085.
137. **Ersöz** et al., *The association between micronutrient and hemogram values and prognostic factors in COVID-19 patients: A single-center experience from Turkey*, International Journal of Clinical Practice, doi:10.1111/ijcp.14078.
138. **Espitia-Hernandez** et al., *Effects of Ivermectin-azithromycin-cholecalciferol combined therapy on COVID-19 infected patients: A proof of concept study*, Biomedical Research, 31:5, www.biomedres.info/biomedical-research/effects-of-ivermectinazithromycincholecalciferol-combined-therapy-on-covid19-infected-patients-a-proof-of-concept-study-14435.html.
139. **Fairfield** et al., *Association of Vitamin D Prescribing and Clinical Outcomes in Adults Hospitalized with COVID-19*, Nutrients, doi:10.3390/nu14153073.
140. **Faniyi** et al., *Vitamin D status and seroconversion for COVID-19 in UK healthcare workers who isolated for COVID-19 like symptoms during the 2020 pandemic*, medRxiv, doi:10.1101/2020.10.05.20206706.
141. **Faria** et al., *Genomics and epidemiology of the P.1 SARS-CoV-2 lineage in Manaus, Brazil*, Science, doi:10.1126/science.abh2644.
142. **Fasano** et al., *COVID-19 in Parkinson's Disease Patients Living in Lombardy, Italy*, Movement Disorders, doi:10.1002/mds.28176.
143. **Fatemi** et al., *Association of vitamin D deficiency with COVID-19 severity and mortality in Iranian people: a prospective observational study*, Acute and Critical Care, doi:10.4266/acc.2021.00605.

144. **Faul** et al., *Vitamin D Deficiency and ARDS after SARS-CoV-2 Infection*, Irish Medical Journal, 113:5, 84, imj.ie/vitamin-d-deficiency-and-ards-after-sars-cov-2-infection/.
145. **Fernandes de Souza** et al., *Lung Inflammation Induced by Inactivated SARS-CoV-2 in C57BL/6 Female Mice Is Controlled by Intranasal Instillation of Vitamin D*, Cells, doi:10.3390/cells12071092.
146. **Ferrer-Sánchez** et al., *Serum 25(OH) Vitamin D Levels in Pregnant Women with Coronavirus Disease 2019 (COVID-19): A Case-Control Study*, International Journal of Environmental Research and Public Health, doi:10.3390/ijerph19073965.
147. **Fiore** et al., *Effectiveness of Vitamin D Supplements among Patients Hospitalized for COVID-19: Results from a Monocentric Matched-Cohort Study*, Healthcare, doi:10.3390/healthcare10050956.
148. **Freitas** et al., *Vitamin D-related polymorphisms and vitamin D levels as risk biomarkers of COVID-19 infection severity*, medRxiv, doi:10.1101/2021.03.22.21254032.
149. **Frish** et al., *The Association of Weight Reduction and Other Variables after Bariatric Surgery with the Likelihood of SARS-CoV-2 Infection*, Journal of Clinical Medicine, doi:10.3390/jcm12124054.
150. **Galaznik** et al., *Assessment of vitamin D deficiency and COVID-19 diagnosis in patients with breast or prostate cancer using electronic medical records*, Journal of Clinical Oncology, doi:10.1200/JCO.2021.39.15_suppl.6589.
151. **Galmés** et al., *Suboptimal Consumption of Relevant Immune System Micronutrients Is Associated with a Worse Impact of COVID-19 in Spanish Populations*, Nutrients, doi:10.3390/nu14112254.
152. **Galmés (B)** et al., *Current State of Evidence: Influence of Nutritional and Nutrigenetic Factors on Immunity in the COVID-19 Pandemic Framework*, Nutrients, doi:10.3390/nu12092738.
153. **Gasmi** et al., *Quercetin in the Prevention and Treatment of Coronavirus Infections: A Focus on SARS-CoV-2*, Pharmaceuticals, doi:10.3390/ph15091049.
154. **Gaudio** et al., *Vitamin D Levels Are Reduced at the Time of Hospital Admission in Sicilian SARS-CoV-2-Positive Patients*, International Journal of Environmental Research and Public Health, doi:10.3390/ijerph18073491.
155. **Gavioli** et al., *An Evaluation of Serum 25-Hydroxy Vitamin D Levels in Patients with COVID-19 in New York City*, Journal of the American College of Nutrition, doi:10.1080/07315724.2020.1869626.
156. **Ghanei** et al., *Low serum levels of zinc and 25-hydroxyvitmain D as potential risk factors for COVID-19 susceptibility: a pilot case-control study*, European Journal of Clinical Nutrition, doi:10.1038/s41430-022-01095-5.
157. **Gholi** et al., *Vitamin D deficiency is associated with increased risk of delirium and mortality among critically ill, elderly covid-19 patients*, Complementary Therapies in Medicine, doi:10.1016/j.ctim.2022.102855.
158. **Giannini** et al., *Effectiveness of In-Hospital Cholecalciferol Use on Clinical Outcomes in Comorbid COVID-19 Patients: A Hypothesis-Generating Study*, Nutrients , doi:10.3390/nu13010219.
159. **Gibbons** et al., *Association between vitamin D supplementation and COVID-19 infection and mortality*, Scientific Reports, doi:10.1038/s41598-022-24053-4.
160. **Golabi** et al., *The Association between Vitamin D and Zinc Status and the Progression of Clinical Symptoms among Outpatients Infected with SARS-CoV-2 and Potentially Non-Infected Participants: A Cross-Sectional Study*, Nutrients, doi:10.3390/nu13103368.
161. **Golabi (B)** et al., *The Association between Vitamin D and Zinc Status and the Progression of Clinical Symptoms among Outpatients Infected with SARS-CoV-2 and Potentially Non-Infected Participants: A Cross-Sectional Study*, Nutrients, doi:10.3390/nu13103368.
162. **Gönen** et al., *Rapid and Effective Vitamin D Supplementation May Present Better Clinical Outcomes in COVID-19 (SARS-CoV-2) Patients by Altering Serum INOS1, IL1B, IFNg, Cathelicidin-LL37, and ICAM1*, Nutrients, doi:10.3390/nu13114047.
163. **Gonzalez** et al., *Vitamin D on admission and disease severity in patients with COVID-19 in the Intensive Care Unit*, Revista de Nutrición Clínica y Metabolismo, doi:10.35454/rncm.v6n2.485.
164. **González-Estevez** et al., *Association of Food Intake Quality with Vitamin D in SARS-CoV-2 Positive Patients from Mexico: A Cross-Sectional Study*, International Journal of Environmental Research and Public Health, doi:10.3390/ijerph18147266.

165. **Gøtzsche**, P., *Bias in double-blind trials*, Doctoral Thesis, University of Copenhagen, www.scientificfreedom.dk/2023/05/16/bias-in-double-blind-trials-doctoral-thesis/.
166. **Green** et al., *A higher frequency of physical activity is associated with reduced rates of SARS-CoV-2 infection*, European Journal of General Practice, doi:10.1080/13814788.2022.2138855.
167. **Guldemir** et al., *Clinical characteristics of bus drivers and field officers infected with COVID-19: A cross-sectional study from Istanbul*, Work, doi:10.3233/wor-220292.
168. **Guldemir (B)** et al., *Clinical characteristics of bus drivers and field officers infected with COVID-19: A cross-sectional study from Istanbul*, Work, doi:10.3233/wor-220292.
169. **Gupta** et al., *Temporal Association of Reduced Serum Vitamin D with COVID-19 Infection: Two Single-Institution Case–Control Studies*, Nutrients, doi:10.3390/nu14132757.
170. **Güven** et al., *The effect of high-dose parenteral vitamin D3 on COVID-19-related in-hospital mortality in critical COVID-19 patients during intensive care unit admission: an observational cohort study*, European Journal of Clinical Nutrition, doi:10.1038/s41430-021-00984-5.
171. **Hafez** et al., *Vitamin D Status in Relation to the Clinical Outcome of Hospitalized COVID-19 Patients*, Frontiers in Medicine, doi:10.3389/fmed.2022.843737.
172. **Hafez (B)** et al., *Factors Influencing Disease Stability and Response to Tocilizumab Therapy in Severe COVID-19: A Retrospective Cohort Study*, Antibiotics, doi:10.3390/antibiotics11081078.
173. **Hafezi** et al., *Vitamin D enhances type I IFN signaling in COVID-19 patients*, Scientific Reports, doi:10.1038/s41598-022-22307-9.
174. **Harbord** et al., *A modified test for small-study effects in meta-analyses of controlled trials with binary endpoints*, Statistics in Medicine, doi:10.1002/sim.2380.
175. **Hariyanto** et al., *Vitamin D supplementation and Covid-19 outcomes: A systematic review, meta-analysis and meta-regression*, Reviews in Medical Virology, doi:10.1002/rmv.2269.
176. **Hastie** et al., *Vitamin D concentrations and COVID-19 infection in UK Biobank*, Diabetes and Metabolic Syndrome: Clinical Research and Reviews, 14:4, 561–565, doi:10.1016/j.dsx.2020.04.050.
177. **Hayden** et al., *Baloxavir Marboxil for Uncomplicated Influenza in Adults and Adolescents*, New England Journal of Medicine, doi:10.1056/NEJMoa1716197.
178. **Hermawan** et al., *Association between 25(OH)D3 Levels and the Presence of COVID-19 Symptoms*, Molecular and Cellular Biomedical Sciences, doi:10.21705/mcbs.v7i1.306.
179. **Hernández** et al., *Vitamin D Status in Hospitalized Patients with SARS-CoV-2 Infection*, The Journal of Clinical Endocrinology & Metabolism, doi:10.1210/clinem/dgaa733.
180. **Hernández (B)** et al., *Vitamin D Status in Hospitalized Patients with SARS-CoV-2 Infection*, The Journal of Clinical Endocrinology & Metabolism, doi:10.1210/clinem/dgaa733.
181. **Hogarth** et al., *Clinical Characteristics and Comorbidities associated with SARS-CoV-2 breakthrough infection in the University of California Healthcare Systems*, The American Journal of the Medical Sciences, doi:10.1016/j.amjms.2023.04.019.
182. **Holt** et al., *Risk factors for developing COVID-19: a population-based longitudinal study (COVIDENCE UK)*, Thorax, doi:10.1136/thoraxjnl-2021-217487.
183. **Hosseini** et al., *Effects of Vitamin D Supplementation on COVID-19 Related Outcomes: A Systematic Review and Meta-Analysis*, Nutrients, doi:10.3390/nu14102134.
184. **Hosseini (B)** et al., *Comparing Serum Levels of Vitamin D and Zinc in Novel Coronavirus–Infected Patients and Healthy Individuals in Northeastern Iran, 2020*, Infectious Diseases in Clinical Practice, doi:10.1097/IPC.0000000000001051.
185. **Hosseini (C)** et al., *Prevention of COVID-19 with Oral Vitamin D supplemental Therapy in Essential healthCare Teams (PROTECT): Ancillary study of a randomised controlled trial*, Research Square, doi:10.21203/rs.3.rs-1588325/v1.

186. **Huang** et al., *Effect of vitamin D status on adult COVID-19 pneumonia induced by Delta variant: A longitudinal, real-world cohort study*, *Frontiers in Medicine*, doi:10.3389/fmed.2023.1121256.
187. **Hunt** et al., *Medications Associated with Lower Mortality in a SARS-CoV-2 Positive Cohort of 26,508 Veterans*, *Journal of General Internal Medicine*, doi:10.1007/s11606-022-07701-3.
188. **Hurst** et al., *Vitamin D insufficiency in COVID-19 and influenza A, and critical illness survivors: a cross-sectional study*, *BMJ Open*, doi:10.1136/bmjopen-2021-055435.
189. **Ikematsu** et al., *Baloxavir Marboxil for Prophylaxis against Influenza in Household Contacts*, *New England Journal of Medicine*, doi:10.1056/NEJMoa1915341.
190. **Im** et al., *Nutritional status of patients with COVID-19*, *Int. J. Infect. Dis.*, doi:10.1016/j.ijid.2020.08.018.
191. **Infante** et al., *Low Vitamin D Status at Admission as a Risk Factor for Poor Survival in Hospitalized Patients With COVID-19: An Italian Retrospective Study*, *Journal of the American College of Nutrition*, doi:10.1080/07315724.2021.1877580.
192. **Israel** et al., *Vitamin D deficiency is associated with higher risks for SARS-CoV-2 infection and COVID-19 severity: a retrospective case-control study*, *Internal and Emergency Medicine*, doi:10.1007/s11739-021-02902-w.
193. **Israel (B)** et al., *Identification of drugs associated with reduced severity of COVID-19: A case-control study in a large population*, *Epidemiology and Global Health Microbiology and Infectious Disease*, doi:10.7554/eLife.68165.
194. **Jabbar** et al., *Vitamin D Serum Levels and Its Association With COVID 19 Infection In Babylon Governorate, Iraq*, *Nat. Volatiles & Essent. Oils*, 8:4, www.nveo.org/index.php/journal/article/view/1046.
195. **Jabeen** et al., *Protective Effect of Vitamin-D Supplementation in Patients of Acute Coronary Syndrome During COVID-19 Pandemic*, *Pakistan Journal of Medical and Health Sciences*, doi:10.53350/pjmhs221631053.
196. **Jadad** et al., *Randomized Controlled Trials: Questions, Answers, and Musings, Second Edition*, doi:10.1002/9780470691922.
197. **Jain** et al., *Demographical Profile and Clinical Outcomes of Covid-19 Patients at a Tertiary Care Centre*, *Journal of Cardiovascular Disease Research*, doi:10.31838/jcdr.2023.14.05.215.
198. **Jain (B)** et al., *Analysis of vitamin D level among asymptomatic and critically ill COVID-19 patients and its correlation with inflammatory markers*, *Nature*, doi:10.1038/s41598-020-77093-z.
199. **Jalavu** et al., *An investigation of the correlation of vitamin D status and management outcomes in patients with severe COVID-19 at a South African tertiary hospital*, *IJID Regions*, doi:10.1016/j.ijregi.2023.05.007.
200. **Jayawardena** et al., *Impact of the vitamin D deficiency on COVID-19 infection and mortality in Asian countries*, *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*, doi:10.1016/j.dsx.2021.03.006.
201. **Jeffreys** et al., *Remdesivir-ivermectin combination displays synergistic interaction with improved in vitro activity against SARS-CoV-2*, *International Journal of Antimicrobial Agents*, doi:10.1016/j.ijantimicag.2022.106542.
202. **Jevalikar** et al., *Lack of association of baseline 25-hydroxyvitamin D levels with disease severity and mortality in Indian patients hospitalized for COVID-19*, *Scientific Reports*, doi:10.1038/s41598-021-85809-y.
203. **Jimenez** et al., *Mortality in Hemodialysis Patients with COVID-19, the Effect of Paricalcitol or Calcimimetics*, *Nutrients*, doi:10.3390/nu13082559.
204. **Jitobaom** et al., *Favipiravir and Ivermectin Showed in Vitro Synergistic Antiviral Activity against SARS-CoV-2*, *Research Square*, doi:10.21203/rs.3.rs-941811/v1.
205. **Jitobaom (B)** et al., *Synergistic anti-SARS-CoV-2 activity of repurposed anti-parasitic drug combinations*, *BMC Pharmacology and Toxicology*, doi:10.1186/s40360-022-00580-8.
206. **Jolliffe** et al., *Effect of a test-and-treat approach to vitamin D supplementation on risk of all cause acute respiratory tract infection and covid-19: phase 3 randomised controlled trial (CORONAVIT)*, *BMJ*, doi:10.1136/bmj-2022-071230.
207. **Jude** et al., *Vitamin D deficiency is associated with higher hospitalisation risk from COVID-19: a retrospective case-control study*, *Journal of Clinical Endocrinology & Metabolism*, doi:10.1210/clinem/dgab439.

208. **Junior** et al., *Chronic diseases, chest computed tomography, and laboratory tests as predictors of severe respiratory failure and death in elderly Brazilian patients hospitalized with COVID-19: a prospective cohort study*, BMC Geriatrics, doi:10.1186/s12877-022-02776-3.
209. **Juraj** et al., *COVID-19 pneumonia patients with 25(OH)D levels lower than 12 ng/ml are at increased risk of death*, International Journal of Infectious Diseases, doi:10.1016/j.ijid.2022.01.044.
210. **Kalichuran** et al., *Vitamin D status and COVID-19 severity*, Southern African Journal of Infectious Diseases, doi:10.4102/sajid.v37i1.359.
211. **Karahan** et al., *Impact of Serum 25(OH) Vitamin D Level on Mortality in Patients with COVID-19 in Turkey*, J. Nutr. Health Aging , doi:10.1007/s12603-020-1479-0.
212. **Karimpour-Razkenari** et al., *Evaluating the Effects of Clinical Characteristics and Therapeutic Regimens on Mortality in Hospitalized Patients with Severe COVID-19*, Journal of Pharmaceutical Care, doi:10.18502/jpc.v10i3.10790.
213. **Karita** et al., *Trajectory of viral load in a prospective population-based cohort with incident SARS-CoV-2 G614 infection*, medRxiv, doi:10.1101/2021.08.27.21262754.
214. **Karonova** et al., *Effect of Cholecalciferol Supplementation on the Clinical Features and Inflammatory Markers in Hospitalized COVID-19 Patients: A Randomized, Open-Label, Single-Center Study*, Nutrients, doi:10.3390/nu14132602.
215. **Karonova (B)** et al., *Vitamin D Status and Immune Response in Hospitalized Patients with Moderate and Severe COVID-19*, Pharmaceuticals, doi:10.3390/ph15030305.
216. **Karonova (C)** et al., *Low 25(OH)D Level Is Associated with Severe Course and Poor Prognosis in COVID-19*, Nutrients, doi:10.3390/nu13093021.
217. **Karonova (D)** et al., *Serum 25(OH)D level in patients with CoVID-19*, Infectology, doi:10.22625/2072-6732-2020-12-3-21-27.
218. **Katz** et al., *Increased risk for Covid-19 in patients with Vitamin D deficiency*, Nutrition, doi:10.1016/j.nut.2020.111106.
219. **Kaufman** et al., *SARS-CoV-2 positivity rates associated with circulating 25-hydroxyvitamin D levels*, PLOS One, doi:10.1371/journal.pone.0239252.
220. **Kaur** et al., *Correlation of Vitamin D Levels with COVID-19 Severity and Outcome*, Indian Journal of Clinical Practice, 32:6, ijcp.in/Admin/CMS/PDF/7.%20ClinicalStudy_2IJCP_Nov2021.pdf.
221. **Kazemi** et al., *Comparison of the cardiovascular system, clinical condition, and laboratory results in COVID-19 patients with and without vitamin D insufficiency*, BMC Infectious Diseases, doi:10.1186/s12879-022-07438-8.
222. **Kerget** et al., *Evaluation of the relationship of serum vitamin D levels in COVID-19 patients with clinical course and prognosis*, Tuberk Toraks, doi:10.5578/tt.70027.
223. **Khalil** et al., *Evaluation of vitamin D in COVID-19 patients*, 1st Samarra International Conference for Pure and Applied Sciences (SICPS2021), doi:10.1063/5.0122108.
224. **Khan** et al., *Oral Co-Supplementation of Curcumin, Quercetin, and Vitamin D3 as an Adjuvant Therapy for Mild to Moderate Symptoms of COVID-19—Results From a Pilot Open-Label, Randomized Controlled Trial*, Frontiers in Pharmacology, doi:10.3389/fphar.2022.898062.
225. **Krishnan** et al., *Clinical comorbidities, characteristics, and outcomes of mechanically ventilated patients in the State of Michigan with SARS-CoV-2 pneumonia*, J Clin Anesth., doi:10.1016/j.jclinane.2020.110005.
226. **Kumar** et al., *Association of vitamin D status with severity of COVID-19*, Journal of Cardiovascular Disease Research, 12:6, jcdonline.org/admin/Uploads/Files/6249729ba2acf0.47779591.pdf.
227. **Kumar (B)** et al., *Combining baloxavir marboxil with standard-of-care neuraminidase inhibitor in patients hospitalised with severe influenza (FLAGSTONE): a randomised, parallel-group, double-blind, placebo-controlled, superiority trial*, The Lancet Infectious Diseases, doi:10.1016/S1473-3099(21)00469-2.
228. **Lakkireddy** et al., *Effect of Short Term High Dose Oral Vitamin D Therapy on the Inflammatory Markers in Patients with COVID 19 Disease*, Archives of Clinical and Biomedical Research, doi:10.26502/acbr.50170273.

229. **Latifi-Pupovci** et al., *Relationship of anti-SARS-CoV-2 IgG antibodies with Vitamin D and inflammatory markers in COVID-19 patients*, Scientific Reports, doi:10.1038/s41598-022-09785-7.
230. **Lau** et al., *Vitamin D Insufficiency is Prevalent in Severe COVID-19*, medRxiv, doi:10.1101/2020.04.24.20075838.
231. **Lázaro** et al., *Vitamin D deficit in type 2 diabetes patients during COVID-19 lockdown with and without supplementation*, Endocrine Abstracts, doi:10.1530/endoabs.70.EP552.
232. **Leal-Martínez** et al., *Effect of a Nutritional Support System to Increase Survival and Reduce Mortality in Patients with COVID-19 in Stage III and Comorbidities: A Blinded Randomized Controlled Clinical Trial*, International Journal of Environmental Research and Public Health, doi:10.3390/ijerph19031172.
233. **Lee** et al., *Analysis of Overall Level of Evidence Behind Infectious Diseases Society of America Practice Guidelines*, Arch Intern Med., 2011, 171:1, 18-22, doi:10.1001/archinternmed.2010.482.
234. **Levitus** et al., *The Effect of Vitamin D Supplementation on Severe COVID-19 Outcomes in Patients With Vitamin D Insufficiency*, Journal of the Endocrine Society, doi: 10.1210/jendso/bvab048.567, academic.oup.com/jes/article/5/Supplement_1/A279/6240740.
235. **Levy** et al., *Frail Older Adults with Presymptomatic SARS-CoV-2 Infection: Clinical Course and Prognosis*, Gerontology, doi:10.1159/000521412.
236. **Li** et al., *Assessment of the Association of Vitamin D Level With SARS-CoV-2 Seropositivity Among Working-Age Adults*, JAMA Network Open, doi:10.1001/jamanetworkopen.2021.11634.
237. **Li (B)** et al., *Metabolic Healthy Obesity, Vitamin D Status, and Risk of COVID-19, Aging and Disease*, doi:10.14336/AD.2020.1108.
238. **Ling** et al., *High-Dose Cholecalciferol Booster Therapy is Associated with a Reduced Risk of Mortality in Patients with COVID-19: A Cross-Sectional Multi-Centre Observational Study*, Nutrients, doi:10.3390/nu12123799.
239. **Livingston** et al., *Detectable respiratory SARS-CoV-2 RNA is associated with low vitamin D levels and high social deprivation*, Int. J. Clinical Practice, doi:10.1111/ijcp.14166.
240. **Lohia** et al., *Exploring the link between vitamin D and clinical outcomes in COVID-19*, American Journal of Physiology-Endocrinology and Metabolism, doi:10.1152/ajpendo.00517.2020.
241. **Lohia (B)** et al., *Exploring the link between vitamin D and clinical outcomes in COVID-19*, American Journal of Physiology-Endocrinology and Metabolism, doi:10.1152/ajpendo.00517.2020.
242. **López-Medina** et al., *Effect of Ivermectin on Time to Resolution of Symptoms Among Adults With Mild COVID-19: A Randomized Clinical Trial*, JAMA, doi:10.1001/jama.2021.3071.
243. **Louca** et al., *Modest effects of dietary supplements during the COVID-19 pandemic: insights from 445 850 users of the COVID-19 Symptom Study app*, BMJ Nutrition, Prevention & Health, doi:10.1136/bmjnph-2021-000250.
244. **Loucera** et al., *Real world evidence of calcifediol or vitamin D prescription and mortality rate of COVID-19 in a retrospective cohort of hospitalized Andalusian patients*, Scientific Reports, doi:10.1038/s41598-021-02701-5.
245. **Luo** et al., *Vitamin D Deficiency Is Associated with COVID-19 Incidence and Disease Severity in Chinese People*, The Journal of Nutrition, doi:10.1093/jn/nxaa332.
246. **Ma** et al., *Associations between predicted vitamin D status, vitamin D intake, and risk of SARS-CoV-2 infection and Coronavirus Disease 2019 severity*, The American Journal of Clinical Nutrition, doi:10.1093/ajcn/nqab389.
247. **Ma (B)** et al., *Habitual use of vitamin D supplements and risk of coronavirus disease 2019 (COVID-19) infection: a prospective study in UK Biobank*, The American Journal of Clinical Nutrition, doi:10.1093/ajcn/nqaa381.
248. **Macaskill** et al., *A comparison of methods to detect publication bias in meta-analysis*, Statistics in Medicine, doi:10.1002/sim.698.
249. **Macaya** et al., *Interaction between age and vitamin D deficiency in severe COVID-19 infection*, Nutr. Hosp., doi:10.20960/nh.03193.

250. **Maghbooli** et al., *Treatment with 25-hydroxyvitamin D3 (calcifediol) is associated with a reduction in the blood neutrophil-to-lymphocyte ratio marker of disease severity in patients hospitalized with COVID-19: a pilot, multicenter, randomized, placebo-controlled double blind clinical trial*, Endocrine Practice, doi:10.1016/j.eprac.2021.09.016.
251. **Maghbooli (B)** et al., *Vitamin D sufficiency, a serum 25-hydroxyvitamin D at least 30 ng/mL reduced risk for adverse clinical outcomes in patients with COVID-19 infection*, PLOS One, doi:10.1371/journal.pone.0239799.
252. **Mahmood** et al., *Coronavirus in HIP Fractures CHIP 2: Is Vitamin D Deficiency Associated with Increased Mortality from COVID-19 Infections in A Hip Fracture Population?*, European Journal of Medical and Health Sciences, doi:10.24018/ejmed.2021.3.6.1159.
253. **Manojlovic** et al., *Association between vitamin D hypovitaminosis and severe forms of COVID-19*, European Review for Medical and Pharmacological Sciences, doi:10.26355/eurrev_202306_32651.
254. **Mansour** et al., *Association of serum zinc level and clinical outcome in Egyptian COVID-19 patients*, The Egyptian Journal of Internal Medicine, doi:10.1186/s43162-022-00159-z.
255. **Mansouri** et al., *The impact of calcitriol and estradiol on the SARS-CoV-2 biological activity: a molecular modeling approach*, Scientific Reports, doi:10.1038/s41598-022-04778-y.
256. **Mardani** et al., *Association of vitamin D with the modulation of the disease severity in COVID-19*, Virus Research, doi:10.1016/j.virusres.2020.198148.
257. **Mariani** et al., *High-dose vitamin D versus placebo to prevent complications in COVID-19 patients: Multicentre randomized controlled clinical trial*, PLOS ONE, doi:10.1371/journal.pone.0267918.
258. **Marik** et al., *Does vitamin D status impact mortality from SARS-CoV-2 infection?*, Med Drug Discov., doi:10.1016/j.medidd.2020.100041.
259. **Martens** et al., *Vitamin D's Effect on Immune Function*, Nutrients, doi:10.3390/nu12051248.
260. **Martínez-Rodríguez** et al., *Evaluation of the usefulness of vitamin D as a predictor of mortality in patients with COVID-19*, Gaceta Médica de México, doi:10.24875/GMM.M22000637.
261. **Matin** et al., *The sufficient vitamin D and albumin level have a protective effect on COVID-19 infection*, Archives of Microbiology, doi:10.1007/s00203-021-02482-5.
262. **Mayurathan** et al., *Association of vitamin D levels with severity and outcome of COVID-19 infection among inward patients at a tertiary care unit in Sri Lanka*, Asian Journal of Internal Medicine, 2:2, ajim.sljol.info/articles/10.4038/ajim.v2i2.84.
263. **Mazziotti** et al., *Vitamin D deficiency, secondary hyperparathyroidism and respiratory insufficiency in hospitalized patients with COVID-19*, J Endocrinol. Invest., doi:10.1007/s40618-021-01535-2.
264. **McLean** et al., *Impact of Late Oseltamivir Treatment on Influenza Symptoms in the Outpatient Setting: Results of a Randomized Trial*, Open Forum Infect. Dis. September 2015, 2:3, doi:10.1093/ofid/ofv100.
265. **medicospelavidacovid19.com.br**,
medicospelavidacovid19.com.br/editoriais/folha-de-s-paulo-revela-numeros-de-david-uip-veja-a-comparacao-com-medicos-que-fazem-tratamento-precoce/.
266. **Meeus**, G., *Online Comment*, twitter.com/gertmeeus_MD/status/1386636373889781761.
267. **Meltzer** et al., *Association of Vitamin D Levels, Race/Ethnicity, and Clinical Characteristics With COVID-19 Test Results*, JAMA Netw Open., doi:10.1001/jamanetworkopen.2021.4117.
268. **Meltzer (B)** et al., *Association of Vitamin D Status and Other Clinical Characteristics With COVID-19 Test Results*, JAMA network open, 3:9, doi:10.1001/jamanetworkopen.2020.19722.
269. **Meltzer (C)** et al., *Association of Vitamin D Levels, Race/Ethnicity, and Clinical Characteristics With COVID-19 Test Results*, JAMA Netw Open., doi:10.1001/jamanetworkopen.2021.4117.
270. **Mendy** et al., *Factors Associated with Hospitalization and Disease Severity in a Racially and Ethnically Diverse Population of COVID-19 Patients*, medRxiv, doi:10.1101/2020.06.25.20137323.
271. **Meneguesso**, A., *Médica defende tratamento precoce da Covid-19*, www.youtube.com/watch?v=X5FCrlm_19U.

272. **Meng** et al., *The role of vitamin D in the prevention and treatment of SARS-CoV-2 infection: A meta-analysis of randomized controlled trials*, Clinical Nutrition, doi:10.1016/j.clnu.2023.09.008.
273. **Merzon** et al., *Low plasma 25(OH) vitamin D level is associated with increased risk of COVID-19 infection: an Israeli population-based study*, The FEBS Journal, doi:doi.org/10.1111/febs.15495.
274. **Mingiano** et al., *Vitamin D Deficiency in COVID-19 Patients and Role of Calcifediol Supplementation*, Nutrients, doi:10.3390/nu15153392.
275. **Mishra** et al., *Vitamin D Deficiency and Comorbidities as Risk Factors of COVID-19 Infection: A Systematic Review and Meta-analysis*, Journal of Preventive Medicine and Public Health, doi:10.3961/jpmph.21.640.
276. **Mohseni** et al., *Do body mass index (BMI) and history of nutritional supplementation play a role in the severity of COVID-19? A retrospective study*, Nutrition & Food Science, doi:10.1108/NFS-11-2020-0421.
277. **Mok** et al., *Calcitriol, the active form of vitamin D, is a promising candidate for COVID-19 prophylaxis*, bioRxiv, doi:10.1101/2020.06.21.162396.
278. **Morad** et al., *Serum vitamin D level in COVID-19 patients and its correlation with disease severity*, Egyptian Rheumatology and Rehabilitation, doi:10.1186/s43166-022-00155-9.
279. **Moreno** et al., *Assessment of regression-based methods to adjust for publication bias through a comprehensive simulation study*, BMC Medical Research Methodology, doi:10.1186/1471-2288-9-2.
280. **Mostafa** et al., *Clinical and Prognostic Significance of Baseline Serum Vitamin D Levels in Hospitalized Egyptian Covid-19 Patients*, International Journal of General Medicine, doi:10.2147/IJGM.S386815.
281. **Mousa** et al., *Effect of vitamin D supplementation on inflammation and nuclear factor kappa-B activity in overweight/obese adults: a randomized placebo-controlled trial*, Scientific Reports, doi:10.1038/s41598-017-15264-1.
282. **Murai** et al., *Effect of a Single High Dose of Vitamin D3 on Hospital Length of Stay in Patients With Moderate to Severe COVID-19: A Randomized Clinical Trial*, JAMA, doi:10.1001/jama.2020.26848.
283. **Nasiri** et al., *Does vitamin D serum level affect prognosis of COVID-19 patients?*, International Journal of Infectious Diseases, doi:10.1016/j.ijid.2021.04.083.
284. **Neves** et al., *Vitamin D deficiency predicts 30-day hospital mortality of adults with COVID-19*, Clinical Nutrition ESPEN, doi:10.1016/j.clnesp.2022.05.027.
285. **Nguyen** et al., *25-hydroxyvitamin D is a predictor of COVID-19 severity of hospitalized patients*, PLOS ONE, doi:10.1371/journal.pone.0268038.
286. **Nichol** et al., *Challenging issues in randomised controlled trials*, Injury, 2010, doi: 10.1016/j.injury.2010.03.033, www.injuryjournal.com/article/S0020-1383(10)00233-0/fulltext.
287. **Nicolescu** et al., *The evaluation of vitamin D deficiency as a risk factor in the case of patients with moderate COVID-19*, Farmacia, doi:10.31925/farmacia.2022.3.17.
288. **Nikniaz** et al., *The impact of vitamin D supplementation on mortality rate and clinical outcomes of COVID-19 patients: A systematic review and meta-analysis*, Pharmaceutical Sciences, doi:10.34172/PS.2021.13.
289. **Nimavat** et al., *Vitamin D deficiency and COVID-19: A case-control study at a tertiary care hospital in India*, Annals of Medicine and Surgery, doi:10.1016/j.amsu.2021.102661.
290. **Nimer** et al., *The impact of vitamin and mineral supplements usage prior to COVID-19 infection on disease severity and hospitalization*, Bosnian Journal of Basic Medical Sciences, doi:10.17305/bjbms.2021.7009.
291. **Nogués** et al., *Calcifediol Treatment and COVID-19-Related Outcomes*, The Journal of Clinical Endocrinology & Metabolism, doi:10.1210/clinem/dgab405.
292. **Nonaka** et al., *SARS-CoV-2 variant of concern P.1 (Gamma) infection in young and middle-aged patients admitted to the intensive care units of a single hospital in Salvador, Northeast Brazil, February 2021*, International Journal of Infectious Diseases, doi:10.1016/j.ijid.2021.08.003.

293. **Ogasawara** et al., *The effect of 1-hydroxy-vitamin D treatment in hospitalized patients with COVID-19: A retrospective study*, Clinical Nutrition, doi:10.1016/j.clnu.2023.08.021.
294. **Orchard** et al., *Vitamin-D levels and intensive care unit outcomes of a cohort of critically ill COVID-19 patients*, Clin Chem Lab Med, doi:10.1515/cclm-2020-1567.
295. **Oristrell** et al., *Vitamin D supplementation and COVID-19 risk: a population-based, cohort study*, Journal of Endocrinological Investigation, doi:10.1007/s40618-021-01639-9.
296. **Oristrell (B)** et al., *Association of Calcitriol Supplementation with Reduced COVID-19 Mortality in Patients with Chronic Kidney Disease: A Population-based Study*, Biomedicines, doi:10.3390/biomedicines9050509.
297. **Ortatatli** et al., *Potential Role of Vitamin D, ACE2 and the Proteases as TMPRSS2 and Furin on SARS-CoV-2 Pathogenesis and COVID-19 Severity*, Archives of Medical Research, doi:10.1016/j.arcmed.2023.02.002.
298. **Ostrov** et al., *Highly Specific Sigma Receptor Ligands Exhibit Anti-Viral Properties in SARS-CoV-2 Infected Cells*, Pathogens, doi:10.3390/pathogens10111514.
299. **Ozturk** et al., *Is there a relationship between vitamin D levels, inflammatory parameters, and clinical severity of COVID-19 infection?*, Bratislava Medical Journal, doi:10.4149/BLL_2022_065.
300. **Panagiotou** et al., *Low serum 25-hydroxyvitamin D (25[OH]D) levels in patients hospitalised with COVID-19 are associated with greater disease severity: results of a local audit of practice*, medRxiv, doi:10.1101/2020.06.21.20136903.
301. **Pande** et al., *Vitamin D Levels and its Association with Inflammatory Markers, Severity and Outcome in Hospitalised COVID-19 Patients - An Indian Perspective*, Journal of Communicable Diseases, doi:10.24321/0019.5138.202227.
302. **Pandya** et al., *Unravelling Vitamin B12 as a potential inhibitor against SARS-CoV-2: A computational approach*, Informatics in Medicine Unlocked, doi:10.1016/j.imu.2022.100951.
303. **Papadimitriou** et al., *Association between population vitamin D status and SARS-CoV-2 related serious-critical illness and deaths: An ecological integrative approach*, World J. Virology, doi:10.5501/wjv.v10.i3.1111.
304. **Parant** et al., *Vitamin D and COVID-19 Severity in Hospitalized Older Patients: Potential Benefit of Prehospital Vitamin D Supplementation*, Nutrients, doi:10.3390/nu14081641.
305. **Parra-Ortega** et al., *25-Hydroxyvitamin D level is associated with mortality in patients with critical COVID-19: a prospective observational study in Mexico City*, Nutrition Research and Practice, doi:10.4162/nrp.2021.15.S1.S32.
306. **Peacock** et al., *The SARS-CoV-2 variant, Omicron, shows rapid replication in human primary nasal epithelial cultures and efficiently uses the endosomal route of entry*, bioRxiv, doi:10.1101/2021.12.31.474653.
307. **Pecina** et al., *Vitamin D Status and Severe COVID-19 Disease Outcomes in Hospitalized Patients*, Journal of Primary Care & Community Health , doi:10.1177/21501327211041206.
308. **Pepkowitz** et al., *Vitamin D Deficiency is Associated with Increased COVID-19 Severity: Prospective Screening of At-Risk Groups is Medically Indicated*, Research Square, doi:10.21203/rs.3.rs-83262/v1.
309. **Peters**, J., *Comparison of Two Methods to Detect Publication Bias in Meta-analysis*, JAMA, doi:10.1001/jama.295.6.676.
310. **Petkovich** et al., *Modified-release oral calcifediol corrects vitamin D insufficiency with minimal CYP24A1 upregulation*, The Journal of Steroid Biochemistry and Molecular Biology, doi:10.1016/j.jsbmb.2014.11.022.
311. **Pickard** et al., *Discovery of re-purposed drugs that slow SARS-CoV-2 replication in human cells*, PLOS Pathogens, doi:10.1371/journal.ppat.1009840.
312. **Pimental** et al., *Low vitamin D levels and increased neutrophil in patients admitted at ICU with COVID-19*, Clinical Nutrition ESPEN, doi:10.1016/j.clnesp.2021.05.021.
313. **Pop-Kostova** et al., *Vitamin D status in patients with COVID-19 – sex differences associated with severity of the disease*, Medical Journal MEDICUS, 28:1, eprints.ugd.edu.mk/31736/.
314. **Protas** et al., *Plasma 25-Hydroxyvitamin D Level and VDR Gene Single Nucleotide Polymorphism rs2228570 Influence on COVID-19 Susceptibility among the Kazakh Ethnic Group—A Pilot Study*, Nutrients, doi:10.3390/nu15071781.

315. **Putra** et al., *Vitamin D Levels among Hospitalized and Non-Hospitalized COVID-19 Patients in Dr. M. Djamil General Hospital Padang*, European Journal of Medical and Health Sciences, doi:10.24018/ejmed.2021.3.6.1131.
316. **Qayyum** et al., *Vitamin D and lumisterol novel metabolites can inhibit SARS-CoV-2 replication machinery enzymes*, Endocrinology and Metabolism, doi:10.1152/ajpendo.00174.2021.
317. **Qu** et al., *Decreased serum vitamin D level as a prognostic marker in patients with COVID-19*, arXiv, doi:10.48550/arXiv.2301.02660.
318. **Quraishi** et al., *Association Between Preoperative 25-Hydroxyvitamin D Level and Hospital-Acquired Infections Following Roux-en-Y Gastric Bypass Surgery*, JAMA Surgery, doi:10.1001/jamasurg.2013.3176.
319. **Rachman** et al., *Impact of vitamin D deficiency in relation to the clinical outcomes of hospitalized COVID-19 patients*, F1000Research, doi:10.12688/f1000research.132214.1.
320. **Radujkovic** et al., *Vitamin D Deficiency and Outcome of COVID-19 Patients*, Nutrients 2020, 12:9, 2757, doi:10.3390/nu12092757.
321. **Raisi-Estabragh** et al., *Greater risk of severe COVID-19 in Black, Asian and Minority Ethnic populations is not explained by cardiometabolic, socioeconomic or behavioural factors, or by 25(OH)-vitamin D status: study of 1326 cases from the UK Biobank*, J. Public Health, doi:10.1093/pubmed/fdaa095.
322. **Ramirez-Sandoval** et al., *Very Low Vitamin D Levels are a Strong Independent Predictor of Mortality in Hospitalized Patients with Severe COVID-19*, Archives of Medical Research, doi:10.1016/j.arcmed.2021.09.006.
323. **Ramos** et al., *Vitamin D, Zinc and Iron in Adult Patients with Covid-19 and Their Action in the Immune Response as Biomarkers*, Global Journal of Health Science, doi:10.5539/gjhs.v14n1p1.
324. **Ranjbar** et al., *Serum level of Vitamin D is associated with COVID-19 mortality rate in hospitalized patients*, Journal of Research in Medical Sciences, doi:10.4103/jrms.JRMS_1151_20.
325. **Rastogi** et al., *Short term, high-dose vitamin D supplementation for COVID-19 disease: a randomised, placebo-controlled, study (SHADE study)*, Postgraduate Medical Journal, doi:10.1136/postgradmedj-2020-139065.
326. **Rathod** et al., *Association of vitamin D with the severity of disease and mortality in COVID-19: Prospective study in central India*, Annals of African Medicine, doi:10.4103/aam.aam_21_22.
327. **Regalia** et al., *Vitamin D Status and SARS-CoV-2 Infection in a Cohort of Kidney Transplanted Patients*, Nutrients, doi:10.3390/nu14020317.
328. **Reis** et al., *Influence of vitamin D status on hospital length of stay and prognosis in hospitalized patients with moderate to severe COVID-19: a multicenter prospective cohort study*, The American Journal of Clinical Nutrition, doi:10.1093/ajcn/nqab151.
329. **Reyes Pérez** et al., *Deficiency of vitamin D is a risk factor of mortality in patients with COVID-19*, Revista de Sanidad Militar, doi:10.35366/93773.
330. **Rhodes** et al., *COVID-19 mortality increases with northerly latitude after adjustment for age suggesting a link with ultraviolet and vitamin D*, BMJ Nutr. Prev. Health, doi:10.1136/bmjnp-2020-000110.
331. **Ribeiro** et al., *Previous vitamin D status and total cholesterol are associated with SARS-CoV-2 infection*, Clinica Chimica Acta, doi:10.1016/j.cca.2021.08.003.
332. **Ricci** et al., *Circulating Vitamin D levels status and clinical prognostic indices in COVID-19 patients*, Respiratory Research, doi:10.1186/s12931-021-01666-3.
333. **Ritsinger** et al., *History of heart failure and chronic kidney disease and risk of all-cause death after COVID-19 during the first three waves of the pandemic in comparison with influenza outbreaks in Sweden: a registry-based, retrospective, case-control study*, BMJ Open, doi:10.1136/bmjopen-2022-069037.
334. **Rodríguez-Vidales** et al., *Severe COVID-19 patients have severe vitamin D deficiency in Northeast Mexico*, Nutrición Hospitalaria, doi:10.20960/nh.03731.

335. **Rothstein**, H., *Publication Bias in Meta-Analysis: Prevention, Assessment and Adjustments*, www.wiley.com/en-ae/Publication+Bias+in+Meta+Analysis:+Prevention,+Assessment+and+Adjustments-p-9780470870143.
336. **Rücker** et al., *Arcsine test for publication bias in meta-analyses with binary outcomes*, *Statistics in Medicine*, doi:10.1002/sim.2971.
337. **Rybakovsky** et al., *Calcitriol modifies tight junctions, improves barrier function, and reduces TNF- α -induced barrier leak in the human lung-derived epithelial cell culture model, 16HBE 14o-*, *Physiological Reports*, doi:10.14814/phy2.15592.
338. **Saeed** et al., *Vitamin D Deficiency and Clinical Outcomes in Patients with COVID-19*, *University of Thi-Qar Journal of Medicine*, 25:1, www.jmed.utq.edu.iq/index.php/main/article/view/380.
339. **Saeed (B)** et al., *Cholecalciferol level and its impact on COVID-19 patients*, *The Egyptian Journal of Internal Medicine*, doi:10.1186/s43162-022-00116-w.
340. **Saheb Sharif-Askari** et al., *Vitamin D modulates systemic inflammation in patients with severe COVID-19*, *Life Sciences*, doi:10.1016/j.lfs.2022.120909.
341. **Said** et al., *The effect of Nigella sativa and vitamin D3 supplementation on the clinical outcome in COVID-19 patients: A randomized controlled clinical trial*, *Frontiers in Pharmacology*, doi:10.3389/fphar.2022.1011522.
342. **Sainz-Amo** et al., *COVID-19 in Parkinson's disease: what holds the key?*, *Journal of Neurology*, doi:10.1007/s00415-020-10272-0.
343. **Sanamandra** et al., *Correlation between Serum Vitamin D3 levels and severity of COVID-19, experience from a COVID-19-dedicated tertiary care hospital from Western India*, *Indian Journal of Endocrinology and Metabolism*, doi:10.4103/ijem.ijem_383_22.
344. **Sánchez-Zuno**, J., *Vitamin D Levels in COVID-19 Outpatients from Western Mexico: Clinical Correlation and Effect of Its Supplementation* *Clinical Medicine*, doi:10.3390/jcm10112378.
345. **Sánchez-Zuno (B)**, J., *Vitamin D Levels in COVID-19 Outpatients from Western Mexico: Clinical Correlation and Effect of Its Supplementation* *Clinical Medicine*, doi:10.3390/jcm10112378.
346. **Sanson** et al., *A combined role for low vitamin D and low albumin circulating levels as strong predictors of worse outcome in COVID-19 patients*, *Irish Journal of Medical Science (1971 -)*, doi:10.1007/s11845-022-02952-9.
347. **Saponaro** et al., *Is There a Crucial Link Between Vitamin D Status and Inflammatory Response in Patients With COVID-19?*, *Frontiers in Immunology*, doi:10.3389/fimmu.2021.745713.
348. **Savitri** et al., *Comparison between Vitamin D Level of Asymptomatic Confirmed Covid-19 Patients with Symptomatic Confirmed Covid-19 Patients in Makassar*, *Annals of the Romanian Society for Cell Biology*, 25:6, www.annalsofscb.ro/index.php/journal/article/view/9130.
349. **Schmidt** et al., *Identification of Clinical Response Predictors of Tocilizumab Treatment in Patients with Severe COVID-19 Based on Single-Center Experience*, *Journal of Clinical Medicine*, doi:10.3390/jcm12062429.
350. **Schmitt** et al., *Oxidative stress status and vitamin D levels of asymptomatic to mild symptomatic COVID-19 infections during the third trimester of pregnancy: A retrospective study in Metz, France*, *Journal of Medical Virology*, doi:10.1002/jmv.27606.
351. **Seal** et al., *Association of Vitamin D Status and COVID-19-Related Hospitalization and Mortality*, *Journal of General Internal Medicine*, doi:10.1007/s11606-021-07170-0.
352. **Seely** et al., *Dietary supplements to reduce symptom severity and duration in people with SARS-CoV-2: a double-blind randomised controlled trial*, *BMJ Open*, doi:10.1136/bmjopen-2023-073761.
353. **Seely (B)** et al., *Dietary supplements to reduce symptom severity and duration in people with SARS-CoV-2: a double-blind randomised controlled trial*, *BMJ Open*, doi:10.1136/bmjopen-2023-073761.
354. **Şengül** et al., *Serum Vitamin D Concentrations and Covid-19 In Pregnant Women, Does Vitamin D Supplementation Impact Results? A Comprehensive Study*, *Cukurova Anestesi ve Cerrahi Bilimler Dergisi*, doi:10.36516/jocass.1185181.
355. **Seven** et al., *Correlation between 25-hydroxy vitamin D levels and COVID-19 severity in pregnant women: a cross-sectional study*, *The Journal of Maternal-Fetal & Neonatal Medicine*, doi:10.1080/14767058.2021.2005564.

356. **Shah** et al., *Does vitamin D supplementation reduce COVID-19 severity? - a systematic review*, QJM: An International Journal of Medicine, doi:10.1093/qjmed/hcac040.
357. **Shahid** et al., *The effects of vitamin D therapy on outcomes for hispanic patients hospitalized for COVID-19*, Abstracts from the 2022 Annual Meeting of the Society of General Internal Medicine, Journal of General Internal Medicine, doi:10.1007/s11606-022-07653-8.
358. **Shamsi** et al., *Survival and Mortality in Hospitalized Children with COVID-19: A Referral Center Experience in Yazd, Iran*, Canadian Journal of Infectious Diseases and Medical Microbiology, doi:10.1155/2023/5205188.
359. **Shannak** et al., *Evaluation of the level of vitamin D3 in the blood serum of patients infected with COVID-19 in Al-Amiriya city*, Technium BioChemMed, doi:10.47577/biochemmed.v3i2.7179.
360. **Sharif** et al., *Impact of Zinc, Vitamins C and D on Disease Prognosis among Patients with COVID-19 in Bangladesh: A Cross-Sectional Study*, Nutrients, doi:10.3390/nu14235029.
361. **Shehab** et al., *Immune-boosting effect of natural remedies and supplements on progress of, and recovery from COVID-19 infection*, Tropical Journal of Pharmaceutical Research, doi:10.4314/tjpr.v21i2.13.
362. **Silva** et al., *Does serum 25-hydroxyvitamin D decrease during acute-phase response? A systematic review*, Nutrition Research, doi:10.1016/j.nutres.2014.12.008.
363. **Sinaci** et al., *Impact of vitamin D on the course of COVID-19 during pregnancy: A case control study*, The Journal of Steroid Biochemistry and Molecular Biology, doi:10.1016/j.jsbmb.2021.105964.
364. **Singh** et al., *Single, High Dose Vitamin D Supplementation in Vitamin D Deficient Severe COVID-19: Randomized, Double-Blind, Placebocontrol Study (Shade-S)*, Abstracts Criticare - IJCCM2022, Indian J. Crit. Care Med., doi:10.5005/ijccm-26-S1-S1.
365. **Sinnberg** et al., *Vitamin C Deficiency in Blood Samples of COVID-19 Patients*, Antioxidants, doi:10.3390/antiox11081580.
366. **Siuka** et al., *The effect of Vitamin D levels on the course of COVID-19 in hospitalized patients – a 1-year prospective cohort study*, F1000Research, doi:10.12688/f1000research.131730.1.
367. **Soliman** et al., *Impact of Vitamin D Therapy on the Progress COVID-19: Six Weeks Follow-Up Study of Vitamin D Deficient Elderly Diabetes Patients*, Proceedings of Singapore Healthcare, doi:10.1177/20101058211041405.
368. **Soltani-Zangbar** et al., *Serum levels of vitamin D and immune system function in patients with COVID-19 admitted to intensive care unit*, Gene Reports, doi:10.1016/j.genrep.2022.101509.
369. **Song** et al., *Vitamin D3 and its hydroxyderivatives as promising drugs against COVID-19: a computational study*, Journal of Biomolecular Structure and Dynamics, doi:10.1080/07391102.2021.1964601.
370. **Sooriyaarachchi** et al., *Impact of vitamin D deficiency on COVID-19*, Clinical Nutrition ESPEN, doi:10.1016/j.clnesp.2021.05.011.
371. **Sposito** et al., *Age differential CD13 and interferon expression in airway epithelia affect SARS-CoV-2 infection - effects of vitamin D*, Mucosal Immunology, doi:10.1016/j.mucimm.2023.08.002.
372. **Stanley** et al., *Meta-regression approximations to reduce publication selection bias*, Research Synthesis Methods, doi:10.1002/jrsm.1095.
373. **Subramanian** et al., *Vitamin D, D-binding protein, free vitamin D and COVID-19 mortality in hospitalized patients*, The American Journal of Clinical Nutrition, doi:10.1093/ajcn/nqac027.
374. **Sulli** et al., *Vitamin D and Lung Outcomes in Elderly COVID-19 Patients*, Nutrients, doi:10.3390/nu13030717.
375. **Sulli (B)** et al., *Vitamin D and Lung Outcomes in Elderly COVID-19 Patients*, Nutrients, doi:10.3390/nu13030717.
376. **Susianti** et al., *Low levels of vitamin D were associated with coagulopathy among hospitalized coronavirus disease-19 (COVID-19) patients: A single-centered study in Indonesia*, Journal of Medical Biochemistry, doi:10.5937/jomb0-30228.
377. **Sweeting** et al., *What to add to nothing? Use and avoidance of continuity corrections in meta-analysis of sparse data*, Statistics in Medicine, doi:10.1002/sim.1761.

378. **Szeto** et al., *Vitamin D Status and COVID-19 Clinical Outcomes in Hospitalized Patients*, Endocrine Research, doi:10.1080/07435800.2020.1867162.
379. **Takase** et al., *Association between 25-hydroxyvitamin D levels and COVID-19 severity*, Clinical Nutrition ESPEN, doi:10.1016/j.clnesp.2022.04.003.
380. **Tallon** et al., *Impact of diabetes status and related factors on COVID-19-associated hospitalization: A nationwide retrospective cohort study of 116,370 adults with SARS-CoV-2 infection*, Diabetes Research and Clinical Practice, doi:10.1016/j.diabres.2022.110156.
381. **Tan** et al., *Association of Vitamin D levels on the Clinical Outcomes of Patients Hospitalized for COVID-19 in a Tertiary Hospital*, Journal of the ASEAN Federation of Endocrine Societies, doi:10.15605/jafes.038.01.07.
382. **Tan (B)** et al., *Cohort study to evaluate the effect of combination Vitamin D, Magnesium and Vitamin B12 (DMB) on progression to severe outcome in older COVID-19 patients*, Nutrition, doi:10.1016/j.nut.2020.111017.
383. **Tehrani** et al., *Evaluation of vitamin D levels in COVID-19 patients referred to Labafinejad hospital in Tehran and its relationship with disease severity and mortality*, Clinical Nutrition, doi:10.1016/j.clnesp.2021.01.014.
384. **Tentolouris** et al., *The effect of vitamin D supplementation on mortality and intensive care unit admission of COVID-19 patients. A systematic review, meta-analysis and meta-regression*, Diabetes/Metabolism Research and Reviews, doi:10.1002/dmrr.3517.
385. **Thairu** et al., *A Comparison of Ivermectin and Non Ivermectin Based Regimen for COVID-19 in Abuja: Effects on Virus Clearance, Days-to-discharge and Mortality*, Journal of Pharmaceutical Research International, doi:10.9734/jpri/2022/v34i44A36328.
386. **Tomasa-Irrigui** et al., *Low Levels of Few Micronutrients May Impact COVID-19 Disease Progression: An Observational Study on the First Wave*, Metabolites, doi:10.3390/metabo11090565.
387. **Topan** et al., *25 Hydroxyvitamin D Serum Concentration and COVID-19 Severity and Outcome—A Retrospective Survey in a Romanian Hospital*, Nutrients, doi:10.3390/nu15051227.
388. **Treanor** et al., *Efficacy and Safety of the Oral Neuraminidase Inhibitor Oseltamivir in Treating Acute Influenza: A Randomized Controlled Trial*, JAMA, 2000, 283:8, 1016-1024, doi:10.1001/jama.283.8.1016.
389. **Tylicki** et al., *Predictors of Mortality in Hemodialyzed Patients after SARS-CoV-2 Infection*, Journal of Clinical Medicine, doi:10.3390/jcm11020285.
390. **Ullah** et al., *COVID-19 in patients with hepatobiliary and pancreatic diseases in East London: a single-centre cohort study*, Pancreatology, doi:10.1016/j.pan.2020.10.005.
391. **Umay** et al., *Comparison of Length of Hospital Stay and Routine Laboratory Parameters in Covid-19 Patients With and Without Serum Vitamin D Deficiency*, Journal of Contemporary Medicine, doi:10.16899/jcm.1319088.
392. **Ünsal** et al., *Retrospective analysis of vitamin D status on inflammatory markers and course of the disease in patients with COVID-19 infection*, Journal of Endocrinological Investigation, doi:10.1007/s40618-021-01566-9.
393. **Ünsal (B)** et al., *Retrospective analysis of vitamin D status on inflammatory markers and course of the disease in patients with COVID-19 infection*, Journal of Endocrinological Investigation, doi:10.1007/s40618-021-01566-9.
394. **Valecha** et al., *The Effect of Vitamin B12, Magnesium and Vitamin D in COVID-19 among Geriatric Patients*, International Journal of Pharmaceutical and Clinical Research, 14:5, impactfactor.org/PDF/IJPCR/14/IJPCR,Vol14,Issue5,Article113.pdf.
395. **van Helmond** et al., *Vitamin D3 Supplementation at 5000 IU Daily for the Prevention of Influenza-like Illness in Healthcare Workers: A Pragmatic Randomized Clinical Trial*, Nutrients, doi:10.3390/nu15010180.
396. **Vanegas-Cedillo** et al., *Serum Vitamin D Levels Are Associated With Increased COVID-19 Severity and Mortality Independent of Whole-Body and Visceral Adiposity*, medRxiv, doi:10.1101/2021.03.12.21253490.
397. **Varikasuvu** et al., *COVID-19 and Vitamin D (Co-VIVID Study): a systematic review and meta-analysis of randomized controlled trials*, Expert Review of Anti-infective Therapy, doi:10.1080/14787210.2022.2035217.

398. **Vasheghani** et al., *The relationship between serum 25-hydroxyvitamin D levels and the severity of COVID-19 disease and its mortality*, Scientific Reports, doi:10.1038/s41598-021-97017-9.
399. **Vasheghani (B)** et al., *The relationship between serum 25-hydroxyvitamin D levels and the severity of COVID-19 disease and its mortality*, Scientific Reports, doi:10.1038/s41598-021-97017-9.
400. **Vásquez-Procopio** et al., *Association between 25-OH Vitamin D Deficiency and COVID-19 Severity in Pregnant Women*, International Journal of Molecular Sciences, doi:10.3390/ijms232315188.
401. **Vassiliou** et al., *Vitamin D deficiency correlates with a reduced number of natural killer cells in intensive care unit (ICU) and non-ICU patients with COVID-19 pneumonia*, Hellenic Journal of Cardiology, doi:10.1016/j.hjc.2020.11.011.
402. **Vassiliou (B)** et al., *Low 25-Hydroxyvitamin D Levels on Admission to the Intensive Care Unit May Predispose COVID-19 Pneumonia Patients to a Higher 28-Day Mortality Risk: A Pilot Study on a Greek ICU Cohort*, Nutrients, doi:10.3390/nu12123773.
403. **Villasis-Keever** et al., *Efficacy and Safety of Vitamin D Supplementation to Prevent COVID-19 in Frontline Healthcare Workers. A Randomized Clinical Trial*, Archives of Medical Research, doi:10.1016/j.arcmed.2022.04.003.
404. **Voelkle** et al., *Prevalence of Micronutrient Deficiencies in Patients Hospitalized with COVID-19: An Observational Cohort Study*, Nutrients, doi:10.3390/nu14091862.
405. **Walk** et al., *Vitamin D - contrary to vitamin K - does not associate with clinical outcome in hospitalized COVID-19 patients*, medRxiv, doi:10.1101/2020.11.07.20227512.
406. **Walrand, S.**, *Autumn COVID-19 surge dates in Europe correlated to latitudes, not to temperature-humidity, pointing to vitamin D as contributing factor*, Nature, doi:10.1038/s41598-021-81419-w.
407. **Wang** et al., *Influence of a High Vitamin D2 Dose on the Prevention and Improvement of Symptomatic COVID-19 in Health Care Workers: A Multicenter Randomized Clinical Trial*, Elsevier BV, doi:10.2139/ssrn.4401710.
408. **Wani** et al., *Impact of Age and Clinico-Biochemical Parameters on Clinical severity of SARS-CoV-2 Infection*, Intervirology, doi:10.1159/000530906.
409. **Willett** et al., *The hyper-transmissible SARS-CoV-2 Omicron variant exhibits significant antigenic change, vaccine escape and a switch in cell entry mechanism*, medRxiv, doi:10.1101/2022.01.03.21268111.
410. **Williams, T.**, *Not All Ivermectin Is Created Equal: Comparing The Quality of 11 Different Ivermectin Sources*, Do Your Own Research, doyourownresearch.substack.com/p/not-all-ivermectin-is-created-equal.
411. **Xie** et al., *Micronutrient perspective on COVID-19: Umbrella review and reanalysis of meta-analyses*, Critical Reviews in Food Science and Nutrition, doi:10.1080/10408398.2023.2174948.
412. **Xu** et al., *A study of impurities in the repurposed COVID-19 drug hydroxychloroquine sulfate by UHPLC-Q/TOF-MS and LC-SPE-NMR*, Rapid Communications in Mass Spectrometry, doi:10.1002/rcm.9358.
413. **Yadav** et al., *Association of Vitamin D Status with COVID-19 Infection and Mortality in the Asia Pacific region: A Cross-Sectional Study*, Indian Journal of Clinical Biochemistry, doi:10.1007/s12291-020-00950-1.
414. **Ye** et al., *Does Serum Vitamin D Level Affect COVID-19 Infection and Its Severity? A Case-Control Study*, Journal of the American College of Nutrition, doi:10.1080/07315724.2020.182600.
415. **Yildiz** et al., *The prognostic significance of vitamin D deficiency in patients with COVID-19 pneumonia*, Bratislava Medical Journal, doi:10.4149/BLL_2021_119.
416. **Yilmaz** et al., *Is vitamin D deficiency a risk factor for COVID-19 in children?*, Pediatric Pulmonology, doi:10.1002/ppul.25106.
417. **Zafar** et al., *Vitamin D levels and mortality with SARS-COV-2 infection: a retrospective two-centre cohort study*, Postgraduate Medical Journal, doi:10.1136/postgradmedj-2021-140564.
418. **Zangeneh** et al., *Survival analysis based on body mass index in patients with Covid-19 admitted to the intensive care unit of Amir Al-Momenin Hospital in Arak – 2021*, Obesity Medicine, doi:10.1016/j.obmed.2022.100420.

419. **Zavascki** et al., *Advanced ventilatory support and mortality in hospitalized patients with COVID-19 caused by Gamma (P.1) variant of concern compared to other lineages: cohort study at a reference center in Brazil*, Research Square, doi:10.21203/rs.3.rs-910467/v1.
420. **Zeidan** et al., *Vitamin D deficiency and vitamin D receptor FokI polymorphism as risk factors for COVID-19*, Pediatric Research, doi:10.1038/s41390-022-02275-6.
421. **Zelzer** et al., *Vitamin D Metabolites and Clinical Outcome in Hospitalized COVID-19 Patients*, Nutrients, doi:10.3390/nu13072129.
422. **Zeraatkar** et al., *Consistency of covid-19 trial preprints with published reports and impact for decision making: retrospective review*, BMJ Medicine, doi:10.1136/bmjmed-2022-0003091.
423. **Zhang** et al., *What's the relative risk? A method of correcting the odds ratio in cohort studies of common outcomes*, JAMA, 80:19, 1690, doi:10.1001/jama.280.19.1690.
424. **Zidrou** et al., *The Relationship Between Vitamin D Status and the Clinical Severity of COVID-19 Infection: A Retrospective Single-Center Analysis*, Cureus, doi:10.7759/cureus.22385.
425. **Zimmerman** et al., *Melatonin and the Optics of the Human Body*, Melatonin Research, doi:10.32794/mr11250016.
426. **Zurita-Cruz** et al., *Efficacy and safety of vitamin D supplementation in hospitalized COVID-19 pediatric patients: A randomized controlled trial*, Frontiers in Pediatrics, doi:10.3389/fped.2022.943529.