

Exercise reduces COVID-19 risk: real-time meta analysis of 68 studies

@CovidAnalysis, July 2025, Version 43
<https://c19early.org/exmeta.html>

Abstract

Significantly lower risk is seen for mortality, ventilation, ICU admission, hospitalization, progression, recovery, and cases. 52 studies from 52 independent teams in 24 countries show significant benefit.

Meta analysis using the most serious outcome reported shows 39% [33-44%] lower risk. Results are similar for higher quality and peer-reviewed studies.

Results are very robust — in exclusion sensitivity analysis 56 of 68 studies must be excluded to avoid finding statistically significant efficacy in pooled analysis.

Results are consistent with the overall risk of all cause mortality based on cardiorespiratory fitness — *Laukkanen* show RR 0.55 [0.50-0.61] for the top vs. bottom tertiles.

Most studies analyze activity levels before infection, comparing regular/moderate exercise and lower/no exercise. Risk may increase with more extreme activity levels. Exercise may also be beneficial after infection. One study shows lower COVID-19 mortality with exercise during hospitalization².


No treatment is 100% effective. Protocols combine safe and effective options with individual risk/benefit analysis and monitoring. All data and sources to reproduce this analysis are in the appendix.

6 other meta analyses show significant improvements with exercise for mortality³⁻⁷, ICU admission³, hospitalization^{3,4,7,8}, severity⁴⁻⁶, and cases⁴.


Serious Outcome Risk



Exercise for COVID-19



Evolution of COVID-19 clinical evidence



EXERCISE FOR COVID-19 — HIGHLIGHTS

Exercise reduces risk with very high confidence for mortality, ICU admission, hospitalization, recovery, cases, and in pooled analysis, and low confidence for ventilation and progression.

9th treatment shown effective in October 2020, now with $p < 0.00000000001$ from 68 studies.

Real-time updates and corrections with a consistent protocol for 172 treatments. Outcome specific analysis and combined evidence from all studies including treatment delay, a primary confounding factor.

		Improvement, RR [CI]	Treatment	Control		
Fernandez	47%	0.53 [0.29-0.90]	death	16/201	62/238	
Late treatment	47%	0.53 [0.29-0.90]	16/201	62/238		47% lower risk
Tau ² = 0.00, I ² = 0.0%, p = 0.014						
		Improvement, RR [CI]	Treatment	Control		
Hamer	28%	0.72 [0.61-0.87]	hosp.	n/a	n/a	
Brawner	74%	0.26 [0.11-0.58]	hosp.	n/a	n/a	
Tret'yakov	98%	0.02 [0.00-0.27]	severe case	0/27	53/266	
Gao	-105%	2.05 [1.39-3.02]	cases	case control		
Ho	35%	0.65 [0.48-0.89]	hosp.	213/123,588	59/14,887	
Halabchi	89%	0.11 [0.01-1.46]	death	0/249	79/4,445	
Zhang	26%	0.74 [0.48-1.14]	death	n/a	n/a	
Li	81%	0.19 [0.05-0.74]	severe case	n/a	n/a	
Tavakoli	69%	0.31 [0.10-1.02]	severe case	3/64	19/124	
Yates	45%	0.55 [0.38-0.79]	death	72/163,912	62/30,119	
Holt	17%	0.83 [0.63-1.09]	cases	15,227 (all patients)		
Cho	53%	0.47 [0.26-0.87]	death	case control		
Sallis	59%	0.41 [0.22-0.76]	death	11/3,118	170/6,984	
Christensen	63%	0.37 [0.16-0.85]	death	543 (n)	529 (n)	
Lobelo	20%	0.80 [0.66-0.97]	hosp.	2,121 (n)	1,648 (n)	
Hegazy	46%	0.54 [0.26-1.15]	m/s case	24/82	7/13	
Latorre-Román	76%	0.24 [0.05-1.04]	hosp.	n/a	n/a	
Marcus	42%	0.58 [0.48-0.71]	symp. case	240/10,627	134/3,708	
Yuan	91%	0.09 [0.01-1.65]	death	0/61	6/103	
Brandenburg	-6%	1.06 [0.23-4.79]	hosp.	102 (n)	39 (n)	
Bielik	30%	0.70 [0.40-1.21]	mod. case	775 (n)	365 (n)	
af Geijerstam	50%	0.50 [0.31-0.81]	death	318,902 (n)	336,271 (n)	
Lee	74%	0.26 [0.07-0.99]	death	2/11,072	32/41,293	
Maltagliati	52%	0.48 [0.25-0.87]	hosp.	n/a	n/a	
Baynouna AlKetbi	98%	0.01 [0.00-0.44]	death	n/a	n/a	
Ahmadi	30%	0.70 [0.54-0.89]	death	160/232,613	112/95,221	
Nguyen	20%	0.80 [0.72-0.88]	symp. case	904/2,836	483/1,111	
Lin	47%	0.53 [0.12-2.33]	cases	n/a	n/a	
de Souza	73%	0.27 [0.07-1.06]	ventilation	3/611	6/327	
Mohsin	19%	0.81 [0.66-0.99]	severe case	86/258	224/544	
Eklblom-Bak	48%	0.52 [0.30-0.92]	severe case	n/a	n/a	
Lengelé	74%	0.26 [0.07-0.80]	cases	23/229	4/12	
Saadeh	9%	0.91 [0.56-1.43]	symp. case	362 (n)	225 (n)	
Hamrouni	29%	0.71 [0.55-0.92]	death	138/106,006	109/47,827	
Huang	47%	0.53 [0.23-1.22]	severe case	7/74	16/90	
Hamdan	16%	0.84 [0.49-1.39]	hosp.	22/128	37/172	
Steenkamp	42%	0.58 [0.50-0.68]	death	29,469 (n)	13,366 (n)	
Gilley	-42%	1.42 [0.60-3.35]	cases	172/1,917	5/79	
Almansour	6%	0.94 [0.40-1.47]	cases	35/71	38/71	
Beydoun	43%	0.57 [0.32-1.01]	cases	1,710 (n)	448 (n)	
Salgado-Aranda	83%	0.17 [0.05-0.56]	death	4/223	41/297	
Paul	38%	0.62 [0.30-1.20]	PASC	1,811 (all patients)		
Kontopoulou	66%	0.34 [0.24-0.47]	recov. time	42 (n)	24 (n)	
Malisoux	63%	0.37 [0.14-0.98]	progression	115 (n)	108 (n)	
Antunes	80%	0.20 [0.03-1.41]	ICU	1/14	9/25	
Tsuzuki	56%	0.44 [0.36-0.52]	severe case	3,340 (n)	1,528 (n)	
Reis	41%	0.59 [0.26-1.26]	hosp.	9/241	29/305	




Figure 1. A. Random effects meta-analysis. This plot shows pooled effects, see the specific outcome analyses for individual outcomes. Analysis validating pooled outcomes for COVID-19 can be found below. Effect extraction is pre-specified, using the most serious outcome reported. For details see the appendix. **B. Timeline of results in exercise studies.** The marked dates indicate the time when efficacy was known with a statistically significant improvement of $\geq 10\%$ from ≥ 3 studies for pooled outcomes and one or more specific outcome. Efficacy based on specific outcomes was delayed by 0.8 months, compared to using pooled outcomes.

Introduction

Exercise

Exercise can improve immune system function, reduce chronic inflammation, improve cardiovascular health, improve comorbidities, enhance lung function, reduce stress, and increase nitric oxide. Prolonged high-intensity workouts may temporarily suppress the immune system. Insufficient physical activity is a risk factor for many diseases, is common around the world, has increasing prevalence over time, and has over two times greater prevalence in high-income countries⁹. For upper respiratory tract infections, research shows lower risk for moderate activity vs. a sedentary lifestyle, however risk may increase with more extreme activity levels¹⁰.

Other infections

Efficacy with exercise has been shown for pneumonia¹¹.

Analysis

We analyze all significant studies reporting COVID-19 outcomes as a function of physical activity levels. Search methods, inclusion criteria, effect extraction criteria (more serious outcomes have priority), all individual study data, PRISMA answers, and statistical methods are detailed in Appendix 1. We present random effects meta-analysis results for all studies, studies within each treatment stage, individual outcomes, peer-reviewed studies, and higher quality studies.

Results

Table 1 summarizes the results for all studies, for peer-reviewed studies, after exclusions, and for specific outcomes. Figure 2 plots individual results by treatment stage. Figure 3, 4, 5, 6, 7, 8, 9, 10, 11, and 12 show forest plots for random effects meta-analysis of all studies with pooled effects, mortality results, ventilation, ICU admission, hospitalization, progression, recovery, cases, peer reviewed studies, and long COVID.

	Relative Risk	Studies	Patients
All studies	0.61 [0.56-0.67] ****	68	1M
After exclusions	0.63 [0.57-0.68] ****	63	1M
Peer-reviewed	0.62 [0.57-0.67] ****	66	1M
Mortality	0.52 [0.43-0.62] ****	19	1M
Ventilation	0.54 [0.43-0.68] ****	2	40K
ICU admission	0.59 [0.53-0.65] ****	4	700K
Hospitalization	0.67 [0.60-0.75] ****	20	1M
Recovery	0.42 [0.30-0.59] ****	3	297
Cases	0.77 [0.69-0.86] ****	25	280K

Table 1. Random effects meta-analysis for all studies, for peer-reviewed studies, after exclusions, and for specific outcomes. Results show the relative risk with increased activity levels and the 95% confidence interval. * $p < 0.05$ **** $p < 0.0001$.





Figure 2. Scatter plot showing the most serious outcome in all studies, and for studies within each stage. Diamonds shows the results of random effects meta-analysis.

68 exercise COVID-19 studies

c19early.org
July 2025




Figure 3. Random effects meta-analysis for all studies. This plot shows pooled effects, see the specific outcome analyses for individual outcomes. Analysis validating pooled outcomes for COVID-19 can be found below. Effect extraction is pre-specified, using the most serious outcome reported. For details see the appendix.

19 exercise COVID-19 mortality results

c19early.org
July 2025




Figure 4. Random effects meta-analysis for mortality results.

2 exercise COVID-19 mechanical ventilation results

c19early.org
July 2025




Figure 5. Random effects meta-analysis for ventilation.

4 exercise COVID-19 ICU results

c19early.org

July 2025




Figure 6. Random effects meta-analysis for ICU admission.

20 exercise COVID-19 hospitalization results

c19early.org

July 2025




Figure 7. Random effects meta-analysis for hospitalization.

2 exercise COVID-19 progression results

c19early.org

July 2025




Figure 8. Random effects meta-analysis for progression.

3 exercise COVID-19 recovery results

c19early.org

July 2025




Figure 9. Random effects meta-analysis for recovery.

25 exercise COVID-19 case results



c19early.org
July 2025Tau² = 0.05, I² = 91.6%, p < 0.0001

Figure 10. Random effects meta-analysis for cases.

66 exercise COVID-19 peer reviewed studies

c19early.org
July 2025

Tau² = 0.07, I² = 79.6%, p < 0.0001Effect extraction pre-specified
(most serious outcome, see appendix)

Favors exercise Favors inactivity

Figure 11. Random effects meta-analysis for peer reviewed studies. Effect extraction is pre-specified, using the most serious outcome reported, see the appendix for details. Analysis validating pooled outcomes for COVID-19 can be found below. Zeraatkar *et al.* analyze 356 COVID-19 trials, finding no significant evidence that preprint results are inconsistent with peer-reviewed studies. They also show extremely long peer-review delays, with a median of 6 months to journal publication. A six month delay was equivalent to around 1.5 million deaths during the first two years of the pandemic. Authors recommend using preprint evidence, with appropriate checks for potential falsified data, which provides higher certainty much earlier. Davidson *et al.* also showed no important difference between meta analysis results of preprints and peer-reviewed publications for COVID-19, based on 37 meta analyses including 114 trials.

6 exercise COVID-19 long COVID results


c19early.org
July 2025

Figure 12. Random effects meta-analysis for long COVID. Effect extraction is pre-specified, using the most serious outcome reported, see the appendix for details. Analysis validating pooled outcomes for COVID-19 can be found below.

Exclusions

To avoid bias in the selection of studies, we analyze all non-retracted studies. Here we show the results after excluding studies with major issues likely to alter results, non-standard studies, and studies where very minimal detail is currently available. Our bias evaluation is based on analysis of each study and identifying when there is a significant chance that limitations will substantially change the outcome of the study. We believe this can be more valuable than checklist-based approaches such as Cochrane GRADE, which can be easily influenced by potential bias, may ignore or underemphasize serious issues not captured in the checklists, and may overemphasize issues unlikely to alter outcomes in specific cases (for example certain specifics of randomization with a very large effect size and well-matched baseline characteristics).

The studies excluded are as below. Figure 13 shows a forest plot for random effects meta-analysis of all studies after exclusions.

Brawner, unadjusted results with no group details.

de Souza, unadjusted results with no group details. Excluded results: mechanical ventilation.

Hegazy, unadjusted results with no group details.

Huang, unadjusted results with no group details. Excluded results: severe case.


Kontopoulou, unadjusted results with no group details.

Mohsin, unadjusted results with no group details.

Tret'yakov, unadjusted results with no group details.

Yuan, excessive unadjusted differences between groups. Excluded results: death.

63 exercise COVID-19 studies after exclusions

c19early.org
July 2025Tau² = 0.06, I² = 79.8%, p < 0.0001Effect extraction pre-specified
(most serious outcome, see appendix)

Favors exercise Favors inactivity

Figure 13. Random effects meta-analysis for all studies after exclusions. This plot shows pooled effects, see the specific outcome analyses for individual outcomes. Analysis validating pooled outcomes for COVID-19 can be found below. Effect extraction is pre-specified, using the most serious outcome reported. For details see the appendix.

Pooled Effects

Pooled effects are no longer required to show efficacy as of November 2020

This section validates the use of pooled effects for COVID-19, which enables earlier detection of efficacy, however pooled effects are no longer required for exercise as of November 2020. Efficacy is now known based on specific outcomes. Efficacy based on specific outcomes was delayed by 0.8 months compared to using pooled outcomes.

Combining studies is required

For COVID-19, delay in clinical results translates into additional death and morbidity, as well as additional economic and societal damage. Combining the results of studies reporting different outcomes is required. There may be no mortality in a trial with low-risk patients, however a reduction in severity or improved viral clearance may translate into lower mortality in a high-risk population. Different studies may report lower severity, improved recovery, and lower mortality, and the significance may be very high when combining the results. "*The studies reported different outcomes*" is not a good reason for disregarding results. Pooling the results of studies reporting different outcomes allows us to use more of the available information. Logically we should, and do, use additional information when evaluating treatments—for example dose-response and treatment delay-response relationships provide additional evidence of efficacy that is considered when reviewing the evidence for a treatment.

Specific outcome and pooled analyses

We present both specific outcome and pooled analyses. In order to combine the results of studies reporting different outcomes we use the most serious outcome reported in each study, based on the thesis that improvement in the most serious outcome provides comparable measures of efficacy for a treatment. A critical advantage of this approach is simplicity and transparency. There are many other ways to combine evidence for different outcomes, along with additional evidence such as dose-response relationships, however these increase complexity.

Ethical and practical issues limit high-risk trials

Trials with high-risk patients may be restricted due to ethics for treatments that are known or expected to be effective, and they increase difficulty for recruiting. Using less severe outcomes as a proxy for more serious outcomes allows faster and safer collection of evidence.

Validating pooled outcome analysis for COVID-19

For many COVID-19 treatments, a reduction in mortality logically follows from a reduction in hospitalization, which follows from a reduction in symptomatic cases, which follows from a reduction in PCR positivity. We can directly test this for COVID-19.

Analysis of the the association between different outcomes across studies from all 172 treatments we cover confirms the validity of pooled outcome analysis for COVID-19. Figure 14 shows that lower hospitalization is very strongly associated with lower mortality ($p < 0.000000000001$). Similarly, Figure 15 shows that improved recovery is very strongly associated with lower mortality ($p < 0.000000000001$). Considering the extremes, *Singh et al.* show an association between viral clearance and hospitalization or death, with $p = 0.003$ after excluding one large outlier from a mutagenic treatment, and based on 44 RCTs including 52,384 patients. Figure 16 shows that improved viral clearance is strongly associated with fewer serious outcomes. The association is very similar to *Singh et al.*, with higher confidence due to the larger number of studies. As with *Singh et al.*, the confidence increases when excluding the outlier treatment, from $p = 0.000000082$ to $p = 0.0000000033$.




Figure 14. Lower hospitalization is associated with lower mortality, supporting pooled outcome analysis.




Figure 15. Improved recovery is associated with lower mortality, supporting pooled outcome analysis.




Figure 14. Improved viral clearance is associated with fewer serious outcomes, supporting pooled outcome analysis.

Pooled outcomes identify efficacy 5 months faster (7 months for RCTs)

Currently, 55 of the treatments we analyze show statistically significant efficacy or harm, defined as $\geq 10\%$ decreased risk or $>0\%$ increased risk from ≥ 3 studies. 88% of these have been confirmed with one or more specific outcomes, with a mean delay of 4.9 months. When restricting to RCTs only, 57% of treatments showing statistically significant efficacy/harm with pooled effects have been confirmed with one or more specific outcomes, with a mean delay of 7.3 months. Figure 17 shows when treatments were found effective during the pandemic. Pooled outcomes often resulted in earlier detection of efficacy.

Time when COVID-19 studies showed efficacy

c19early.org
July 2025




Figure 17. The time when studies showed that treatments were effective, defined as statistically significant improvement of $\geq 10\%$ from ≥ 3 studies. Pooled results typically show efficacy earlier than specific outcome results. Results from all studies often shows efficacy much earlier than when restricting to RCTs. Results reflect conditions as used in trials to date, these depend on the population treated, treatment delay, and treatment regimen.

Limitations

Pooled analysis could hide efficacy, for example a treatment that is beneficial for late stage patients but has no effect on viral clearance may show no efficacy if most studies only examine viral clearance. In practice, it is rare for a non-antiviral treatment to report viral clearance and to not report clinical outcomes; and in practice other sources of heterogeneity such as difference in treatment delay is more likely to hide efficacy.

Summary

Analysis validates the use of pooled effects and shows significantly faster detection of efficacy on average. However, as with all meta analyses, it is important to review the different studies included. We also present individual outcome analyses, which may be more informative for specific use cases.

Discussion

Results for other infections

Efficacy with exercise has also been shown for pneumonia¹¹.

Exercise post-infection

Most studies analyze activity levels before infection, comparing regular/moderate exercise and lower/no exercise. Risk may increase with more extreme activity levels. Exercise may also be beneficial after infection. One study shows lower COVID-19 mortality with exercise during hospitalization². When appropriate and within limits, exercise may be beneficial even for later stage patients, for example a non-COVID-19 RCT with critical patients under mechanical ventilation (APACHE II 22) shows lower mortality with exercise²³.

Notes

6 other meta analyses show significant improvements with exercise for mortality³⁻⁷, ICU admission³, hospitalization^{3,4,7,8}, severity⁴⁻⁶, and cases⁴.

Reviews

Multiple reviews cover exercise for COVID-19, presenting additional background on mechanisms and related results, including²⁴⁻²⁶.

Perspective

Results compared with other treatments

SARS-CoV-2 infection and replication involves a complex interplay of 100+ host and viral proteins and other factors²⁷⁻³⁴, providing many therapeutic targets. Over 9,000 compounds have been predicted to reduce COVID-19 risk³⁵, either by directly minimizing infection or replication, by supporting immune system function, or by minimizing secondary complications. Exercise can improve immune system function, reduce chronic inflammation, improve cardiovascular health, improve comorbidities, enhance lung function, reduce stress, and increase nitric oxide. Prolonged high-intensity workouts may temporarily suppress the immune system. Figure 18 shows an overview of the results for exercise in the context of multiple COVID-19 treatments, and Figure 19 shows a plot of efficacy vs. cost for COVID-19 treatments.

Efficacy in COVID-19 studies (pooled effects)

c19early.org
July 2025




Figure 18. Scatter plot showing results within the context of multiple COVID-19 treatments. Diamonds shows the results of random effects meta-analysis. 0.6% of 9,000+ proposed treatments show efficacy³⁶.

Efficacy vs. cost for COVID-19 treatments

● Lifestyle / free
● No prescription
● Prescription required
● High-cost

c19early.org
July 2025




Figure 19. Efficacy vs. cost for COVID-19 treatments.

Conclusion

Exercise can improve immune system function, reduce chronic inflammation, improve cardiovascular health, improve comorbidities, enhance lung function, reduce stress, and increase nitric oxide. Prolonged high-intensity workouts may temporarily suppress the immune system.

More physically active people have reduced risk for COVID-19. Significantly lower risk is seen for mortality, ventilation, ICU admission, hospitalization, progression, recovery, and cases. 52 studies from 52 independent teams in 24 countries show significant benefit. Meta analysis using the most serious outcome reported shows 39% [33-44%] lower risk. Results are similar for higher quality and peer-reviewed studies. Results are very robust — in exclusion sensitivity analysis 56 of 68 studies must be excluded to avoid finding statistically significant efficacy in pooled analysis. Results are consistent with the overall risk of all cause mortality based on cardiorespiratory fitness — *Laukkanen* show RR 0.55 [0.50-0.61] for the top vs. bottom tertiles.


Most studies analyze activity levels before infection, comparing regular/moderate exercise and lower/no exercise. Risk may increase with more extreme activity levels. Exercise may also be beneficial after infection. One study shows lower COVID-19 mortality with exercise during hospitalization².

6 other meta analyses show significant improvements with exercise for mortality³⁻⁷, ICU admission³, hospitalization^{3,4,7,8}, severity⁴⁻⁶, and cases⁴.

Efficacy with exercise has also been shown for pneumonia¹¹.

Study Notes


af Geijerstam



Prospective study of 1,559,187 men in Sweden with cardiorespiratory fitness levels measured on military conscription, showing high cardiorespiratory fitness associated with lower risk of COVID-19 hospitalization, ICU admission, and death.

Ahmadi

Exercise for COVID-19 Ahmadi et al. Prophylaxis



Retrospective 468,569 adults in the UK, showing significantly lower COVID-19 mortality with physical activity.

Akbar


Exercise for COVID-19 Akbar et al. Prophylaxis



Retrospective 10,000 adults in Qatar, showing lower risk of COVID-19 cases with increased leisure time physical activity, without statistical significance. Authors do not analyze COVID-19 severity.

Almansour


Exercise for COVID-19 Almansour et al. Prophylaxis



Retrospective 142 patients in Saudi Arabia, showing no significant difference in cases with physical activity.

Antunes

Exercise for COVID-19 Antunes et al. Prophylaxis



Does physical activity reduce risk for COVID-19?

Retrospective 39 patients in Brazil (September - December 2020)

Shorter hospitalization with higher activity levels (p=0.03)


Antunes et al., Sport Sciences for Health, Jun 2022

c19early.org

Retrospective 39 hospitalized COVID-19 survivors >60 years old, showing shorter hospitalization for patients with active lifestyles before COVID-19 symptoms.

Baynouna AlKetbi

Exercise Baynouna AlKetbi et al. Prophylaxis



Does physical activity reduce risk for COVID-19?

Retrospective study in United Arab Emirates

Lower mortality with higher activity levels (p=0.049)


Baynouna AlKetbi et al., J. Epidemiol., Aug 2021

c19early.org

Retrospective 234 COVID-19 cases in the United Arab Emirates, showing lower risk of mortality with increased physical activity.

Beydoun

Exercise for COVID-19 Beydoun et al. Prophylaxis



Does physical activity reduce risk for COVID-19?

Retrospective 2,158 patients in the USA

Fewer cases with higher activity levels (not stat. sig., p=0.055)


Beydoun et al., American J. Infection ..., Mar 2022

c19early.org

Retrospective 2,830 people in the USA, showing lower risk of COVID-19 with a history of moderate/vigorous exercise.

Bielik

Exercise for COVID-19 Bielik et al. Prophylaxis



Does physical activity reduce risk for COVID-19?

Retrospective 1,140 patients in Slovakia (December - December 2020)

Fewer moderate/severe cases with higher activity levels (not stat. sig., $p=0.1$)


Bielik et al., Int. J. Environmental R., Jul 2021

c19early.org

Retrospective 1,544 participants in Slovakia, showing a lower risk of more severe COVID-19 for physically active participants, without statistical significance.

Brandenburg

Exercise for COVID-19 Brandenburg et al. Prophylaxis



Does physical activity reduce risk for COVID-19?

Retrospective 211 patients in multiple countries

Lower severe cases with higher activity levels (not stat. sig., $p=0.3$)


Brandenburg et al., J. Physical Activi., Jul 2021

c19early.org

Retrospective 263 COVID+ patients, showing lower hospitalization with higher self-reported cardiorespiratory fitness, but no significant differences for physical activity. Participants in the study were healthier and more fit than the general population.

Brawner

Exercise for COVID-19 Brawner et al. Prophylaxis



Does physical activity reduce risk for COVID-19?

Retrospective study in the USA (February - May 2020)

Lower hospitalization with higher activity levels ($p=0.0012$)


Brawner et al., Mayo Clinic Proceedings, Oct 2020

c19early.org

Retrospective 246 COVID-19 patients in the USA, showing the risk of hospitalization inversely associated with maximal exercise capacity. Adjusted results are only provided for MET as a continuous variable.

Cardoso


Exercise for COVID-19 Cardoso et al. Prophylaxis



Case control study with 307 severe COVID-19 ICU patients and 307 matched COVID-19 outpatients in Brazil, showing significantly higher risk of severe cases with low physical activity.

Cho


Exercise for COVID-19 Cho et al. Prophylaxis



Retrospective 6,288 COVID+ patients and 125,772 matched controls in South Korea, showing significantly lower risk of COVID-19 infection and mortality with higher physical activity.

Christensen


Exercise for COVID-19 Christensen et al. Prophylaxis



Prospective study of 2,690 adults in the UK Biobank showing lower cardiorespiratory fitness associated with COVID-19 mortality.

de Souza

Exercise for COVID-19 de Souza et al. Prophylaxis



Does physical activity reduce risk for COVID-19?

Retrospective 938 patients in Brazil (June - August 2020)

Lower hospitalization with higher activity levels (p=0.046)


de Souza et al., J. Science and Medici..., Sep 2021

c19early.org

Retrospective survey of 938 COVID-19 recovered patients in Brazil, showing lower hospitalization with physical activity. NCT04396353.

Ekblom-Bak

Exercise for COVID-19 Ekblom-Bak et al. Prophylaxis



Does physical activity reduce risk for COVID-19?

Retrospective study in Sweden

Lower severe cases with higher activity levels (p=0.023)


Ekblom-Bak et al., Int. J. Behavioral ..., Oct 2021

c19early.org

Retrospective 857 severe COVID-19 cases and matched controls in Sweden, showing lower risk of severe COVID-19 with higher cardiorespiratory fitness.

Fernandez

Exercise Fernandez et al. LATE TREATMENT



Is **late** treatment with exercise beneficial for COVID-19?

Retrospective 439 patients in Chile

Lower mortality with exercise (p=0.018)


Fernandez et al., J. Applied Physiology, Feb 2023

c19early.org

Retrospective 439 severe COVID-19 hospitalized patients with hypertension, 201 receiving a supervised exercise program, showing significantly lower mortality with exercise. Exercise included of aerobic, breathing, and musculoskeletal exercises, 3 to 4 times per week. There were significantly more control patients on beta-adrenergic blockers and thiazide diuretics.


There are many possible mechanisms of action, including improved circulation, stress reduction, hormone regulation, improved sleep, increased antioxidant levels, and increased nitric oxide levels in the respiratory system. Over-exercising may be detrimental and lead to impaired immune function.

Feter




Analysis of 237 COVID-19 patients in Brazil, showing lower risk of long COVID with physical activity.

Frish



Retrospective 3,038 bariatric surgery patients in Israel, showing higher risk of SARS-CoV-2 infection with vitamin D deficiency, and lower risk with physical activity.


Gao



Case control study in China with 105 cases and 210 matched controls, showing COVID-19 cases associated with physical activity ≥ 5 times per week. Authors note that people may choose gyms for exercise in winter, leading to higher exposure risk.

Gilley

Exercise for COVID-19 Gilley et al. Prophylaxis



Does physical activity reduce risk for COVID-19?

Retrospective 1,996 patients in the USA (September - December 2020)

More cases with higher activity levels (not stat. sig., $p=0.55$)


Gilley et al., JMIR Mental Health, Feb 2022

c19early.org

Retrospective survey of 1,997 college students in the USA, showing no significant difference in COVID-19 cases with exercise in unadjusted results.

Green

Exercise for COVID-19 Green et al. Prophylaxis



Does physical activity reduce risk for COVID-19?

Retrospective 113,075 patients in Israel (February - December 2020)

Fewer cases with higher activity levels ($p<0.000001$)


Green et al., European J. General Prac., Nov 2022

c19early.org

Retrospective 113,075 people in Israel, showing lower risk of COVID-19 cases with physical activity and a dose dependent response.

Halabchi

Exercise for COVID-19 Halabchi et al. Prophylaxis



Does physical activity reduce risk for COVID-19?

Retrospective 4,694 patients in Iran

Lower hospitalization with higher activity levels ($p=0.044$)

Halabchi et al., J. Physical Activity ..., Dec 2020

c19early.org

Retrospective 4,694 COVID-19 patients in Iran, showing lower risk of hospitalization and mortality with regular sports participation.

Hamdan

Exercise for COVID-19 Hamdan et al. Prophylaxis



Does physical activity reduce risk for COVID-19?

Retrospective 300 patients in Palestine

Lower hospitalization with higher activity levels (not stat. sig., $p=0.53$)


Hamdan et al., J. Int. Medical Research, Dec 2021

c19early.org

Retrospective 300 participants in Palestine, showing lower risk of hospitalization with physical activity, without statistical significance.

Hamer

Exercise for COVID-19 Hamer et al. Prophylaxis



Does physical activity reduce risk for COVID-19?

Retrospective study in the United Kingdom

Lower hospitalization with higher activity levels ($p=0.0004$)

Hamer et al., Brain, Behavior, and Imm., Jul 2020

c19early.org

UK Biobank retrospective analysis of 387,109 people, showing lower risk of COVID-19 hospitalization with physical activity.

Hamrouni

Exercise for COVID-19 Hamrouni et al. Prophylaxis



Does physical activity reduce risk for COVID-19?

Prospective study of 153,833 patients in the United Kingdom

Lower mortality with higher activity levels ($p=0.0093$)


Hamrouni et al., BMJ Open, November 2021

c19early.org

Prospective UK Biobank analysis, showing a history of low physical activity associated with COVID-19 mortality.

Hegazy

Exercise for COVID-19 Hegazy et al. Prophylaxis



Does physical activity reduce risk for COVID-19?

Retrospective 73 patients in Egypt (May 2021 - February 2022)

Fewer moderate/severe cases with higher activity levels ($p=0.0096$)

Hegazy et al., BMC Nutrition, October 2023

c19early.org

Retrospective 68 COVID-19 patients showing physical activity and healthier nutrition associated with lower COVID-19 severity.

Hegazy

Exercise for COVID-19 Hegazy et al. Prophylaxis



Does physical activity reduce risk for COVID-19?

Retrospective 200 patients in Egypt

Fewer moderate/severe cases with higher activity levels (not stat. sig., $p=0.11$)


Hegazy et al., British J. Nutrition, Jun 2021

c19early.org

Analysis of 200 mild and moderate COVID-19 outpatients showing an association between higher ESSAP scores (measuring exercise, sugar and prebiotic consumption, sleep, and antibiotic use) and milder COVID-19 disease. Authors find increased risk with daily yogurt containing probiotics. Probiotic intake based on yogurt only may be inaccurate. Authors hypothesize that commercial yogurt products may not contain sufficient beneficial bacteria or may be contaminated. Other research shows that probiotic food labels are often misleading — of 26 probiotic foods tested, only 5 contained Bifidobacterium in sufficient concentration for exhibiting a therapeutic effect³⁷. For sleep, authors compare <8 hours and ≥8 hours, while sleep for less than or longer than a recommended range may indicate increased risk.

Ho

Exercise for COVID-19 Ho et al. Prophylaxis



Does physical activity reduce risk for COVID-19?

Retrospective 138,475 patients in the United Kingdom


Lower hospitalization with higher activity levels ($p=0.007$)

Ho et al., BMJ Open, November 2020

c19early.org


UK Biobank retrospective 235,928 participants using walking pace as a proxy for physical fitness, showing lower risk of COVID-19 hospitalization with an average vs. slow walking pace.

Holt




Prospective survey-based study with 15,227 people in the UK, showing reduced risk of COVID-19 cases with lower impact physical activity.

Huang



Retrospective 164 COVID-19 patients and 188 controls in China, showing lower risk of cases with regular exercise.


Kapusta



Retrospective 568 convalescent COVID-19 patients in Poland, showing lower risk of severe cases with regular physical activity in the 3 months before COVID-19.

Kontopoulou

Exercise for COVID-19 Kontopoulou et al. Prophylaxis



Does physical activity reduce risk for COVID-19?

Retrospective 66 patients in Greece (November - December 2020)

Faster recovery with higher activity levels ($p < 0.000001$)


Kontopoulou et al., J. Personalized Me., Apr 2022

c19early.org

Retrospective 66 hospitalized COVID-19 patients in Greece, showing significantly improved recovery with a history of exercise in unadjusted results. Exercise after hospitalization was also associated with lower levels of dyspnea one month post hospitalization.

Latorre-Román

Exercise Latorre-Román et al. Prophylaxis



Does physical activity reduce risk for COVID-19?

Retrospective study in Spain

Lower hospitalization with higher activity levels (not stat. sig., $p = 0.05$)


Latorre-Román et al., Research in Spor., Jun 2021

c19early.org

Retrospective 420 people in Spain, showing lower risk of COVID-19 hospitalization with a history of physical activity.

Lee

Exercise for COVID-19 Lee et al. Prophylaxis



Does physical activity reduce risk for COVID-19?

Retrospective 52,365 patients in South Korea (Jan - Jul 2020)

Lower mortality ($p = 0.046$) and severe cases ($p = 0.03$)


Lee et al., British J. Sports Medicine, Jul 2021

c19early.org

Retrospective 212,768 adults in South Korea, showing lower risk of COVID-19 cases, severity, and mortality with physical activity. Notably, results for aerobic and muscle strengthening activities combined were much better than results for either one in isolation.

Lengelé


Exercise for COVID-19 Lengelé et al. Prophylaxis



Analysis of 241 adults >65yo in Belgium, showing lower risk of COVID-19 with a history of physical activity.

Li


Exercise for COVID-19 Li et al. Prophylaxis



Mendelian randomization study showing lower risk of severe COVID-19 with physical activity.

Lin

Exercise for COVID-19 Lin et al. Prophylaxis



Prospective survey analysis of 28,575 people in 99 countries, showing a lower risk of COVID-19 with a exercise, without statistical significance.

Lobelo

Exercise for COVID-19 Lobelo et al. Prophylaxis



Does physical activity reduce risk for COVID-19?

Retrospective 5,712 patients in Georgia (March - October 2020)

Lower hospitalization with higher activity levels ($p=0.022$)


Lobelo et al., BMJ Open, May 2021

c19early.org

Retrospective 5,712 COVID-19 patients in the USA, showing higher risk of COVID-19 hospitalization with a history of physical inactivity.

Malisoux

Exercise for COVID-19 Malisoux et al. Prophylaxis



Does physical activity reduce risk for COVID-19?

Retrospective 224 patients in Luxembourg (May 2020 - June 2021)

Lower progression with higher activity levels ($p=0.045$)


Malisoux et al., BMJ Open, April 2022

c19early.org

Retrospective 452 participants in Luxembourg, showing lower risk of moderate cases with higher physical activity.

Maltagliati

Exercise for COVID-19 Maltagliati et al. Prophylaxis



Does physical activity reduce risk for COVID-19?

Retrospective study in multiple countries

Lower hospitalization with higher activity levels ($p=0.02$)


Maltagliati et al., J. Sports Sciences, Aug 2021

c19early.org

Retrospective 3,139 adults >50 in Europe, with 66 COVID-19 hospitalizations, showing lower risk of hospitalization with higher physical activity and with higher muscle strength. Note that model 2 includes muscle strength which is correlated with physical activity³⁸.

Marcus


Exercise for COVID-19 Marcus et al. Prophylaxis



Prospective survey based study with 14,335 participants, showing lower risk of viral symptoms with regular exercise.

Mohsin


Exercise for COVID-19 Mohsin et al. Prophylaxis



Retrospective 1,500 COVID+ patients in Bangladesh, showing lower risk of severe cases with regular exercise in unadjusted results.

Muñoz-Vergara


Exercise Muñoz-Vergara et al. Prophylaxis



Prospective study of 61,557 adults aged 45+ years showing reduced risk of COVID-19 diagnosis and hospitalization for those meeting physical activity guidelines of ≥ 7.5 MET-hours/week before the pandemic compared to inactive individuals.

Nguyen

Exercise for COVID-19 Nguyen et al. Prophylaxis



Does physical activity reduce risk for COVID-19?

Retrospective 3,947 patients in Vietnam (February - March 2020)

Fewer symptomatic cases with higher activity levels ($p=0.00011$)


Nguyen et al., Nutrients, September 2021

c19early.org

Analysis of 3,947 participants in Vietnam, showing significantly lower risk of COVID-19-like symptoms with physical activity and with a healthy diet. The combination of being physically active and eating healthy reduced risk further compared to either alone. The analyzed period was Feb 14 to Mar 2, 2020, which may have been before testing was widely available.

Park

Exercise for COVID-19 Park et al. Prophylaxis



Does physical activity reduce risk for COVID-19?

Retrospective study in South Korea (January - August 2020)

Fewer cases with higher activity levels ($p=0.016$)


Park et al., Frontiers in Public Health, Feb 2023

c19early.org

Retrospective 4,363 COVID-19 patients and 67,125 controls in South Korea, showing higher risk of mortality and cases with insufficient physical activity.

Paul

Exercise Paul et al. Prophylaxis LONG COVID



Does physical activity reduce the risk of long COVID (PASC)?

Retrospective 1,811 patients in the United Kingdom

Lower PASC with higher activity levels (not stat. sig., $p=0.16$)


Paul et al., medRxiv, April 2022

c19early.org

Retrospective 1,811 COVID-19 patients in the UK, showing lower risk of self-reported long COVID with 3+ hours of exercise per week in the month before infection, without statistical significance ($p=0.16$).

Pavlidou


Exercise for COVID-19 Pavlidou et al. Prophylaxis



Retrospective 5,197 Greek adults over 65. After adjustment for confounders, COVID-19 infection was independently associated with poor sleep, low physical activity, low Mediterranean diet adherence, living in urban areas, smoking, obesity, depression, anxiety, stress, and poor health-related quality of life.

Pitanga


Exercise for COVID-19 Pitanga et al. Prophylaxis



Retrospective 4,476 participants in Brazil, showing lower risk of COVID-19 cases with a history of physical activity, statistically significant only for those following specific practices to protect against COVID-19.

Pływaczewska-Jakubowska


Exercise Pływaczewska-Jakubowska et al. Prophylaxis



Retrospective 1,847 COVID+ patients in Poland, showing no significant difference in moderate/severe cases with physical activity. Hospitalized patients were excluded.

Reis

Exercise for COVID-19 Reis et al. Prophylaxis



Does physical activity reduce risk for COVID-19?

Retrospective 546 patients in the USA (December 2020 - February 2021)

Lower hospitalization with higher activity levels (not stat. sig., $p=0.18$)


Reis et al., American J. Lifestyle Med., Oct 2022

c19early.org

Retrospective 546 COVID+ patients in the USA, showing lower risk of hospitalization with higher frequency of strength training, without statistical significance.

Rocha

Exercise Rocha et al. Prophylaxis LONG COVID



Does physical activity reduce the risk of long COVID (PASC)?

Retrospective 2,919 patients in Brazil (December 2020 - March 2021)

Lower PASC with higher activity levels ($p=0.05$)


Rocha et al., BMC Sports Science, Medi., Dec 2023

c19early.org

Retrospective 2,919 non-hospitalized COVID-19 patients in Brazil showing remaining physically active before and after COVID-19 infection reduces the probability of experiencing long COVID symptoms, particularly those affecting the musculoskeletal, neurological and respiratory systems.

Saadeh

Exercise for COVID-19 Saadeh et al. Prophylaxis



Does physical activity reduce risk for COVID-19?

Retrospective 587 patients in Sweden (March - June 2020)


No significant difference in symptomatic cases

Saadeh et al., Aging Clinical and Expe., Oct 2021

c19early.org


Retrospective 904 patients in Sweden, showing higher risk of COVID-19-like symptoms with poor muscle strength. Risk was slightly higher for physical inactivity, without statistical significance.

Salgado-Aranda




Retrospective 520 COVID-19 patients in Spain, showing significantly lower mortality with a history of physical activity.

Sallis




Retrospective 48,440 COVID-19 patients in the USA, showing significantly lower mortality, ICU admission, and hospitalization with exercise.

Sanchez




Retrospective 29,875 university staff and students in Spain, 3,662 with data, showing lower risk of COVID-19 symptoms for people that exercise. Exercise more than 5 days/week was the most protective, and intense exercise was more effective than moderate exercise.

Schmidt




Retrospective 5,338 individuals with confirmed contact with a COVID-19 patient, showing lower risk of COVID-19 with exercise.

Steenkamp




Retrospective 65,361 COVID-19 patients in South Africa, showing significantly lower hospitalization, ICU admission, ventilation, and mortality with exercise.

Sutkowska




Prospective study of 131 hospitalized patients in Poland, showing lower mortality and improved recovery with a history of higher physical activity.

Tavakol




Retrospective 206 patients in Iran, showing COVID-19 disease severity associated with lower physical activity.

Tret'yakov




Retrospective 293 COVID+ patients in Russia, showing lower risk of severe COVID-19 for individuals who regularly practice aerobic training in unadjusted results.

Tsuzuki




Retrospective 4,868 elderly COVID-19 patients in Japan, showing higher risk of severe cases with poor physical activity status.

Wang




Prospective study of 68,896 UK Biobank participants with COVID-19 showing adherence to a healthy lifestyle prior to infection, characterized by 10 factors including adequate physical activity and sleep, not smoking, and a healthy BMI, was associated with a significantly lower risk of mortality, hospitalization, and post-COVID multisystem sequelae. Risk decreased monotonically for increasing numbers of healthy lifestyle factors from 5-10. Reduced risks were evident across cardiovascular, metabolic, neurologic, respiratory, and other disorders over 210 days following infection, during both acute and post-acute phases, regardless of age, sex, ethnicity, test setting, vaccination status, or SARS-CoV-2 variant.

Wang




Prospective analysis of 32,249 women, showing lower risk of PASC with a healthy lifestyle, in a dose-dependent manner. Participants with 5 or 6 healthy lifestyle factors had significantly lower COVID-19 hospitalization and PASC. BMI and sleep were independently associated with risk of PASC.

Yates




UK Biobank retrospective 412,596 people, showing severe COVID-19 and COVID-19 mortality inversely associated with self-reported walking pace.

Young




Retrospective 194,191 COVID-19 patients in the USA, showing lower risk of hospitalization and mortality with physical activity, with a dose response relationship.

Yuan




Retrospective 164 COVID-19 patients in China, showing physical inactivity associated with an increased risk of severe COVID-19.

Zhang



UK Biobank retrospective showing significantly lower COVID-19 cases with objectively measured physical activity.

Šebić



Retrospective 100 COVID-19 patients in Bosnia and Herzegovina, showing lower symptom severity and faster recovery with a history of regular physical activity.

Appendix 1. Methods and Data

We perform ongoing searches of PubMed, medRxiv, Europe PMC, ClinicalTrials.gov, The Cochrane Library, Google Scholar, Research Square, ScienceDirect, Oxford University Press, the reference lists of other studies and meta-analyses, and submissions to the site c19early.org. Search terms are (exercise OR "physical activity") AND COVID-19. Automated searches are performed twice daily, with all matches reviewed for inclusion. All studies regarding the use of exercise for COVID-19 that report a comparison with a control group are included in the main analysis. Sensitivity analysis is performed, excluding studies with major issues, epidemiological studies, and studies with minimal available information. Studies with major unexplained data issues, for example major outcome data that is impossible to be correct with no response from the authors, are excluded. This is a living analysis and is updated regularly.

We extracted effect sizes and associated data from all studies. If studies report multiple kinds of effects then the most serious outcome is used in pooled analysis, while other outcomes are included in the outcome specific analyses. For example, if effects for mortality and cases are reported then they are both used in specific outcome analyses, while mortality is used for pooled analysis. If symptomatic results are reported at multiple times, we use the latest time, for

example if mortality results are provided at 14 days and 28 days, the results at 28 days have preference. Mortality alone is preferred over combined outcomes. Outcomes with zero events in both arms are not used, the next most serious outcome with one or more events is used. For example, in low-risk populations with no mortality, a reduction in mortality with treatment is not possible, however a reduction in hospitalization, for example, is still valuable. Clinical outcomes are considered more important than viral outcomes. When basically all patients recover in both treatment and control groups, preference for viral clearance and recovery is given to results mid-recovery where available. After most or all patients have recovered there is little or no room for an effective treatment to do better, however faster recovery is valuable. An IPD meta-analysis confirms that intermediate viral load reduction is more closely associated with hospitalization/death than later viral load reduction³⁹. If only individual symptom data is available, the most serious symptom has priority, for example difficulty breathing or low SpO₂ is more important than cough. When results provide an odds ratio, we compute the relative risk when possible, or convert to a relative risk according to *Zhang (B) et al.* Reported confidence intervals and *p*-values are used when available, and adjusted values are used when provided. If multiple types of adjustments are reported propensity score matching and multivariable regression has preference over propensity score matching or weighting, which has preference over multivariable regression. Adjusted results have preference over unadjusted results for a more serious outcome when the adjustments significantly alter results. When needed, conversion between reported *p*-values and confidence intervals followed *Altman, Altman (B)*, and Fisher's exact test was used to calculate *p*-values for event data. If continuity correction for zero values is required, we use the reciprocal of the opposite arm with the sum of the correction factors equal to 1⁴³. Results are expressed with RR < 1.0 favoring treatment, and using the risk of a negative outcome when applicable (for example, the risk of death rather than the risk of survival). If studies only report relative continuous values such as relative times, the ratio of the time for the treatment group versus the time for the control group is used. Calculations are done in Python (3.13.5) with scipy (1.16.0), pythonmeta (1.26), numpy (2.3.1), statsmodels (0.14.4), and plotly (6.2.0).




Figure 20. Mid-recovery results can more accurately reflect efficacy when almost all patients recover. *Mateja et al.* confirm that intermediate viral load results more accurately reflect hospitalization/death.

Forest plots are computed using PythonMeta⁴⁴ with the DerSimonian and Laird random effects model (the fixed effect assumption is not plausible in this case) and inverse variance weighting. Results are presented with 95% confidence intervals. Heterogeneity among studies was assessed using the *I*² statistic. Mixed-effects meta-regression results are computed with R (4.4.0) using the metafor (4.6-0) and rms (6.8-0) packages, and using the most serious sufficiently powered outcome. For all statistical tests, a *p*-value less than 0.05 was considered statistically significant. Grobid 0.8.2 is used to parse PDF documents.

We have classified studies as early treatment if most patients are not already at a severe stage at the time of treatment (for example based on oxygen status or lung involvement), and treatment started within 5 days of the onset of symptoms. If studies contain a mix of early treatment and late treatment patients, we consider the treatment time of patients contributing most to the events (for example, consider a study where most patients are treated early but late treatment patients are included, and all mortality events were observed with late treatment patients). We note that a shorter time may be preferable. Antivirals are typically only considered effective when used within a shorter timeframe, for example 0-36 or 0-48 hours for oseltamivir, with longer delays not being effective^{45,46}.

We received no funding, this research is done in our spare time. We have no affiliations with any pharmaceutical companies or political parties.

A summary of study results is below. Please submit updates and corrections at <https://c19early.org/exmeta.html>.

Late treatment

Effect extraction follows pre-specified rules as detailed above and gives priority to more serious outcomes. For pooled analyses, the first (most serious) outcome is used, which may differ from the effect a paper focuses on. Other outcomes are used in outcome specific analyses.

<i>Fernandez</i> , 2/2/2023, retrospective, Chile, peer-reviewed, 10 authors.	risk of death, 47.5% lower, RR 0.53, $p = 0.02$, high activity levels 16 of 201 (8.0%), low activity levels 62 of 238 (26.1%), NNT 5.5, adjusted per study, odds ratio converted to relative risk, multivariable.
---	--

Prophylaxis

Effect extraction follows pre-specified rules as detailed above and gives priority to more serious outcomes. For pooled analyses, the first (most serious) outcome is used, which may differ from the effect a paper focuses on. Other outcomes are used in outcome specific analyses.

<i>af Geijerstam</i> , 7/5/2021, prospective, Sweden, peer-reviewed, 9 authors, study period March 2020 - September 2020.	risk of death, 50.0% lower, OR 0.50, $p = 0.005$, high vs. low fitness, model 7, RR approximated with OR.
	risk of ICU admission, 40.0% lower, OR 0.60, $p < 0.001$, high vs. low fitness, model 7, RR approximated with OR.
	risk of hospitalization, 27.0% lower, OR 0.73, $p < 0.001$, high vs. low fitness, model 7, RR approximated with OR.
<i>Ahmadi</i> , 8/31/2021, retrospective, United Kingdom, peer-reviewed, 5 authors.	risk of death, 30.0% lower, RR 0.70, $p = 0.005$, adjusted per study, sufficient vs. inactive, model 2, multivariable.
<i>Akbar</i> , 11/7/2023, retrospective, Qatar, peer-reviewed, mean age 40.3, 9 authors, study period March 2020 - September 2020.	risk of case, 7.0% lower, OR 0.93, $p = 0.40$, high activity levels 3,333, low activity levels 3,333, adjusted per study, T3 vs. T1, multivariable, model 2, RR approximated with OR.
<i>Almansour</i> , 2/17/2022, retrospective, Saudi Arabia, peer-reviewed, 12 authors, study period April 2020 - June 2020.	risk of case, 5.7% lower, RR 0.94, $p = 0.85$, high activity levels 35 of 71 (49.3%), low activity levels 38 of 71 (53.5%), NNT 24, adjusted per study, odds ratio converted to relative risk, multivariable.
<i>Antunes</i> , 6/11/2022, retrospective, Brazil, peer-reviewed, survey, 5 authors, study period September 2020 - December 2020.	risk of ICU admission, 80.2% lower, RR 0.20, $p = 0.06$, high activity levels 1 of 14 (7.1%), low activity levels 9 of 25 (36.0%), NNT 3.5.
	risk of miscellaneous, 40.5% lower, RR 0.60, $p = 0.48$, high activity levels 3 of 14 (21.4%), low activity levels 9 of 25 (36.0%), NNT 6.9, CT abnormalities >50%.
	risk of miscellaneous, 72.5% lower, RR 0.27, $p = 0.04$, high activity levels 2 of 14 (14.3%), low activity levels 13 of 25 (52.0%), NNT 2.7, CT abnormalities 25-50%.
	hospitalization time, 43.4% lower, relative time 0.57, $p = 0.03$, high activity levels 14, low activity levels 25.
	miscellaneous, 25.5% lower, relative time 0.74, $p = 0.02$, high activity levels 14, low activity levels 25.
<i>Baynouna AlKetbi</i> , 8/23/2021, retrospective, United Arab Emirates, peer-reviewed, 16 authors.	risk of death, 98.5% lower, OR 0.01, $p = 0.049$, adjusted per study, multivariable, RR approximated with OR.
<i>Beydoun</i> , 3/12/2022, retrospective, USA, peer-reviewed, survey, 7 authors.	risk of case, 43.0% lower, OR 0.57, $p = 0.05$, high activity levels 1,710, low activity levels 448, adjusted per study, multivariable, >1/week vs. none, model 2, RR approximated with OR.
	risk of case, 62.0% lower, OR 0.38, $p = 0.010$, high activity levels 672, low activity levels 448, adjusted per study, multivariable, 1-4/mon vs. none, model 2, RR approximated with OR.

<i>Bielik</i> , 7/4/2021, retrospective, Slovakia, peer-reviewed, survey, 3 authors, study period 7 December, 2020 - 12 December, 2020.	risk of moderate case, 30.4% lower, RR 0.70, $p = 0.10$, high activity levels 775, low activity levels 365, adjusted per study, physically active group.
	risk of case, 9.1% higher, RR 1.09, $p = 0.36$, high activity levels 775, low activity levels 365, adjusted per study, physically active group.
<i>Brandenburg</i> , 7/1/2021, retrospective, multiple countries, peer-reviewed, survey, 4 authors.	risk of hospitalization, 6.0% higher, OR 1.06, $p = 0.60$, high activity levels 102, low activity levels 39, adjusted per study, multivariable, PA, >1h vigorous vs. no/low, RR approximated with OR.
	risk of hospitalization, 78.0% lower, OR 0.22, $p = 0.05$, high activity levels 177, low activity levels 34, adjusted per study, multivariable, CRF, 6.2-8.7 vs. >10, RR approximated with OR.
	risk of hospitalization, 64.0% lower, OR 0.36, $p = 0.04$, high activity levels 97, low activity levels 34, adjusted per study, multivariable, CRF, 8.7-10 vs. >10, RR approximated with OR.
	risk of severe case, 35.0% lower, OR 0.65, $p = 0.30$, high activity levels 102, low activity levels 39, adjusted per study, multivariable, PA, >1h vigorous vs. no/low, RR approximated with OR.
	risk of severe case, 24.0% lower, OR 0.76, $p = 0.60$, high activity levels 52, low activity levels 34, adjusted per study, multivariable, CRF, 4.4-6.2 vs. >10, RR approximated with OR.
<i>Brawner</i> , 10/10/2020, retrospective, USA, peer-reviewed, 10 authors, study period 29 February, 2020 - 30 May, 2020, excluded in exclusion analyses: unadjusted results with no group details.	risk of hospitalization, 74.2% lower, OR 0.26, $p = 0.001$, unadjusted, inverted to make OR<1 favor high activity levels, highest fitness quartile vs. lowest fitness quartile, RR approximated with OR.
<i>Cardoso</i> , 5/9/2023, retrospective, Brazil, peer-reviewed, 6 authors, study period April 2020 - February 2022.	risk of severe case, 73.0% lower, OR 0.27, $p < 0.001$, high activity levels 307, low activity levels 307, adjusted per study, inverted to make OR<1 favor high activity levels, case control OR, moderate/high vs. low physical activity, multivariable.
<i>Cho</i> , 4/6/2021, retrospective, South Korea, peer-reviewed, 9 authors.	risk of death, 53.0% lower, OR 0.47, $p = 0.01$, high activity levels 17 of 48 (35.4%) cases, 3,223 of 4,536 (71.1%) controls, case control OR, moderate to vigorous vs. inactive.
	risk of case, 10.0% lower, OR 0.90, $p < 0.001$, high activity levels 3,223 of 4,536 (71.1%) cases, 68,609 of 92,587 (74.1%) controls, NNT 142, case control OR, moderate to vigorous vs. inactive.
<i>Christensen</i> , 5/5/2021, prospective, United Kingdom, peer-reviewed, 5 authors, study period 16 March, 2020 - 26 July, 2020.	risk of death, 63.0% lower, RR 0.37, $p = 0.02$, high activity levels 543, low activity levels 529, adjusted per study, high fitness vs. low fitness, multivariable.
	risk of case, 23.0% lower, RR 0.77, $p = 0.20$, high activity levels 55 of 543 (10.1%), low activity levels 77 of 529 (14.6%), NNT 23, adjusted per study, high fitness vs. low fitness, multivariable.
<i>de Souza</i> , 9/30/2021, retrospective, Brazil, peer-reviewed, 8 authors, study period June 2020 - August 2020, trial NCT04396353 (history).	risk of mechanical ventilation, 73.2% lower, RR 0.27, $p = 0.07$, high activity levels 3 of 611 (0.5%), low activity levels 6 of 327 (1.8%), NNT 74, unadjusted, excluded in exclusion analyses: unadjusted results with no group details.

	risk of hospitalization, 34.3% lower, RR 0.66, $p = 0.046$, high activity levels 49 of 611 (8.0%), low activity levels 42 of 327 (12.8%), NNT 21, adjusted per study, sufficient vs. insufficient, model 3, multivariable.
<i>Eklom-Bak</i> , 10/19/2021, retrospective, Sweden, peer-reviewed, 13 authors.	risk of severe case, 47.6% lower, OR 0.52, $p = 0.02$, inverted to make $OR < 1$ favor high activity levels, case control OR, model 3, high vs. very low CRF.
<i>Feter</i> , 6/13/2023, retrospective, Brazil, peer-reviewed, survey, mean age 37.1, 17 authors.	risk of PASC, 26.0% lower, RR 0.74, $p = 0.02$, high activity levels 52, low activity levels 95, adjusted per study, before and during pandemic, multivariable.
	risk of PASC, 17.0% lower, RR 0.83, $p = 0.04$, high activity levels 67, low activity levels 170, adjusted per study, during pandemic, multivariable.
<i>Frish</i> , 6/15/2023, retrospective, Israel, peer-reviewed, 7 authors, study period 1 February, 2020 - 31 December, 2020.	risk of case, 53.0% lower, OR 0.47, $p = 0.04$, high activity levels 212, low activity levels 1,202, adjusted per study, >3 times per week vs. none, multivariable, RR approximated with OR.
<i>Gao</i> , 11/5/2020, retrospective, China, peer-reviewed, survey, median age 55.0, 11 authors, study period 10 February, 2020 - 1 March, 2020.	risk of case, 105.0% higher, HR 2.05, $p < 0.001$, high activity levels 59 of 105 (56.2%) cases, 69 of 210 (32.9%) controls, case control OR, Cox proportional hazards.
<i>Gilley</i> , 2/10/2022, retrospective, USA, peer-reviewed, survey, 21 authors, study period September 2020 - December 2020, trial NCT04766788 (history).	risk of case, 41.8% higher, RR 1.42, $p = 0.55$, high activity levels 172 of 1,917 (9.0%), low activity levels 5 of 79 (6.3%), unadjusted.
<i>Green</i> , 11/7/2022, retrospective, Israel, peer-reviewed, 9 authors, study period 1 February, 2020 - 31 December, 2020.	risk of case, 41.7% lower, RR 0.58, $p < 0.001$, high activity levels 1,267 of 11,144 (11.4%), low activity levels 16,198 of 101,931 (15.9%), adjusted per study, odds ratio converted to relative risk, >3 times per week vs. none, multivariable.
<i>Halabchi (B)</i> , 12/1/2020, retrospective, Iran, peer-reviewed, 8 authors.	risk of death, 88.8% lower, RR 0.11, $p = 0.08$, high activity levels 0 of 249 (0.0%), low activity levels 79 of 4,445 (1.8%), NNT 56, adjusted per study, odds ratio converted to relative risk, multivariable.
	risk of hospitalization, 28.3% lower, RR 0.72, $p = 0.04$, high activity levels 30 of 249 (12.0%), low activity levels 878 of 4,445 (19.8%), adjusted per study, odds ratio converted to relative risk, multivariable.
<i>Hamdan</i> , 12/23/2021, retrospective, Palestine, peer-reviewed, survey, mean age 30.5, 7 authors.	risk of hospitalization, 16.4% lower, RR 0.84, $p = 0.53$, high activity levels 22 of 128 (17.2%), low activity levels 37 of 172 (21.5%), NNT 23, adjusted per study, odds ratio converted to relative risk, multivariable.
<i>Hamer</i> , 7/31/2020, retrospective, United Kingdom, peer-reviewed, 4 authors.	risk of hospitalization, 27.5% lower, RR 0.72, $p < 0.001$, adjusted per study, inverted to make $RR < 1$ favor high activity levels, model 2, sufficient vs. no activity, multivariable.
	risk of hospitalization, 33.8% lower, RR 0.66, $p < 0.001$, adjusted per study, inverted to make $RR < 1$ favor high activity levels, model 1, sufficient vs. no activity, multivariable.
<i>Hamrouni</i> , 11/3/2021, prospective, United Kingdom, peer-reviewed, 5 authors.	risk of death, 29.0% lower, RR 0.71, $p = 0.009$, high activity levels 138 of 106,006 (0.1%), low activity levels 109 of 47,827 (0.2%), adjusted per study, inverted to make $RR < 1$ favor high activity levels, odds ratio converted to relative risk, high vs. low physical activity, multivariable.

<p><i>Hegazy</i>, 10/2/2023, retrospective, Egypt, peer-reviewed, 7 authors, study period May 2021 - February 2022, trial NCT04447144 (history), excluded in exclusion analyses: unadjusted results with no group details.</p>	<p>risk of moderate case, 54.0% lower, RR 0.46, $p = 0.010$, high activity levels 15 of 50 (30.0%), low activity levels 15 of 23 (65.2%), NNT 2.8, active vs. inactive.</p>
	<p>risk of moderate case, 97.1% lower, RR 0.03, $p = 0.02$, high activity levels 0 of 7 (0.0%), low activity levels 30 of 61 (49.2%), NNT 2.0, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm), moderate vs. low/inactive.</p>
<p><i>Hegazy (B)</i>, 6/7/2021, retrospective, Egypt, peer-reviewed, 10 authors, trial NCT04447144 (history).</p>	<p>risk of moderate/severe case, 45.6% lower, RR 0.54, $p = 0.11$, high activity levels 24 of 82 (29.3%), low activity levels 7 of 13 (53.8%), NNT 4.1, >10min/day vs. none.</p>
<p><i>Ho</i>, 11/19/2020, retrospective, United Kingdom, peer-reviewed, survey, 13 authors.</p>	<p>risk of hospitalization, 34.6% lower, RR 0.65, $p = 0.007$, high activity levels 213 of 123,588 (0.2%), low activity levels 59 of 14,887 (0.4%), adjusted per study, inverted to make $RR < 1$ favor high activity levels, model 2, average vs. slow walking pace, multivariable.</p>
<p><i>Holt</i>, 3/30/2021, prospective, United Kingdom, peer-reviewed, 34 authors, study period 1 May, 2020 - 5 February, 2021, trial NCT04330599 (history) (COVIDENCE UK).</p>	<p>risk of case, 17.0% lower, OR 0.83, $p = 0.18$, adjusted per study, fully adjusted, ≥ 2 hours lower impact physical activity vs. 0 hours, RR approximated with OR.</p>
<p><i>Huang</i>, 11/30/2021, retrospective, China, peer-reviewed, survey, 5 authors, study period 10 February, 2020 - 28 March, 2020.</p>	<p>risk of severe case, 46.8% lower, RR 0.53, $p = 0.18$, high activity levels 7 of 74 (9.5%), low activity levels 16 of 90 (17.8%), NNT 12, unadjusted, exercise habit, ≥ 1 time per week, excluded in exclusion analyses: unadjusted results with no group details.</p>
	<p>risk of severe case, 8.0% lower, RR 0.92, $p = 1.00$, high activity levels 3 of 23 (13.0%), low activity levels 20 of 141 (14.2%), NNT 88, unadjusted, ≥ 30 minutes ≥ 3 times per week, excluded in exclusion analyses: unadjusted results with no group details.</p>
	<p>risk of case, 65.9% lower, OR 0.34, $p = 0.004$, adjusted per study, inverted to make $OR < 1$ favor high activity levels, case control OR, regular exercise, multivariable.</p>
<p><i>Kapusta</i>, 12/12/2022, retrospective, Poland, peer-reviewed, survey, mean age 70.4, 7 authors, study period 1 March, 2020 - 30 August, 2020, trial NCT05018052 (history).</p>	<p>risk of severe case, 70.9% lower, OR 0.29, $p = 0.001$, high activity levels 181, low activity levels 387, inverted to make $OR < 1$ favor high activity levels, RR approximated with OR.</p>
<p><i>Kontopoulou</i>, 4/17/2022, retrospective, Greece, peer-reviewed, survey, 4 authors, study period November 2020 - December 2020, excluded in exclusion analyses: unadjusted results with no group details.</p>	<p>recovery time, 66.2% lower, relative time 0.34, $p < 0.001$, high activity levels mean 22.0 (± 14.0) $n=42$, low activity levels mean 65.0 (± 32.0) $n=24$.</p>
	<p>relative dyspnea after hospitalization, 66.7% better, RR 0.33, $p < 0.001$, high activity levels mean 1.0 (± 1.0) $n=42$, low activity levels mean 3.0 (± 1.0) $n=24$, inverted to make $RR < 1$ favor high activity levels.</p>
<p><i>Latorre-Román</i>, 6/15/2021, retrospective, Spain, peer-reviewed, survey, 7 authors.</p>	<p>risk of hospitalization, 76.0% lower, OR 0.24, $p = 0.05$, moderate physical activity, >150 min per week, RR approximated with OR.</p>
	<p>risk of hospitalization, 87.0% lower, OR 0.13, $p = 0.07$, moderate physical activity, 30-150 min per week, RR approximated with OR.</p>

Lee, 7/22/2021, retrospective, South Korea, peer-reviewed, 25 authors, study period 1 January, 2020 - 31 July, 2020.	risk of death, 74.0% lower, RR 0.26, $p = 0.046$, high activity levels 2 of 11,072 (0.0%), low activity levels 32 of 41,293 (0.1%), NNT 1683, adjusted per study, odds ratio converted to relative risk, model 2, aerobic and muscle strengthening vs. insufficient aerobic and muscle strengthening, multivariable.
	risk of severe case, 57.8% lower, RR 0.42, $p = 0.03$, high activity levels 39 of 11,072 (0.4%), low activity levels 273 of 41,293 (0.7%), adjusted per study, odds ratio converted to relative risk, model 2, aerobic and muscle strengthening vs. insufficient aerobic and muscle strengthening, multivariable.
	risk of case, 15.6% lower, RR 0.84, $p = 0.03$, high activity levels 291 of 11,072 (2.6%), low activity levels 1,293 of 41,293 (3.1%), NNT 199, adjusted per study, odds ratio converted to relative risk, model 2, aerobic and muscle strengthening vs. insufficient aerobic and muscle strengthening, multivariable.
Lengel�, 10/23/2021, prospective, Belgium, peer-reviewed, median age 75.6, 8 authors, study period March 2020 - April 2021.	risk of case, 73.6% lower, RR 0.26, $p = 0.03$, high activity levels 23 of 229 (10.0%), low activity levels 4 of 12 (33.3%), NNT 4.3, inverted to make $RR < 1$ favor high activity levels, odds ratio converted to relative risk.
Li (B), 2/3/2021, retrospective, United Kingdom, peer-reviewed, 2 authors, per SD increase.	risk of severe case, 81.0% lower, OR 0.19, $p = 0.02$, RR approximated with OR.
	risk of hospitalization, 56.0% lower, OR 0.44, $p = 0.07$, RR approximated with OR.
Lin, 9/21/2021, prospective, multiple countries, peer-reviewed, survey, 19 authors, study period 26 March, 2020 - 8 October, 2020.	risk of case, 47.4% lower, OR 0.53, $p = 0.40$, inverted to make $OR < 1$ favor high activity levels, exercise ≥ 1 /month vs. exercise < 1 /month, RR approximated with OR.
Lobelo, 5/19/2021, retrospective, Georgia, peer-reviewed, 7 authors, study period 3 March, 2020 - 29 October, 2020.	risk of hospitalization, 20.0% lower, OR 0.80, $p = 0.02$, high activity levels 2,121, low activity levels 1,648, adjusted per study, inverted to make $OR < 1$ favor high activity levels, active vs. inactive, multivariable, RR approximated with OR.
Malisoux, 4/29/2022, retrospective, Luxembourg, peer-reviewed, survey, median age 42.0, 6 authors, study period May 2020 - June 2021, trial NCT04380987 (history).	risk of progression, 63.0% lower, OR 0.37, $p = 0.045$, high activity levels 115, low activity levels 108, moderate case, > 82 vs. < 30 MET-hour/week, RR approximated with OR.
	risk of progression, 52.0% lower, OR 0.48, $p = 0.14$, high activity levels 116, low activity levels 108, moderate case, $> 52-82$ vs. < 30 MET-hour/week, RR approximated with OR.
	risk of progression, 43.0% lower, OR 0.57, $p = 0.28$, high activity levels 113, low activity levels 108, moderate case, $30-52$ vs. < 30 MET-hour/week, RR approximated with OR.
Maltagliati, 8/11/2021, retrospective, multiple countries, peer-reviewed, survey, 8 authors.	risk of hospitalization, 52.0% lower, OR 0.48, $p = 0.02$, adjusted per study, model 1, more than once a week vs. hardly ever or never, multivariable, RR approximated with OR.
Marcus, 6/17/2021, prospective, multiple countries, peer-reviewed, survey, 12 authors, study period 26 March, 2020 - 3 May, 2020.	risk of symptomatic case, 42.1% lower, RR 0.58, $p < 0.001$, high activity levels 240 of 10,627 (2.3%), low activity levels 134 of 3,708 (3.6%), NNT 74, adjusted per study, odds ratio converted to relative risk, multivariable.
Mohsin, 9/30/2021, retrospective, Bangladesh, peer-reviewed, survey, 10 authors, study period November 2020 - April 2021, excluded in exclusion analyses: unadjusted results with no group details.	risk of severe case, 19.0% lower, RR 0.81, $p = 0.04$, high activity levels 86 of 258 (33.3%), low activity levels 224 of 544 (41.2%), NNT 13, exercise > 30 minutes.

	risk of severe case, 0.9% higher, RR 1.01, $p = 0.91$, high activity levels 290 of 698 (41.5%), low activity levels 224 of 544 (41.2%), exercise <30 minutes.
<i>Muñoz-Vergara</i> , 2/13/2024, prospective, USA, peer-reviewed, 7 authors.	risk of hospitalization, 26.7% lower, RR 0.73, $p = 0.002$, high activity levels 332 of 42,159 (0.8%), low activity levels 203 of 12,405 (1.6%), adjusted per study, odds ratio converted to relative risk, sufficiently active vs. inactive, multivariable, model 3.
	risk of case, 9.1% lower, RR 0.91, $p = 0.004$, high activity levels 3,898 of 42,159 (9.2%), low activity levels 1,293 of 12,405 (10.4%), NNT 85, adjusted per study, odds ratio converted to relative risk, sufficiently active vs. inactive, multivariable, model 3.
<i>Nguyen</i> , 9/18/2021, retrospective, Vietnam, peer-reviewed, survey, 17 authors, study period 14 February, 2020 - 2 March, 2020.	risk of symptomatic case, 20.3% lower, RR 0.80, $p < 0.001$, high activity levels 904 of 2,836 (31.9%), low activity levels 483 of 1,111 (43.5%), NNT 8.6, adjusted per study, odds ratio converted to relative risk, active vs. inactive, COVID-19-like symptoms, multivariable.
<i>Park</i> , 2/14/2023, retrospective, South Korea, peer-reviewed, survey, 4 authors, study period 1 January, 2020 - 14 August, 2020.	risk of death, 25.6% lower, OR 0.74, $p = 0.08$, inverted to make OR<1 favor high activity levels, sufficient vs. insufficient PA, model 3, RR approximated with OR.
	risk of death, 38.4% lower, OR 0.62, $p = 0.02$, inverted to make OR<1 favor high activity levels, sufficient vs. insufficient PA, model 2, RR approximated with OR.
	risk of case, 7.2% lower, OR 0.93, $p = 0.02$, inverted to make OR<1 favor high activity levels, sufficient vs. insufficient PA, model 3, RR approximated with OR.
	risk of case, 10.4% lower, OR 0.90, $p < 0.001$, inverted to make OR<1 favor high activity levels, sufficient vs. insufficient PA, model 2, RR approximated with OR.
<i>Paul</i> , 4/13/2022, retrospective, United Kingdom, preprint, survey, 2 authors.	risk of long COVID, 38.1% lower, RR 0.62, $p = 0.16$, adjusted per study, odds ratio converted to relative risk, 3+ hours per week vs. none, multivariable, model 4, control prevalence approximated with overall prevalence.
	risk of long COVID, 4.1% lower, RR 0.96, $p = 0.89$, adjusted per study, odds ratio converted to relative risk, ≤2 hours per week vs. none, multivariable, model 4, control prevalence approximated with overall prevalence.
<i>Pavlidou</i> , 11/9/2023, retrospective, Greece, peer-reviewed, 14 authors.	risk of case, 42.2% lower, OR 0.58, $p = 0.001$, high activity levels 902, low activity levels 4,295, adjusted per study, inverted to make OR<1 favor high activity levels, high vs. low/moderate IPAQ, multivariable, RR approximated with OR.
<i>Pitanga</i> , 10/29/2022, retrospective, Brazil, peer-reviewed, survey, 11 authors.	risk of case, 33.0% lower, OR 0.67, $p = 0.05$, high activity levels 1,469, low activity levels 1,552, combined results with and without protection practices, RR approximated with OR.
<i>Pływaczewska-Jakubowska</i> , 10/24/2022, retrospective, Poland, peer-reviewed, median age 51.0, 5 authors, study period May 2020 - January 2022.	risk of moderate/severe case, 11.0% lower, OR 0.89, $p = 0.30$, high activity levels 490, low activity levels 1,357, adjusted per study, multivariable, model 3, RR approximated with OR.

	risk of PASC, 14.0% lower, OR 0.86, $p = 0.24$, high activity levels 389, low activity levels 1,128, adjusted per study, multivariable, model 3, RR approximated with OR.
Reis, 10/24/2022, retrospective, USA, peer-reviewed, survey, 6 authors, study period December 2020 - February 2021.	risk of hospitalization, 40.7% lower, RR 0.59, $p = 0.18$, high activity levels 9 of 241 (3.7%), low activity levels 29 of 305 (9.5%), adjusted per study, inverted to make $RR < 1$ favor high activity levels, odds ratio converted to relative risk, strength training 2+ /week vs. <2, multivariable.
Rocha, 12/14/2023, retrospective, Brazil, peer-reviewed, 6 authors, study period December 2020 - March 2021.	risk of PASC, 20.0% lower, OR 0.80, $p = 0.05$, high activity levels 388, low activity levels 2,096, RR approximated with OR.
	risk of PASC, 30.0% lower, OR 0.70, $p = 0.046$, high activity levels 388, low activity levels 2,096, musculoskeletal, RR approximated with OR.
	risk of PASC, 39.0% lower, OR 0.61, $p = 0.007$, high activity levels 388, low activity levels 2,096, neurological, RR approximated with OR.
	risk of PASC, 42.0% lower, OR 0.58, $p = 0.03$, high activity levels 388, low activity levels 2,096, respiratory, RR approximated with OR.
	risk of PASC, 5.0% lower, OR 0.95, $p = 0.79$, high activity levels 388, low activity levels 2,096, sensory, RR approximated with OR.
	risk of PASC, 73.0% lower, OR 0.27, $p = 0.19$, high activity levels 388, low activity levels 2,096, digestive, RR approximated with OR.
Saadeh, 10/30/2021, retrospective, Sweden, peer-reviewed, 6 authors, study period March 2020 - June 2020.	risk of symptomatic case, 9.1% lower, OR 0.91, $p = 0.71$, high activity levels 362, low activity levels 225, adjusted per study, inverted to make $OR < 1$ favor high activity levels, 2+ symptoms, Table 8, physically active vs. inactive, multivariable, RR approximated with OR.
	risk of symptomatic case, 3.8% lower, OR 0.96, $p = 0.85$, high activity levels 362, low activity levels 225, adjusted per study, inverted to make $OR < 1$ favor high activity levels, 1+ symptoms, Table 2, model 2, physically active vs. inactive, multivariable, RR approximated with OR.
Salgado-Aranda, 3/14/2022, retrospective, Spain, peer-reviewed, 15 authors, study period 15 February, 2020 - 15 April, 2020.	risk of death, 83.1% lower, HR 0.17, $p = 0.003$, high activity levels 4 of 223 (1.8%), low activity levels 41 of 297 (13.8%), NNT 8.3, inverted to make $HR < 1$ favor high activity levels, active vs. sedentary, Cox proportional hazards.
Sallis, 4/13/2021, retrospective, USA, peer-reviewed, 8 authors.	risk of death, 59.2% lower, RR 0.41, $p = 0.005$, high activity levels 11 of 3,118 (0.4%), low activity levels 170 of 6,984 (2.4%), adjusted per study, inverted to make $RR < 1$ favor high activity levels, odds ratio converted to relative risk, consistently active vs. consistently inactive, multivariable.
	risk of ICU admission, 41.5% lower, RR 0.58, $p = 0.006$, high activity levels 32 of 3,118 (1.0%), low activity levels 195 of 6,984 (2.8%), adjusted per study, inverted to make $RR < 1$ favor high activity levels, odds ratio converted to relative risk, consistently active vs. consistently inactive, multivariable.

	risk of hospitalization, 53.0% lower, RR 0.47, $p < 0.001$, high activity levels 99 of 3,118 (3.2%), low activity levels 732 of 6,984 (10.5%), adjusted per study, inverted to make RR<1 favor high activity levels, odds ratio converted to relative risk, consistently active vs. consistently inactive, multivariable.
Sanchez, 4/25/2023, retrospective, Spain, peer-reviewed, 3 authors, trial NCT04624048 (history).	risk of symptomatic case, 54.1% lower, OR 0.46, $p < 0.001$, inverted to make OR<1 favor high activity levels, exercise vs. no exercise before COVID-19, RR approximated with OR.
Schmidt, 6/21/2023, retrospective, Germany, peer-reviewed, 8 authors, CoCo-Fakt trial.	risk of case, 31.1% lower, OR 0.69, $p = 0.02$, high activity levels 956, low activity levels 2,705, adjusted per study, inverted to make OR<1 favor high activity levels, above guidelines vs. below guidelines, multivariable, RR approximated with OR.
	risk of case, 34.5% lower, OR 0.66, $p = 0.02$, high activity levels 956, low activity levels 1,113, adjusted per study, inverted to make OR<1 favor high activity levels, above guidelines vs. meeting guidelines, multivariable, RR approximated with OR.
	risk of case, 22.7% lower, OR 0.77, $p = 0.02$, high activity levels 3,658, low activity levels 1,680, adjusted per study, inverted to make OR<1 favor high activity levels, exercise vs. no exercise, multivariable, RR approximated with OR.
	risk of case, 21.6% lower, OR 0.78, $p = 0.03$, high activity levels 3,371, low activity levels 1,716, adjusted per study, inverted to make OR<1 favor high activity levels, moderate-to-vigorous vs. low intensity, multivariable, RR approximated with OR.
Steenkamp, 2/9/2022, retrospective, South Africa, peer-reviewed, 10 authors, study period 19 March, 2020 - 30 June, 2021.	risk of death, 42.0% lower, RR 0.58, $p < 0.001$, high activity levels 29,469, low activity levels 13,366, adjusted per study, high activity vs. low activity, poisson regression, multivariable.
	risk of mechanical ventilation, 45.0% lower, RR 0.55, $p < 0.001$, high activity levels 29,469, low activity levels 13,366, adjusted per study, high activity vs. low activity, poisson regression, multivariable.
	risk of ICU admission, 41.0% lower, RR 0.59, $p < 0.001$, high activity levels 29,469, low activity levels 13,366, adjusted per study, high activity vs. low activity, poisson regression, multivariable.
	risk of hospitalization, 34.0% lower, RR 0.66, $p < 0.001$, high activity levels 29,469, low activity levels 13,366, adjusted per study, high activity vs. low activity, poisson regression, multivariable.
Sutkowska, 6/14/2023, prospective, Poland, peer-reviewed, 14 authors, study period 31 January, 2022 - 11 February, 2022, trial NCT05200767 (history).	risk of death, 62.0% lower, HR 0.38, $p = 0.21$, high activity levels 71, low activity levels 60, inverted to make HR<1 favor high activity levels, IPAQ 1/2 vs. IPAQ 0, Cox proportional hazards.
	risk of no recovery, 61.0% lower, HR 0.39, $p = 0.19$, high activity levels 71, low activity levels 60, IPAQ 1/2 vs. IPAQ 0, Cox proportional hazards.
Tavakol, 2/4/2021, retrospective, Iran, peer-reviewed, 9 authors, study period 20 March, 2020 - 24 April, 2020.	risk of severe case, 68.5% lower, RR 0.31, $p = 0.05$, high activity levels 3 of 64 (4.7%), low activity levels 19 of 124 (15.3%), NNT 9.4, adjusted per study, odds ratio converted to relative risk, moderate to high activity versus low activity, multivariable.

Tret'yakov, 10/26/2020, retrospective, Russia, peer-reviewed, 8 authors, excluded in exclusion analyses: unadjusted results with no group details.	risk of severe case, 98.3% lower, RR 0.02, $p = 0.007$, high activity levels 0 of 27 (0.0%), low activity levels 53 of 266 (19.9%), NNT 5.0, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm).
Tsuzuki, 7/5/2022, retrospective, Japan, preprint, 4 authors, study period 1 January, 2022 - 16 May, 2022.	risk of severe case, 56.3% lower, OR 0.44, $p < 0.001$, high activity levels 3,340, low activity levels 1,528, adjusted per study, inverted to make OR<1 favor high activity levels, good vs. poor physical activity status, multivariable, RR approximated with OR.
Wang, 1/31/2024, prospective, United Kingdom, peer-reviewed, 10 authors.	risk of death, 30.0% lower, HR 0.70, $p < 0.001$, high activity levels 57,930, low activity levels 10,966, adjusted per study, ≥ 150 min/wk moderate or ≥ 75 min/wk vigorous vs. < 75 min/wk vigorous, multivariable.
	risk of hospitalization, 12.0% lower, HR 0.88, $p < 0.001$, high activity levels 57,930, low activity levels 10,966, adjusted per study, ≥ 150 min/wk moderate or ≥ 75 min/wk vigorous vs. < 75 min/wk vigorous, multivariable.
	risk of PASC, 14.0% lower, HR 0.86, $p < 0.001$, high activity levels 57,930, low activity levels 10,966, adjusted per study, ≥ 150 min/wk moderate or ≥ 75 min/wk vigorous vs. < 75 min/wk vigorous, multivariable.
Wang (B), 2/6/2023, prospective, USA, peer-reviewed, survey, mean age 64.7, 8 authors, study period April 2020 - November 2021.	risk of PASC, 10.7% lower, RR 0.89, $p = 0.20$, high activity levels 274 of 691 (39.7%), low activity levels 283 of 594 (47.6%), NNT 13, adjusted per study, inverted to make RR<1 favor high activity levels, ≥ 210 vs. 0-30, multivariable, model 2.
	risk of PASC, 49.0% lower, RR 0.51, $p = 0.002$, high activity levels 188, low activity levels 66, 5 or 6 healthy lifestyle factors vs. 0.
Yates, 2/26/2021, retrospective, United Kingdom, peer-reviewed, 7 authors.	risk of death, 45.3% lower, RR 0.55, $p = 0.001$, high activity levels 72 of 163,912 (0.0%), low activity levels 62 of 30,119 (0.2%), adjusted per study, inverted to make RR<1 favor high activity levels, odds ratio converted to relative risk, multivariable.
	risk of severe case, 46.7% lower, RR 0.53, $p < 0.001$, high activity levels 291 of 163,912 (0.2%), low activity levels 180 of 30,119 (0.6%), adjusted per study, inverted to make RR<1 favor high activity levels, odds ratio converted to relative risk, multivariable.
Young, 12/14/2022, retrospective, USA, peer-reviewed, 7 authors, study period 1 January, 2020 - 31 May, 2021.	risk of death, 74.4% lower, OR 0.26, $p < 0.001$, high activity levels 11,279, low activity levels 29,099, inverted to make OR<1 favor high activity levels, always active vs. always inactive, RR approximated with OR.
	risk of death, 65.3% lower, OR 0.35, $p < 0.001$, high activity levels 11,279, low activity levels 83,452, inverted to make OR<1 favor high activity levels, always active vs. mostly inactive, RR approximated with OR.
	risk of death, 47.9% lower, OR 0.52, $p < 0.001$, high activity levels 11,279, low activity levels 42,490, inverted to make OR<1 favor high activity levels, always active vs. some activity, RR approximated with OR.
	risk of death, 35.5% lower, OR 0.65, $p = 0.002$, high activity levels 11,279, low activity levels 27,871, inverted to make OR<1 favor high activity levels, always active vs. consistently active, RR

	approximated with OR.
	risk of hospitalization, 47.6% lower, OR 0.52, $p < 0.001$, high activity levels 11,279, low activity levels 29,099, inverted to make OR<1 favor high activity levels, always active vs. always inactive, RR approximated with OR.
	risk of hospitalization, 41.9% lower, OR 0.58, $p < 0.001$, high activity levels 11,279, low activity levels 83,452, inverted to make OR<1 favor high activity levels, always active vs. mostly inactive, RR approximated with OR.
	risk of hospitalization, 30.1% lower, OR 0.70, $p < 0.001$, high activity levels 11,279, low activity levels 42,490, inverted to make OR<1 favor high activity levels, always active vs. some activity, RR approximated with OR.
	risk of hospitalization, 20.0% lower, OR 0.80, $p < 0.001$, high activity levels 11,279, low activity levels 27,871, inverted to make OR<1 favor high activity levels, always active vs. consistently active, RR approximated with OR.
Yuan, 6/20/2021, retrospective, China, peer-reviewed, 9 authors, study period 15 February, 2020 - 14 March, 2020.	risk of death, 90.5% lower, RR 0.09, $p = 0.09$, high activity levels 0 of 61 (0.0%), low activity levels 6 of 103 (5.8%), NNT 17, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm), excluded in exclusion analyses: excessive unadjusted differences between groups.
	risk of severe case, 70.0% lower, RR 0.30, $p = 0.03$, high activity levels 3 of 61 (4.9%), low activity levels 26 of 103 (25.2%), NNT 4.9, adjusted per study, inverted to make RR<1 favor high activity levels, odds ratio converted to relative risk, multivariable.
Zhang (C), 12/6/2020, retrospective, United Kingdom, peer-reviewed, 9 authors.	risk of death, 26.0% lower, OR 0.74, $p = 0.17$, adjusted per study, AMPA, per SD increase, multivariable, RR approximated with OR.
	risk of case, 18.0% lower, OR 0.82, $p = 0.01$, adjusted per study, AMPA, per SD increase, multivariable, RR approximated with OR.
Šebić, 7/15/2023, retrospective, Bosnia and Herzegovina, peer-reviewed, 5 authors.	risk of oxygen therapy, 89.5% lower, RR 0.11, $p = 0.045$, high activity levels 0 of 53 (0.0%), low activity levels 4 of 47 (8.5%), NNT 12, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm).
	risk of hospitalization, 91.4% lower, RR 0.09, $p = 0.02$, high activity levels 0 of 53 (0.0%), low activity levels 5 of 47 (10.6%), NNT 9.4, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm).
	risk of progression, 83.9% lower, RR 0.16, $p < 0.001$, high activity levels 4 of 53 (7.5%), low activity levels 22 of 47 (46.8%), NNT 2.5, pneumonia.
	no recovery, 47.3% lower, RR 0.53, $p < 0.001$, high activity levels 22 of 53 (41.5%), low activity levels 37 of 47 (78.7%), NNT 2.7, day 14.

Supplementary Data

Supplementary Data

References

1. **Laukkanen** et al., Objectively Assessed Cardiorespiratory Fitness and All-Cause Mortality Risk, *Mayo Clinic Proceedings*, doi:10.1016/j.mayocp.2022.02.029.
2. **Fernandez** et al., Intrahospital supervised exercise training improves survival rate among hypertensive COVID-19 patients, *Journal of Applied Physiology*, doi:10.1152/jappphysiol.00544.2022.
3. **Rahmati** et al., Baseline physical activity is associated with reduced mortality and disease outcomes in COVID-19: A systematic review and meta-analysis, *Reviews in Medical Virology*, doi:10.1002/rmv.2349.
4. **Ezzatvar** et al., Physical activity and risk of infection, severity and mortality of COVID-19: a systematic review and non-linear dose-response meta-analysis of data from 1 853 610 adults, *British Journal of Sports Medicine*, doi:10.1136/bjsports-2022-105733.
5. **Sittichai** et al., Effects of physical activity on the severity of illness and mortality in COVID-19 patients: A systematic review and meta-analysis, *Frontiers in Physiology*, doi:10.3389/fphys.2022.1030568.
6. **Liu** et al., Baseline physical activity and the risk of severe illness and mortality from COVID-19: A dose-response meta-analysis, *Preventive Medicine Reports*, doi:10.1016/j.pmedr.2023.102130.
7. **Halabchi** et al., Association between physical activity and risk of COVID-19 infection or clinical outcomes of the patients with COVID-19; A systematic review and meta-analysis, *Journal of Preventive Medicine and Hygiene*, doi:10.15167/2421-4248/jpmh2023.64.2.2625.
8. **Li** et al., Association of physical activity and the risk of COVID-19 hospitalization: a dose-response meta-analysis, *medRxiv*, doi:10.1101/2022.06.22.22276789.
9. **Guthold** et al., Worldwide trends in insufficient physical activity from 2001 to 2016: a pooled analysis of 358 population-based surveys with 1.9 million participants, *The Lancet Global Health*, doi:10.1016/S2214-109X(18)30357-7.
10. **Nieman** et al., The compelling link between physical activity and the body's defense system, *Journal of Sport and Health Science*, doi:10.1016/j.jshs.2018.09.009.
11. **Kunutsor** et al., High fitness levels, frequent sauna bathing and risk of pneumonia in a cohort study: Are there potential implications for COVID-19?, *European Journal of Clinical Investigation*, doi:10.1111/eci.13490.
12. **Zeraatkar** et al., Consistency of covid-19 trial preprints with published reports and impact for decision making: retrospective review, *BMJ Medicine*, doi:10.1136/bmjmed-2022-0003091.
13. **Davidson** et al., No evidence of important difference in summary treatment effects between COVID-19 preprints and peer-reviewed publications: a meta-epidemiological study, *Journal of Clinical Epidemiology*, doi:10.1016/j.jclinepi.2023.08.011.
14. **Brawner** et al., Inverse Relationship of Maximal Exercise Capacity to Hospitalization Secondary to Coronavirus Disease 2019, *Mayo Clinic Proceedings*, doi:10.1016/j.mayocp.2020.10.003.
15. **de Souza** et al., Association of physical activity levels and the prevalence of COVID-19-associated hospitalization, *Journal of Science and Medicine in Sport*, doi:10.1016/j.jsams.2021.05.011.
16. **Hegazy** et al., Beneficial role of healthy eating Index-2015 score & physical activity on COVID-19 outcomes, *BMC Nutrition*, doi:10.1186/s40795-023-00727-8.
17. **Huang** et al., Reduced Sleep in the Week Prior to Diagnosis of COVID-19 is Associated with the Severity of COVID-19, *Nature and Science of Sleep*, doi:10.2147/NSS.S263488.
18. **Kontopoulou** et al., Exercise Preferences and Benefits in Patients Hospitalized with COVID-19, *Journal of Personalized Medicine*, doi:10.3390/jpm12040645.
19. **Mohsin** et al., Lifestyle and Comorbidity-Related Risk Factors of Severe and Critical COVID-19 Infection: A Comparative Study Among Survived COVID-19 Patients in Bangladesh, *Infection and Drug Resistance*, doi:10.2147/IDR.S331470.
20. **Tret'yakov** et al., COVID-19 in individuals adapted to aerobic exercise, *Pulmonologiya*, doi:10.18093/0869-0189-2020-30-5-553-560.
21. **Yuan** et al., Does pre-existent physical inactivity have a role in the severity of COVID-19?, *Therapeutic Advances in Respiratory Disease*, doi:10.1177/17534666211025221.
22. **Singh** et al., The relationship between viral clearance rates and disease progression in early symptomatic COVID-19: a systematic review and meta-regression analysis, *Journal of Antimicrobial Chemotherapy*, doi:10.1093/jac/dkac045.
23. **Zhang** et al., Effects of the High-Intensity Early Mobilization on Long-Term Functional Status of Patients with Mechanical Ventilation in the Intensive Care Unit, *Critical Care Research and Practice*, doi:10.1155/2024/4118896.
24. **Nindenshuti** et al., Changes in Diet, Physical Activity, Alcohol Consumption, and Tobacco Use in Adults During the COVID-19 Pandemic: A Systematic Review, *INQUIRY: The Journal of Health Care Organization, Provision, and Financing*, doi:10.1177/00469580231175780.
25. **ÓhAiseadha** et al., Unintended Consequences of COVID-19 Non-Pharmaceutical Interventions (NPIs) for Population Health and Health Inequalities, *International Journal of*

- Environmental Research and Public Health, doi:10.3390/ijerph20075223.
26. **Larenas-Linnemann** et al., *Enhancing innate immunity against virus in times of COVID-19: Trying to untangle facts from fictions*, World Allergy Organization Journal, doi:10.1016/j.waojou.2020.100476.
 27. **Dugied** et al., *Multimodal SARS-CoV-2 interactome sketches the virus-host spatial organization*, Communications Biology, doi:10.1038/s42003-025-07933-z.
 28. **Malone** et al., *Structures and functions of coronavirus replication–transcription complexes and their relevance for SARS-CoV-2 drug design*, Nature Reviews Molecular Cell Biology, doi:10.1038/s41580-021-00432-z.
 29. **Murigneux** et al., *Proteomic analysis of SARS-CoV-2 particles unveils a key role of G3BP proteins in viral assembly*, Nature Communications, doi:10.1038/s41467-024-44958-0.
 30. **Lv** et al., *Host proviral and antiviral factors for SARS-CoV-2*, Virus Genes, doi:10.1007/s11262-021-01869-2.
 31. **Lui** et al., *Nsp1 facilitates SARS-CoV-2 replication through calcineurin-NFAT signaling*, Virology, doi:10.1128/mbio.00392-24.
 32. **Niarakis** et al., *Drug-target identification in COVID-19 disease mechanisms using computational systems biology approaches*, Frontiers in Immunology, doi:10.3389/fimmu.2023.1282859.
 33. **Katiyar** et al., *SARS-CoV-2 Assembly: Gaining Infectivity and Beyond*, Viruses, doi:10.3390/v16111648.
 34. **Wu** et al., *Decoding the genome of SARS-CoV-2: a pathway to drug development through translation inhibition*, RNA Biology, doi:10.1080/15476286.2024.2433830.
 35. **c19early.org**, c19early.org/treatments.html.
 36. **c19early.org (B)**, c19early.org/timeline.html.
 37. **Hazan** et al., *Probiotics Counterfeit! Study Finds Most Labels Mislead Customers*, American Journal of Gastroenterology, doi:10.14309/01.ajg.0000950600.34429.5f.
 38. **eurapa.biomedcentral.com**, eurapa.biomedcentral.com/articles/10.1186/s11556-021-00260-2.
 39. **Mateja** et al., *The choice of viral load endpoint in early phase trials of COVID-19 treatments aiming to reduce 28-day hospitalization and/or death*, The Journal of Infectious Diseases, doi:10.1093/infdis/jiaf282.
 40. **Zhang (B)** et al., *What's the relative risk? A method of correcting the odds ratio in cohort studies of common outcomes*, JAMA, 80:19, 1690, doi:10.1001/jama.280.19.1690.
 41. **Altman, D.**, *How to obtain the P value from a confidence interval*, BMJ, doi:10.1136/bmj.d2304.
 42. **Altman (B)** et al., *How to obtain the confidence interval from a P value*, BMJ, doi:10.1136/bmj.d2090.
 43. **Sweeting** et al., *What to add to nothing? Use and avoidance of continuity corrections in meta-analysis of sparse data*, Statistics in Medicine, doi:10.1002/sim.1761.
 44. **Deng, H.**, PyMeta, Python module for meta-analysis, www.pymeta.com/.
 45. **Treanor** et al., *Efficacy and Safety of the Oral Neuraminidase Inhibitor Oseltamivir in Treating Acute Influenza: A Randomized Controlled Trial*, JAMA, 2000, 283:8, 1016-1024, doi:10.1001/jama.283.8.1016.
 46. **McLean** et al., *Impact of Late Oseltamivir Treatment on Influenza Symptoms in the Outpatient Setting: Results of a Randomized Trial*, Open Forum Infect. Dis. September 2015, 2:3, doi:10.1093/ofid/ofv100.
 47. **af Geijerstam** et al., *Fitness, strength and severity of COVID-19: a prospective register study of 1 559 187 Swedish conscripts*, BMJ Open, doi:10.1136/bmjopen-2021-051316.
 48. **Ahmadi** et al., *Lifestyle risk factors and infectious disease mortality, including COVID-19, among middle aged and older adults: Evidence from a community-based cohort study in the United Kingdom*, Brain, Behavior, and Immunity, doi:10.1016/j.bbi.2021.04.022.
 49. **Akbar** et al., *The Association between Lifestyle Factors and COVID-19: Findings from Qatar Biobank*, Nutrients, doi:10.3390/nu16071037.
 50. **Almansour** et al., *The Influence of Physical Activity on COVID-19 Prevention Among Quarantined Individuals: A Case–Control Study*, Journal of Multidisciplinary Healthcare, doi:10.2147/JMDH.S352753.
 51. **Antunes** et al., *The influence of physical activity level on the length of stay in hospital in older men survivors of COVID-19*, Sport Sciences for Health, doi:10.1007/s11332-022-00948-7.
 52. **Baynouna AlKetbi** et al., *Risk Factors for SARS-CoV-2 Infection Severity in Abu Dhabi*, Journal of Epidemiology and Global Health, doi:10.1007/s44197-021-00006-4.
 53. **Beydoun** et al., *Socio-demographic, lifestyle and health characteristics as predictors of self-reported Covid-19 history among older adults: 2006-2020 Health and Retirement Study*, American Journal of Infection Control, doi:10.1016/j.ajic.2022.02.021.
 54. **Bielik** et al., *A Possible Preventive Role of Physically Active Lifestyle during the SARS-CoV-2 Pandemic; Might Regular Cold-Water Swimming and Exercise Reduce the Symptom Severity of COVID-19?*, International Journal of Environmental Research and Public Health, doi:10.3390/ijerph18137158.
 55. **Brandenburg** et al., *Does Higher Self-Reported Cardiorespiratory Fitness Reduce the Odds of Hospitalization From COVID-19?*, Journal of Physical Activity and Health, doi:10.1123/jpah.2020-0817.
 56. **Cardoso** et al., *Patterns of physical activity and SARS-CoV-2 severe pneumonia: A case–control study*, Medicina Clínica, doi:10.1016/j.medcli.2023.04.031.
 57. **Cho** et al., *Physical Activity and the Risk of COVID-19 Infection and Mortality: A Nationwide Population-Based Case-Control Study*, Journal of Clinical Medicine, doi:10.3390/jcm10071539.
 58. **Christensen** et al., *The association of estimated cardiorespiratory fitness with COVID-19 incidence and mortality: A cohort study*, PLOS ONE, doi:10.1371/journal.pone.0250508.
 59. **Eklblom-Bak** et al., *Cardiorespiratory fitness and lifestyle on severe COVID-19 risk in 279,455 adults: a case control study*, International Journal of Behavioral Nutrition and Physical

- Activity, doi:10.1186/s12966-021-01198-5.
60. **Feter** et al., Physical activity and long COVID: findings from the Prospective Study About Mental and Physical Health in Adults cohort, Public Health, doi:10.1016/j.puhe.2023.05.011.
 61. **Frish** et al., The Association of Weight Reduction and Other Variables after Bariatric Surgery with the Likelihood of SARS-CoV-2 Infection, Journal of Clinical Medicine, doi:10.3390/jcm12124054.
 62. **Gao** et al., The impact of individual lifestyle and status on the acquisition of COVID-19: A case—Control study, PLOS ONE, doi:10.1371/journal.pone.0241540.
 63. **Gilley** et al., Risk Factors for COVID-19 in College Students Identified by Physical, Mental, and Social Health Reported During the Fall 2020 Semester: Observational Study Using the Roadmap App and Fitbit Wearable Sensors, JMIR Mental Health, doi:10.2196/34645.
 64. **Green** et al., A higher frequency of physical activity is associated with reduced rates of SARS-CoV-2 infection, European Journal of General Practice, doi:10.1080/13814788.2022.2138855.
 65. **Halabchi (B)** et al., Regular Sports Participation as a Potential Predictor of Better Clinical Outcome in Adult Patients With COVID-19: A Large Cross-Sectional Study, Journal of Physical Activity and Health, doi:10.1123/jpah.2020-0392.
 66. **Hamdan** et al., Risk factors associated with hospitalization owing to COVID-19: a cross-sectional study in Palestine, Journal of International Medical Research, doi:10.1177/03000605211064405.
 67. **Hamer** et al., Lifestyle risk factors, inflammatory mechanisms, and COVID-19 hospitalization: A community-based cohort study of 387,109 adults in UK, Brain, Behavior, and Immunity, doi:10.1016/j.bbi.2020.05.059.
 68. **Hamrouni** et al., Associations of obesity, physical activity level, inflammation and cardiometabolic health with COVID-19 mortality: a prospective analysis of the UK Biobank cohort, BMJ Open, doi:10.1136/bmjopen-2021-055003.
 69. **Hegazy (B)** et al., Beyond probiotic legend: ESSAP gut microbiota health score to delineate SARS-COV-2 infection severity, British Journal of Nutrition, doi:10.1017/S0007114521001926.
 70. **Ho** et al., Modifiable and non-modifiable risk factors for COVID-19, and comparison to risk factors for influenza and pneumonia: results from a UK Biobank prospective cohort study, BMJ Open, doi:10.1136/bmjopen-2020-040402.
 71. **Holt** et al., Risk factors for developing COVID-19: a population-based longitudinal study (COVIDENCE UK), Thorax, doi:10.1136/thoraxjnl-2021-217487.
 72. **Kapusta** et al., Do selected lifestyle parameters affect the severity and symptoms of COVID-19 among elderly patients? The retrospective evaluation of individuals from the STOP-COVID registry of the PoLoCOV study, Journal of Infection and Public Health, doi:10.1016/j.jiph.2022.12.008.
 73. **Latorre-Román** et al., Protective role of physical activity patterns prior to COVID-19 confinement with the severity/duration of respiratory pathologies consistent with COVID-19 symptoms in Spanish populations, Research in Sports Medicine, doi:10.1080/15438627.2021.1937166.
 74. **Lee** et al., Physical activity and the risk of SARS-CoV-2 infection, severe COVID-19 illness and COVID-19 related mortality in South Korea: a nationwide cohort study, British Journal of Sports Medicine, doi:10.1136/bjsports-2021-104203.
 75. **Lengelé** et al., Frailty but not sarcopenia nor malnutrition increases the risk of developing COVID-19 in older community-dwelling adults, Aging Clinical and Experimental Research, doi:10.1007/s40520-021-01991-z.
 76. **Li (B)** et al., Modifiable lifestyle factors and severe COVID-19 risk: a Mendelian randomisation study, BMC Medical Genomics, doi:10.1186/s12920-021-00887-1.
 77. **Lin** et al., Predictors of incident SARS-CoV-2 infections in an international prospective cohort study, BMJ Open, doi:10.1136/bmjopen-2021-052025.
 78. **Lobelo** et al., Clinical, behavioural and social factors associated with racial disparities in COVID-19 patients from an integrated healthcare system in Georgia: a retrospective cohort study, BMJ Open, doi:10.1136/bmjopen-2020-044052.
 79. **Malisoux** et al., Associations between physical activity prior to infection and COVID-19 disease severity and symptoms: results from the prospective Predi-COVID cohort study, BMJ Open, doi:10.1136/bmjopen-2021-057863.
 80. **Maltagliati** et al., Muscle strength explains the protective effect of physical activity against COVID-19 hospitalization among adults aged 50 years and older, Journal of Sports Sciences, doi:10.1080/02640414.2021.1964721.
 81. **Marcus** et al., Predictors of incident viral symptoms ascertained in the era of COVID-19, PLOS ONE, doi:10.1371/journal.pone.0253120.
 82. **Muñoz-Vergara** et al., Prepandemic Physical Activity and Risk of COVID-19 Diagnosis and Hospitalization in Older Adults, JAMA Network Open, doi:10.1001/jamanetworkopen.2023.55808.
 83. **Nguyen** et al., Single and Combinative Impacts of Healthy Eating Behavior and Physical Activity on COVID-19-like Symptoms among Outpatients: A Multi-Hospital and Health Center Survey, Nutrients, doi:10.3390/nu13093258.
 84. **Park** et al., Pre-pandemic physical activity as a predictor of infection and mortality associated with COVID-19: Evidence from the National Health Insurance Service, Frontiers in Public Health, doi:10.3389/fpubh.2023.1072198.
 85. **Paul** et al., Health behaviours the month prior to COVID-19 infection and the development of self-reported long COVID and specific long COVID symptoms: A longitudinal analysis of 1,811 UK adults, medRxiv, doi:10.1101/2022.04.12.22273792.
 86. **Pavlidou** et al., Association of COVID-19 Infection with Sociodemographic, Anthropometric and Lifestyle Factors: A Cross-Sectional Study in an Older Adults' Population Aged over 65 Years Old, Diseases, doi:10.3390/diseases11040165.
 87. **Pitanga** et al., Leisure Time Physical Activity and SARS-CoV-2 Infection among ELSA-Brasil Participants, International Journal of Environmental Research and Public Health, doi:10.3390/ijerph192114155.

88. **Pływaczewska-Jakubowska** et al., *Lifestyle, course of COVID-19, and risk of Long-COVID in non-hospitalized patients*, *Frontiers in Medicine*, doi:10.3389/fmed.2022.1036556.
89. **Reis** et al., *The Association between Lifestyle Risk Factors and COVID-19 Hospitalization in a Healthcare Institution*, *American Journal of Lifestyle Medicine*, doi:10.1177/15598276221135541.
90. **Rocha** et al., *Physical activity status prevents symptoms of long covid: Sulcovid-19 survey*, *BMC Sports Science, Medicine and Rehabilitation*, doi:10.1186/s13102-023-00782-5.
91. **Saadeh** et al., *Associations of pre-pandemic levels of physical function and physical activity with COVID-19-like symptoms during the outbreak*, *Aging Clinical and Experimental Research*, doi:10.1007/s40520-021-02006-7.
92. **Salgado-Aranda** et al., *Influence of Baseline Physical Activity as a Modifying Factor on COVID-19 Mortality: A Single-Center, Retrospective Study*, *Infectious Diseases and Therapy*, doi:10.1007/s40121-021-00418-6.
93. **Sallis** et al., *Physical inactivity is associated with a higher risk for severe COVID-19 outcomes: a study in 48 440 adult patients*, *British Journal of Sports Medicine*, doi:10.1136/bjsports-2021-104080.
94. **Sanchez** et al., *Influence of Physical Exercise on the Severity of COVID-19*, *Fisioterapia*, doi:10.1016/j.ft.2023.04.003.
95. **Schmidt** et al., *Self-Reported Pre-Pandemic Physical Activity and Likelihood of COVID-19 Infection: Data from the First Wave of the CoCo-Fakt Survey*, *Sports Medicine - Open*, doi:10.1186/s40798-023-00592-6.
96. **Steenkamp** et al., *Small steps, strong shield: directly measured, moderate physical activity in 65 361 adults is associated with significant protective effects from severe COVID-19 outcomes*, *British Journal of Sports Medicine*, doi:10.1136/bjsports-2021-105159.
97. **Sutkowska** et al., *Physical Activity Modifies the Severity of COVID-19 in Hospitalized Patients—Observational Study*, *Journal of Clinical Medicine*, doi:10.3390/jcm12124046.
98. **Tavakol** et al., *Relationship between physical activity, healthy lifestyle and COVID-19 disease severity; a cross-sectional study*, *Journal of Public Health*, doi:10.1007/s10389-020-01468-9.
99. **Tsuzuki** et al., *Impact of dementia, living in a long-term care facility, and physical activity status on COVID-19 severity in older adults*, *medRxiv*, doi:10.1101/2022.07.01.22277144.
100. **Wang** et al., *Modifiable lifestyle factors and the risk of post-COVID-19 multisystem sequelae, hospitalization, and death*, *Nature Communications*, doi:10.1038/s41467-024-50495-7.
101. **Wang (B)** et al., *Adherence to Healthy Lifestyle Prior to Infection and Risk of Post-COVID-19 Condition*, *JAMA Internal Medicine*, doi:10.1001/jamainternmed.2022.6555.
102. **Yates** et al., *Obesity, walking pace and risk of severe COVID-19 and mortality: analysis of UK Biobank*, *International Journal of Obesity*, doi:10.1038/s41366-021-00771-z.
103. **Young** et al., *Associations of Physical Inactivity and COVID-19 Outcomes Among Subgroups*, *American Journal of Preventive Medicine*, doi:10.1016/j.amepre.2022.10.007.
104. **Zhang (C)** et al., *Physical activity and COVID-19: an observational and Mendelian randomisation study*, *Journal of Global Health*, doi:10.7189/jogh-10-020514.
105. **Šebić** et al., *Influence of the Level of Physical Activity on Symptoms and Duration of Recovery From Covid-19*, *Sports Science and Health*, doi:10.7251/SSH2301078S.