

Lactoferrin for COVID-19: real-time meta analysis of 8 studies

@CovidAnalysis, March 2024, Version 15
<https://c19early.org/lfmeta.html>

Abstract



Meta analysis using the most serious outcome reported shows 24% [-24-53%] lower risk, without reaching statistical significance. Results are worse for Randomized Controlled Trials and higher quality studies. Early treatment is more effective than late treatment.

3 studies from 3 independent teams in 2 countries show statistically significant improvements.

No treatment or intervention is 100% effective. All practical, effective, and safe means should be used based on risk/benefit analysis. Multiple treatments are typically used in combination, and other treatments are more effective. The quality of non-prescription supplements can vary widely *Crawford, Crighton*.

All data to reproduce this paper and sources are in the appendix.

Lactoferrin for COVID-19




HIGHLIGHTS

Lactoferrin reduces risk for COVID-19 with very low confidence for viral clearance, however increased risk is seen with very low confidence for ventilation and cases.


We show traditional outcome specific analyses and combined evidence from all studies, incorporating treatment delay, a primary confounding factor in COVID-19 studies.

Real-time updates and corrections, transparent analysis with all results in the same format, consistent protocol for 66 treatments.

8 lactoferrin COVID-19 studies



Efficacy in COVID-19 lactoferrin studies (pooled effects)







Figure 1. A. Random effects meta-analysis. This plot shows pooled effects, see the specific outcome analyses for individual outcomes, and the heterogeneity section for discussion. Effect extraction is pre-specified, using the most serious outcome reported. For details of effect extraction see the [appendix](#). **B. Scatter plot showing the most serious outcome in all studies, and for studies within each stage.** Diamonds shows the results of random effects meta-analysis. **C. Results within the context of multiple COVID-19 treatments.** 0.6% of 6,686 proposed treatments show efficacy [c19early.org](#). **D. Timeline of results in lactoferrin studies.**

Introduction

Immediate treatment recommended. SARS-CoV-2 infection primarily begins in the upper respiratory tract and may progress to the lower respiratory tract, other tissues, and the nervous and cardiovascular systems, which may lead to cytokine storm, pneumonia, ARDS, neurological issues *Scardua-Silva, Yang*, cardiovascular complications *Eberhardt*, organ failure, and death. Minimizing replication as early as possible is recommended.

Many treatments are expected to modulate infection. SARS-CoV-2 infection and replication involves the complex interplay of 50+ host and viral proteins and other factors *Note A, Malone, Murigneux, Lv, Lui*, providing many therapeutic targets for which many existing compounds have known activity. Scientists have predicted that over 6,000

compounds may reduce COVID-19 risk [c19early.org](#) (B), either by directly minimizing infection or replication, by supporting immune system function, or by minimizing secondary complications.

Other infections. Efficacy with lactoferrin has been shown for respiratory infections [Pasinato](#).

Other infections. Efficacy with lactoferrin has been shown in preclinical research for respiratory infections [Pasinato](#).

Analysis. We analyze all significant controlled studies of lactoferrin for COVID-19. Search methods, inclusion criteria, effect extraction criteria (more serious outcomes have priority), all individual study data, PRISMA answers, and statistical methods are detailed in Appendix 1. We present random effects meta-analysis results for all studies, studies within each treatment stage, individual outcomes, Randomized Controlled Trials (RCTs), and higher quality studies.

Treatment timing. Figure 2 shows stages of possible treatment for COVID-19. Prophylaxis refers to regularly taking medication before becoming sick, in order to prevent or minimize infection. Early Treatment refers to treatment immediately or soon after symptoms appear, while Late Treatment refers to more delayed treatment.




Figure 2. Treatment stages.

Preclinical Research

2 *In Silico* studies support the efficacy of lactoferrin [Cutone, Miotto](#).

9 *In Vitro* studies support the efficacy of lactoferrin [Alves, Andreu, Cutone, Kobayashi-Sakamoto, Mirabelli, Ostrov, Piacentini, Salaris, Yazawa](#).

An *In Vivo* animal study supports the efficacy of lactoferrin [Berkowitz](#).

Preclinical research is an important part of the development of treatments, however results may be very different in clinical trials. Preclinical results are not used in this paper.

Results

Table 1 summarizes the results for all stages combined, for Randomized Controlled Trials, after exclusions, and for specific outcomes. Table 2 shows results by treatment stage. Figure 3, 4, 5, 6, 7, 8, and 9 show forest plots for random effects meta-analysis of all studies with pooled effects, mortality results, ventilation, hospitalization, recovery, cases, and viral clearance.

	<i>Improvement</i>	<i>Studies</i>	<i>Patients</i>	<i>Authors</i>
All studies	24% [-24-53%]	8	1,419	150
After exclusions	18% [-34-50%]	7	872	132
Randomized Controlled Trials	-18% [-90-27%]	5	687	92
Mortality	17% [-161-73%]	3	921	85
Hospitalization	61% [-164-94%]	2	277	22
Recovery	-7% [-53-25%]	4	549	81
Cases	-36% [-218-42%]	2	259	19
Viral	28% [-22-58%]	3	294	54
RCT mortality	-16% [-115-38%]	2	374	67

Table 1. Random effects meta-analysis for all stages combined, for Randomized Controlled Trials, after exclusions, and for specific outcomes. Results show the percentage improvement with treatment and the 95% confidence interval. *** $p < 0.001$.

	<i>Early treatment</i>	<i>Late treatment</i>	<i>Prophylaxis</i>
All studies	47% [27-61%] ***	19% [-81-64%]	-36% [-218-42%]
After exclusions	47% [27-61%] ***	-6% [-92-41%]	-36% [-218-42%]
Randomized Controlled Trials	-203% [-7216-87%]	-6% [-92-41%]	-36% [-218-42%]
Mortality	-203% [-7216-87%]	37% [-198-87%]	
Hospitalization	61% [-164-94%]		
Recovery	11% [-56-49%]	-30% [-86-9%]	
Cases			-36% [-218-42%]
Viral	28% [-22-58%]		
RCT mortality	-203% [-7216-87%]	-12% [-110-41%]	

Table 2. Random effects meta-analysis results by treatment stage. Results show the percentage improvement with treatment, the 95% confidence interval, and the number of studies for the stage. *** $p < 0.001$.

8 lactoferrin COVID-19 studies

c19early.org
March 2024




Figure 3. Random effects meta-analysis for all studies with pooled effects. This plot shows pooled effects, see the specific outcome analyses for individual outcomes, and the heterogeneity section for discussion. Effect extraction is pre-specified, using the most serious outcome reported. For details of effect extraction see the appendix.

3 lactoferrin COVID-19 mortality results

c19early.org
March 2024




Figure 4. Random effects meta-analysis for mortality results.

1 lactoferrin COVID-19 mechanical ventilation result

c19early.org
March 2024





Figure 5. Random effects meta-analysis for ventilation.

2 lactoferrin COVID-19 hospitalization results

c19early.org
March 2024




¹ CT: study uses combined treatment

Figure 6. Random effects meta-analysis for hospitalization.

4 lactoferrin COVID-19 recovery results

c19early.org
March 2024



¹ CT: study uses combined treatment

Tau² = 0.03, I² = 24.2%, p = 0.72


Figure 7. Random effects meta-analysis for recovery.

2 lactoferrin COVID-19 case results



Figure 8. Random effects meta-analysis for cases.

3 lactoferrin COVID-19 viral clearance results



¹ CT: study uses combined treatment

Tau² = 0.19, I² = 88.6%, p = 0.22

Figure 9. Random effects meta-analysis for viral clearance.

Randomized Controlled Trials (RCTs)

Figure 10 shows a comparison of results for RCTs and non-RCT studies. Figure 11 and 12 show forest plots for random effects meta-analysis of all Randomized Controlled Trials and RCT mortality results. RCT results are included in Table 1 and Table 2.

RCTs have many potential biases. Bias in clinical research may be defined as something that tends to make conclusions differ systematically from the truth. RCTs help to make study groups more similar and can provide a higher level of evidence, however they are subject to many biases ^{Jadad}, and analysis of double-blind RCTs has identified extreme levels of bias ^{Gotzsche}. For COVID-19, the overhead may delay treatment, dramatically compromising efficacy; they may encourage monotherapy for simplicity at the cost of efficacy which may rely on combined or synergistic effects; the participants that sign up may not reflect real world usage or the population that benefits most in terms of age, comorbidities, severity of illness, or other factors; standard of care may be compromised and unable to evolve quickly based on emerging research for new diseases; errors may be made in randomization and medication delivery; and investigators may have hidden agendas or vested interests influencing design, operation, analysis, and the potential for fraud. All of these biases have been observed with COVID-19 RCTs. There is no guarantee that a specific RCT provides a higher level of evidence.

Conflicts of interest for COVID-19 RCTs. RCTs are expensive and many RCTs are funded by pharmaceutical companies or interests closely aligned with pharmaceutical companies. For COVID-19, this creates an incentive to show efficacy for patented commercial products, and an incentive to show a lack of efficacy for inexpensive treatments. The bias is expected to be significant, for example *Als-Nielsen et al.* analyzed 370 RCTs from Cochrane reviews, showing that trials funded by for-profit organizations were 5 times more likely to recommend the

experimental drug compared with those funded by nonprofit organizations. For COVID-19, some major philanthropic organizations are largely funded by investments with extreme conflicts of interest for and against specific COVID-19 interventions.

RCTs for novel acute diseases requiring rapid treatment. High quality RCTs for novel acute diseases are more challenging, with increased ethical issues due to the urgency of treatment, increased risk due to enrollment delays, and more difficult design with a rapidly evolving evidence base. For COVID-19, the most common site of initial infection is the upper respiratory tract. Immediate treatment is likely to be most successful and may prevent or slow progression to other parts of the body. For a non-prophylaxis RCT, it makes sense to provide treatment in advance and instruct patients to use it immediately on symptoms, just as some governments have done by providing medication kits in advance. Unfortunately, no RCTs have been done in this way. Every treatment RCT to date involves delayed treatment. Among the 66 treatments we have analyzed, 63% of RCTs involve very late treatment 5+ days after onset. No non-prophylaxis COVID-19 RCTs match the potential real-world use of early treatments (they may more accurately represent results for treatments that require visiting a medical facility, e.g., those requiring intravenous administration).

RCT bias for widely available treatments. RCTs have a bias against finding an effect for interventions that are widely available — patients that believe they need the intervention are more likely to decline participation and take the intervention. RCTs for lactoferrin are more likely to enroll low-risk participants that do not need treatment to recover, making the results less applicable to clinical practice. This bias is likely to be greater for widely known treatments, and may be greater when the risk of a serious outcome is overstated. This bias does not apply to the typical pharmaceutical trial of a new drug that is otherwise unavailable.

Non-RCT studies have been shown to be reliable. Evidence shows that non-RCT trials can also provide reliable results. *Concato et al.* found that well-designed observational studies do not systematically overestimate the magnitude of the effects of treatment compared to RCTs. *Anglemyer et al.* summarized reviews comparing RCTs to observational studies and found little evidence for significant differences in effect estimates. *Lee et al.* showed that only 14% of the guidelines of the Infectious Diseases Society of America were based on RCTs. Evaluation of studies relies on an understanding of the study and potential biases. Limitations in an RCT can outweigh the benefits, for example excessive dosages, excessive treatment delays, or Internet survey bias could have a greater effect on results. Ethical issues may also prevent running RCTs for known effective treatments. For more on issues with RCTs see *Deaton, Nichol*.

Using all studies identifies efficacy 5.7+ months faster for COVID-19. Currently, 44 of the treatments we analyze show statistically significant efficacy or harm, defined as $\geq 10\%$ decreased risk or $>0\%$ increased risk from ≥ 3 studies. Of the 44 treatments with statistically significant efficacy/harm, 28 have been confirmed in RCTs, with a mean delay of 5.7 months. When considering only low cost treatments, 23 have been confirmed with a delay of 6.9 months. For the 16 unconfirmed treatments, 3 have zero RCTs to date. The point estimates for the remaining 13 are all consistent with the overall results (benefit or harm), with 10 showing $>20\%$. The only treatments showing $>10\%$ efficacy for all studies, but $<10\%$ for RCTs are sotrovimab and aspirin.

Summary. We need to evaluate each trial on its own merits. RCTs for a given medication and disease may be more reliable, however they may also be less reliable. For off-patent medications, very high conflict of interest trials may be more likely to be RCTs, and more likely to be large trials that dominate meta analyses.




Figure 10. Results for RCTs and non-RCT studies.

5 lactoferrin COVID-19 Randomized Controlled Trials

c19early.org

March 2024



Figure 11. Random effects meta-analysis for all Randomized Controlled Trials. This plot shows pooled effects, see the specific outcome analyses for individual outcomes, and the heterogeneity section for discussion. Effect extraction is pre-specified, using the most serious outcome reported. For details of effect extraction see the appendix.

2 lactoferrin COVID-19 RCT mortality results

c19early.org

March 2024




Figure 12. Random effects meta-analysis for RCT mortality results.

Exclusions

To avoid bias in the selection of studies, we analyze all non-retracted studies. Here we show the results after excluding studies with major issues likely to alter results, non-standard studies, and studies where very minimal detail is currently available. Our bias evaluation is based on analysis of each study and identifying when there is a significant chance that limitations will substantially change the outcome of the study. We believe this can be more valuable than checklist-based approaches such as Cochrane GRADE, which may underemphasize serious issues not captured in the checklists, overemphasize issues unlikely to alter outcomes in specific cases (for example, lack of blinding for an objective mortality outcome, or certain specifics of randomization with a very large effect size), and can be easily influenced by potential bias.

The studies excluded are as below. Figure 13 shows a forest plot for random effects meta-analysis of all studies after exclusions.

Rosa, excessive unadjusted differences between groups. Excluded results: no recovery.

Shousha, confounding by indication, unadjusted results and treatment used selectively per official protocol; unadjusted results with no group details.

7 lactoferrin COVID-19 studies after exclusions




Figure 13. Random effects meta-analysis for all studies after exclusions. This plot shows pooled effects, see the specific outcome analyses for individual outcomes, and the heterogeneity section for discussion. Effect extraction is pre-specified, using the most serious outcome reported. For details of effect extraction see the appendix.

Heterogeneity

Heterogeneity in COVID-19 studies arises from many factors including:

Treatment delay. The time between infection or the onset of symptoms and treatment may critically affect how well a treatment works. For example an antiviral may be very effective when used early but may not be effective in late stage disease, and may even be harmful. Oseltamivir, for example, is generally only considered effective for influenza when used within 0-36 or 0-48 hours *McLean, Treanor*. Baloxavir studies for influenza also show that treatment delay is critical — *Ikematsu* report an 86% reduction in cases for post-exposure prophylaxis, *Hayden* show a 33 hour reduction in the time to alleviation of symptoms for treatment within 24 hours and a reduction of 13 hours for treatment within 24-48 hours, and *Kumar* report only 2.5 hours improvement for inpatient treatment.

<i>Treatment delay</i>	<i>Result</i>
Post exposure prophylaxis	86% fewer cases <i>Ikematsu</i>
<24 hours	-33 hours symptoms <i>Hayden</i>
24-48 hours	-13 hours symptoms <i>Hayden</i>
Inpatients	-2.5 hours to improvement <i>Kumar</i>

Table 3. Studies of baloxavir for influenza show that early treatment is more effective.

Figure 14 shows a mixed-effects meta-regression for efficacy as a function of treatment delay in COVID-19 studies from 66 treatments, showing that efficacy declines rapidly with treatment delay. Early treatment is critical for COVID-19.




Figure 14. Early treatment is more effective. Meta-regression showing efficacy as a function of treatment delay in COVID-19 studies from 66 treatments.

Patient demographics. Details of the patient population including age and comorbidities may critically affect how well a treatment works. For example, many COVID-19 studies with relatively young low-comorbidity patients show all patients recovering quickly with or without treatment. In such cases, there is little room for an effective treatment to improve results (as in *López-Medina*).

Effect measured. Efficacy may differ significantly depending on the effect measured, for example a treatment may be very effective at reducing mortality, but less effective at minimizing cases or hospitalization. Or a treatment may have no effect on viral clearance while still being effective at reducing mortality.

Variants. There are many different variants of SARS-CoV-2 and efficacy may depend critically on the distribution of variants encountered by the patients in a study. For example, the Gamma variant shows significantly different characteristics *Faria, Karita, Nonaka, Zavascki*. Different mechanisms of action may be more or less effective depending on variants, for example the viral entry process for the omicron variant has moved towards TMPRSS2-independent fusion, suggesting that TMPRSS2 inhibitors may be less effective *Peacock, Willett*.

Regimen. Effectiveness may depend strongly on the dosage and treatment regimen.

Other treatments. The use of other treatments may significantly affect outcomes, including anything from supplements, other medications, or other kinds of treatment such as prone positioning.

Medication quality. The quality of medications may vary significantly between manufacturers and production batches, which may significantly affect efficacy and safety. *Williams* analyze ivermectin from 11 different sources, showing highly variable antiparasitic efficacy across different manufacturers. *Xu* analyze a treatment from two different manufacturers, showing 9 different impurities, with significantly different concentrations for each manufacturer. Non-prescription supplements may show very wide variations in quality *Crawford, Crighton*.

Pooled outcome analysis. We present both pooled analyses and specific outcome analyses. Notably, pooled analysis often results in earlier detection of efficacy as shown in Figure 15. For many COVID-19 treatments, a reduction in mortality logically follows from a reduction in hospitalization, which follows from a reduction in symptomatic cases, etc. An antiviral tested with a low-risk population may report zero mortality in both arms, however a reduction in severity and improved viral clearance may translate into lower mortality among a high-risk population, and including these results in pooled analysis allows faster detection of efficacy. Trials with high-risk patients may also be restricted due to ethical concerns for treatments that are known or expected to be effective.

Pooled analysis enables using more of the available information. While there is much more information available, for example dose-response relationships, the advantage of the method used here is simplicity and transparency. Note that pooled analysis could hide efficacy, for example a treatment that is beneficial for late stage patients but has no effect on viral replication or early stage disease could show no efficacy in pooled analysis if most studies only examine viral clearance. While we present pooled results, we also present individual outcome analyses, which may be more informative for specific use cases.

Pooled outcomes identify efficacy faster. Currently, 44 of the treatments we analyze show statistically significant efficacy or harm, defined as $\geq 10\%$ decreased risk or $> 0\%$ increased risk from ≥ 3 studies. 88% of treatments showing statistically significant efficacy/harm with pooled effects have been confirmed with one or more specific outcomes, with a mean delay of 3.6 months. When restricting to RCTs only, 50% of treatments showing statistically significant efficacy/harm with pooled effects have been confirmed with one or more specific outcomes, with a mean delay of 6.1 months.

Time when COVID-19 studies showed efficacy

c19early.org

March 2024




Figure 15. The time when studies showed that treatments were effective, defined as statistically significant improvement of $\geq 10\%$ from ≥ 3 studies. Pooled results typically show efficacy earlier than specific outcome results. Results from all studies often shows efficacy much earlier than when restricting to RCTs. Results reflect conditions as used in trials to date, these depend on the population treated, treatment delay, and treatment regimen.

Meta analysis. The distribution of studies will alter the outcome of a meta analysis. Consider a simplified example where everything is equal except for the treatment delay, and effectiveness decreases to zero or below with increasing delay. If there are many studies using very late treatment, the outcome may be negative, even though early treatment is very effective. This may have a greater effect than pooling different outcomes such as mortality and hospitalization. For example a treatment may have 50% efficacy for mortality but only 40% for hospitalization when used within 48 hours. However efficacy could be 0% when used late.

All meta analyses combine heterogeneous studies, varying in population, variants, and potentially all factors above, and therefore may obscure efficacy by including studies where treatment is less effective. Generally, we expect the estimated effect size from meta analysis to be less than that for the optimal case. Looking at all studies is valuable for providing an overview of all research, important to avoid cherry-picking, and informative when a positive result is found despite combining less-optimal situations. However, the resulting estimate does not apply to specific cases such as early treatment in high-risk populations. While we present results for all studies, we also present treatment time and individual outcome analyses, which may be more informative for specific use cases.

Discussion

Results for other viruses. Efficacy with lactoferrin has also been shown for respiratory infections ^{Pasinato}. Efficacy with lactoferrin has also been shown in preclinical research for respiratory infections ^{Pasinato}.

Publication bias. Publishing is often biased towards positive results, however evidence suggests that there may be a negative bias for inexpensive treatments for COVID-19. Both negative and positive results are very important for COVID-19, media in many countries prioritizes negative results for inexpensive treatments (inverting the typical incentive for scientists that value media recognition), and there are many reports of difficulty publishing positive results *Boulware, Meeus, Meneguzzo*. For lactoferrin, there is currently not enough data to evaluate publication bias with high confidence.

One method to evaluate bias is to compare prospective vs. retrospective studies. Prospective studies are more likely to be published regardless of the result, while retrospective studies are more likely to exhibit bias. For example, researchers may perform preliminary analysis with minimal effort and the results may influence their decision to continue. Retrospective studies also provide more opportunities for the specifics of data extraction and adjustments to influence results.

Figure 16 shows a scatter plot of results for prospective and retrospective studies.




Figure 16. Prospective vs. retrospective studies. The diamonds show the results of random effects meta-analysis.

Funnel plot analysis. Funnel plots have traditionally been used for analyzing publication bias. This is invalid for COVID-19 acute treatment trials — the underlying assumptions are invalid, which we can demonstrate with a simple example. Consider a set of hypothetical perfect trials with no bias. Figure 17 plot A shows a funnel plot for a simulation of 80 perfect trials, with random group sizes, and each patient's outcome randomly sampled (10% control event probability, and a 30% effect size for treatment). Analysis shows no asymmetry ($p > 0.05$). In plot B, we add a single typical variation in COVID-19 treatment trials — treatment delay. Consider that efficacy varies from 90% for treatment within 24 hours, reducing to 10% when treatment is delayed 3 days. In plot B, each trial's treatment delay is randomly selected. Analysis now shows highly significant asymmetry, $p < 0.0001$, with six variants of Egger's test all showing $p < 0.05$ *Egger, Harbord, Macaskill, Moreno, Peters, Rothstein, Rücker, Stanley*. Note that these tests fail even though treatment delay is uniformly distributed. In reality treatment delay is more complex — each trial has a different distribution of delays across patients, and the distribution across trials may be biased (e.g., late treatment trials may be more common). Similarly, many other variations in trials may produce asymmetry, including dose, administration, duration of treatment, differences in SOC, comorbidities, age, variants, and bias in design, implementation, analysis, and reporting.




Figure 17. Example funnel plot analysis for simulated perfect trials.

Conflicts of interest. Pharmaceutical drug trials often have conflicts of interest whereby sponsors or trial staff have a financial interest in the outcome being positive. Lactoferrin for COVID-19 lacks this because it is an inexpensive and widely available supplement. In contrast, most COVID-19 lactoferrin trials have been run by physicians on the front lines with the primary goal of finding the best methods to save human lives and minimize the collateral damage caused by COVID-19. While pharmaceutical companies are careful to run trials under optimal conditions (for example, restricting patients to those most likely to benefit, only including patients that can be treated soon after onset when necessary, and ensuring accurate dosing), not all lactoferrin trials represent the optimal conditions for efficacy.

Limitations. Summary statistics from meta analysis necessarily lose information. As with all meta analyses, studies are heterogeneous, with differences in treatment delay, treatment regimen, patient demographics, variants, conflicts of interest, standard of care, and other factors. We provide analyses by specific outcomes and by treatment delay, and we aim to identify key characteristics in the forest plots and summaries. Results should be viewed in the context of study characteristics.

Some analyses classify treatment based on early or late administration, as done here, while others distinguish between mild, moderate, and severe cases. Viral load does not indicate degree of symptoms — for example patients may have a high viral load while being asymptomatic. With regard to treatments that have antiviral properties, timing of treatment is critical — late administration may be less helpful regardless of severity.

Details of treatment delay per patient is often not available. For example, a study may treat 90% of patients relatively early, but the events driving the outcome may come from 10% of patients treated very late. Our 5 day cutoff for early treatment may be too conservative, 5 days may be too late in many cases.

Comparison across treatments is confounded by differences in the studies performed, for example dose, variants, and conflicts of interest. Trials affiliated with special interests may use designs better suited to the preferred outcome.

In some cases, the most serious outcome has very few events, resulting in lower confidence results being used in pooled analysis, however the method is simpler and more transparent. This is less critical as the number of studies increases. Restriction to outcomes with sufficient power may be beneficial in pooled analysis and improve accuracy when there are few studies, however we maintain our pre-specified method to avoid any retrospective changes.

Studies show that combinations of treatments can be highly synergistic and may result in many times greater efficacy than individual treatments alone *Alsaïdi, Andreani, De Forni, Fiaschi, Jeffreys, Jitobaom, Jitobaom (B), Ostrov, Said, Thairu, Wan*. Therefore standard of care may be critical and benefits may diminish or disappear if standard of care does not include certain treatments.

This real-time analysis is constantly updated based on submissions. Accuracy benefits from widespread review and submission of updates and corrections from reviewers. Less popular treatments may receive fewer reviews.

No treatment, vaccine, or intervention is 100% available and effective for all current and future variants. Efficacy may vary significantly with different variants and within different populations. All treatments have potential side effects. Propensity to experience side effects may be predicted in advance by qualified physicians. We do not provide medical advice. Before taking any medication, consult a qualified physician who can compare all options, provide personalized advice, and provide details of risks and benefits based on individual medical history and situations.

Notes. 1 of 8 studies combine treatments. The results of lactoferrin alone may differ. 1 of 5 RCTs use combined treatment. Currently all studies are peer-reviewed.


Reviews. Multiple reviews cover lactoferrin for COVID-19, presenting additional background on mechanisms and related results, including *Campione, Chang, Mattar, Rosa (B)*.

Conclusion

Meta analysis using the most serious outcome reported shows 24% [-24-53%] lower risk, without reaching statistical significance. Results are worse for Randomized Controlled Trials and higher quality studies. Early treatment is more effective than late treatment. 3 studies from 3 independent teams in 2 countries show statistically significant improvements.


Study Notes

Algahtani




Algahtani: RCT 54 hospitalized patients in Egypt, showing no significant differences in recovery with lactoferrin treatment. 200mg lactoferrin orally once daily (group 1) or 200mg lactoferrin orally twice daily (group 2).

Campione




Campione (B): Small prospective study in Italy with 32 lactoferrin patients, 32 SOC, and 28 patients with no treatment, showing significantly faster viral clearance and improved recovery with treatment. Oral and intranasal lactoferrin.

Mann




Mann: RCT 156 mild/moderate COVID-19 patients, 77 treated with hen egg white and bovine colostrum, showing faster recovery of severe symptoms with treatment. There were no significant differences in overall symptom duration, viral clearance, or post-COVID symptoms. Only one participant progressed to severe COVID-19.

Matino




Matino: RCT 218 hospitalized patients in Italy, showing no significant differences with lactoferrin treatment. Authors note that in several previous studies showing clinical improvement, lactoferrin was given at an earlier stage of disease. Authors also note that potential benefits with the late treatment in this study could be masked by other SOC medications - corticosteroids may have masked immunomodulatory effects of lactoferrin, and there may be heparin-dependent reduction in lactoferrin antiviral activity. 800mg oral bovine lactoferrin daily.

Navarro




Navarro: Early terminated low-risk patient prophylaxis RCT in Peru, showing no significant difference in cases with lactoferrin. There were no moderate or severe cases.

Pasinato




Pasinato: RCT 50 preschool children, 25 treated with bovine lactoferrin (bLf) prophylaxis, showing significantly lower frequency and duration of respiratory infections during the active phase with treatment. The only COVID-19 specific results reported are the number as patients with COVID, 1 vs. 2 for treatment vs. control. bLf 400mg bid for 4 months.

Rosa



Rosa: Retrospective survey based study in Italy with 82 patients treated with lactoferrin, and 39 control patients, showing significantly faster viral clearance with treatment. There was no significant difference in recovery time overall, however the treatment group had significantly more moderate condition patients (39% versus 8%), and improved recovery was seen with treatment as age increased. Median dose for asymptomatic patients was 400mg/day, for paucisymptomatic patients 600mg/day, and for moderate condition patients 1000mg three times a day.

Shousha



Shousha: Retrospective 547 hospitalized COVID+ patients in Egypt, showing lower mortality with lactoferrin treatment (without statistical significance).

Appendix 1. Methods and Data

We perform ongoing searches of PubMed, medRxiv, Europe PMC, ClinicalTrials.gov, The Cochrane Library, Google Scholar, Research Square, ScienceDirect, Oxford University Press, the reference lists of other studies and meta-analyses, and submissions to the site c19early.org. Search terms are lactoferrin and COVID-19 or SARS-CoV-2. Automated searches are performed twice daily, with all matches reviewed for inclusion. All studies regarding the use of lactoferrin for COVID-19 that report a comparison with a control group are included in the main analysis. Sensitivity analysis is performed, excluding studies with major issues, epidemiological studies, and studies with minimal available information. This is a living analysis and is updated regularly.

We extracted effect sizes and associated data from all studies. If studies report multiple kinds of effects then the most serious outcome is used in pooled analysis, while other outcomes are included in the outcome specific analyses. For example, if effects for mortality and cases are both reported, the effect for mortality is used, this may be different to the effect that a study focused on. If symptomatic results are reported at multiple times, we used the latest time, for example if mortality results are provided at 14 days and 28 days, the results at 28 days have preference. Mortality alone is preferred over combined outcomes. Outcomes with zero events in both arms are not used, the next most serious outcome with one or more events is used. For example, in low-risk populations with no mortality, a reduction in mortality with treatment is not possible, however a reduction in hospitalization, for example, is still valuable. Clinical outcomes are considered more important than viral test status. When basically all patients recover in both treatment and control groups, preference for viral clearance and recovery is given to results mid-recovery where available. After most or all patients have recovered there is little or no room for an effective treatment to do better, however faster recovery is valuable. If only individual symptom data is available, the most serious symptom has priority, for example difficulty breathing or low SpO₂ is more important than cough. When results provide an odds ratio, we compute the relative risk when possible, or convert to a relative risk according to *Zhang*. Reported confidence intervals and *p*-values were used when available, using adjusted values when provided. If multiple types of adjustments are reported propensity score matching and multivariable regression has preference over propensity score matching or weighting, which has preference over multivariable regression. Adjusted results have preference over unadjusted results for a more serious outcome when the adjustments significantly alter results. When needed, conversion between reported *p*-values and confidence intervals followed *Altman, Altman (B)*, and Fisher's exact test was used to calculate *p*-values for event data. If continuity correction for zero values is required, we use the reciprocal of the opposite arm with the sum of the correction factors equal to 1 *Sweeting*. Results are expressed with RR < 1.0 favoring treatment, and using the risk of a negative outcome when applicable (for example, the risk of death rather than the risk of survival). If studies only report relative continuous values such as relative times, the ratio of the time for the treatment group versus the time for the control group is used. Calculations are done in Python (3.12.2) with scipy (1.12.0), pythonmeta (1.26), numpy (1.26.4), statsmodels (0.14.1), and plotly (5.19.0).

Forest plots are computed using PythonMeta *Deng* with the DerSimonian and Laird random effects model (the fixed effect assumption is not plausible in this case) and inverse variance weighting. Results are presented with 95% confidence intervals. Heterogeneity among studies was assessed using the *I*² statistic. Mixed-effects meta-regression results are computed with R (4.1.2) using the metafor (3.0-2) and rms (6.2-0) packages, and using the most serious sufficiently powered outcome. For all statistical tests, a *p*-value less than 0.05 was considered statistically significant. Grobid 0.8.0 is used to parse PDF documents.

We have classified studies as early treatment if most patients are not already at a severe stage at the time of treatment (for example based on oxygen status or lung involvement), and treatment started within 5 days of the onset of symptoms. If studies contain a mix of early treatment and late treatment patients, we consider the treatment time of patients contributing most to the events (for example, consider a study where most patients are treated early but late treatment patients are included, and all mortality events were observed with late treatment patients). We note that a shorter time may be preferable. Antivirals are typically only considered effective when used within a shorter timeframe, for example 0-36 or 0-48 hours for oseltamivir, with longer delays not being effective *McLean, Treanor*.

We received no funding, this research is done in our spare time. We have no affiliations with any pharmaceutical companies or political parties.

A summary of study results is below. Please submit updates and corrections at <https://c19early.org/lfmeta.html>.

Early treatment

Effect extraction follows pre-specified rules as detailed above and gives priority to more serious outcomes. For pooled analyses, the first (most serious) outcome is used, which may differ from the effect a paper focuses on. Other outcomes are used in outcome specific analyses.

Campione (B), 10/19/2021, prospective, Italy, peer-reviewed, 32 authors.

time to viral-, 47.5% lower, relative time 0.53, *p* < 0.001, treatment 32, control 32, vs. SOC.

	time to viral-, 56.3% lower, relative time 0.44, $p < 0.001$, treatment 32, control 28, vs. untreated.
<p><i>Mann</i>, 7/20/2023, Double Blind Randomized Controlled Trial, placebo-controlled, South Africa, peer-reviewed, 14 authors, study period 28 July, 2021 - 5 July, 2022, this trial uses multiple treatments in the treatment arm (combined with bovine colostrum and egg white) - results of individual treatments may vary, trial DOH-27-062021-9191.</p>	risk of death, 202.6% higher, RR 3.03, $p = 0.49$, treatment 1 of 77 (1.3%), control 0 of 79 (0.0%), continuity correction due to zero event (with reciprocal of the contrasting arm).
	risk of severe case, 202.6% higher, RR 3.03, $p = 0.49$, treatment 1 of 77 (1.3%), control 0 of 79 (0.0%), continuity correction due to zero event (with reciprocal of the contrasting arm).
	risk of hospitalization, 48.7% lower, RR 0.51, $p = 1.00$, treatment 1 of 77 (1.3%), control 2 of 79 (2.5%), NNT 81, COVID-19.
	risk of hospitalization, 48.7% lower, RR 0.51, $p = 0.68$, treatment 2 of 77 (2.6%), control 4 of 79 (5.1%), NNT 41, all cause.
	relative symptom score, 25.8% better, RR 0.74, $p = 0.24$, treatment 77, control 79, mid-recovery, day 7.
	relative symptom score, 60.0% better, RR 0.40, $p = 0.047$, treatment 77, control 79, day 28.
	relative symptom score, 2.3% better, RR 0.98, $p = 0.91$, treatment 77, control 79, day 14.
	relative symptom score, 18.4% better, RR 0.82, $p = 0.40$, treatment 77, control 79, day 11-13.
	relative symptom score, 4.4% better, RR 0.96, $p = 0.75$, treatment 77, control 79, day 4.
	risk of no viral clearance, 11.2% higher, RR 1.11, $p = 0.36$, treatment 49 of 60 (81.7%), control 36 of 49 (73.5%), day 11-3.
	risk of PASC, 7.6% lower, RR 0.92, $p = 0.84$, treatment 15 of 67 (22.4%), control 16 of 66 (24.2%), NNT 54, day 42.
<p><i>Rosa</i>, 9/21/2021, retrospective, Italy, peer-reviewed, 8 authors, study period October 2020 - March 2021.</p>	risk of hospitalization, 75.6% lower, RR 0.24, $p = 0.32$, treatment 0 of 82 (0.0%), control 1 of 39 (2.6%), NNT 39, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm).
	recovery time, 40.0% higher, relative time 1.40, $p = 0.50$, treatment 82, control 39, excluded in exclusion analyses: excessive unadjusted differences between groups.
	time to viral-, 39.4% lower, relative time 0.61, $p = 0.02$, treatment 82, control 39, inverted to make $RR < 1$ favor treatment, Cox regression, primary outcome.

Late treatment

Effect extraction follows pre-specified rules as detailed above and gives priority to more serious outcomes. For pooled analyses, the first (most serious) outcome is used, which may differ from the effect a paper focuses on. Other outcomes are used in outcome specific analyses.

<p><i>Algahtani</i>, 8/19/2021, Randomized Controlled Trial, Egypt, peer-reviewed, 6 authors, study period 8 July, 2020 - 18 September, 2020.</p>	<p>risk of unresolved fever, 25.0% lower, RR 0.75, $p = 1.00$, treatment 3 of 36 (8.3%), control 2 of 18 (11.1%), NNT 36, day 7.</p>
	<p>risk of unresolved fatigue, 33.3% lower, RR 0.67, $p = 0.67$, treatment 4 of 36 (11.1%), control 3 of 18 (16.7%), NNT 18, day 7.</p>
	<p>risk of unresolved cough, no change, RR 1.00, $p = 1.00$, treatment 8 of 36 (22.2%), control 4 of 18 (22.2%), day 7.</p>
	<p>risk of unresolved headache, no change, RR 1.00, $p = 1.00$, treatment 4 of 36 (11.1%), control 2 of 18 (11.1%), day 7.</p>
	<p>risk of unresolved loss of smell/taste, 25.0% lower, RR 0.75, $p = 0.72$, treatment 6 of 36 (16.7%), control 4 of 18 (22.2%), NNT 18, day 7.</p>
<p><i>Matino</i>, 3/4/2023, Double Blind Randomized Controlled Trial, placebo-controlled, Italy, peer-reviewed, 53 authors, study period January 2021 - May 2021, average treatment delay 6.0 days, trial NCT04847791 (history) (LAC).</p>	<p>risk of death, 11.5% higher, RR 1.12, $p = 0.85$, treatment 18 of 113 (15.9%), control 15 of 105 (14.3%), day 28.</p>
	<p>risk of death, 23.9% higher, RR 1.24, $p = 0.69$, treatment 16 of 113 (14.2%), control 12 of 105 (11.4%), day 14.</p>
	<p>risk of mechanical ventilation, 44.5% higher, RR 1.45, $p = 0.39$, treatment 14 of 113 (12.4%), control 9 of 105 (8.6%).</p>
	<p>risk of death/ICU, 6.2% higher, RR 1.06, $p = 0.87$, treatment 24 of 113 (21.2%), control 21 of 105 (20.0%).</p>
<p><i>Shousha</i>, 10/28/2021, retrospective, Egypt, peer-reviewed, 18 authors, study period 15 April, 2020 - 29 July, 2020, excluded in exclusion analyses: confounding by indication, unadjusted results and treatment used selectively per official protocol; unadjusted results with no group details.</p>	<p>not reaching NEWS2 ≤ 2 or discharge within 14 days, 33.6% higher, RR 1.34, $p = 0.12$, treatment 46 of 113 (40.7%), control 32 of 105 (30.5%).</p>
	<p>risk of death, 79.1% lower, RR 0.21, $p = 0.11$, treatment 1 of 46 (2.2%), control 52 of 501 (10.4%), NNT 12, unadjusted.</p>

Prophylaxis

Effect extraction follows pre-specified rules as detailed above and gives priority to more serious outcomes. For pooled analyses, the first (most serious) outcome is used, which may differ from the effect a paper focuses on. Other outcomes are used in outcome specific analyses.

<p><i>Navarro</i>, 12/7/2022, Double Blind Randomized Controlled Trial, placebo-controlled, Peru, peer-reviewed, median age 37.0, 14 authors, study period October 2020 - February 2021, trial NCT04526821 (history) (LF-COVID).</p>	<p>risk of symptomatic case, 58.7% higher, RR 1.59, $p = 0.34$, treatment 11 of 104 (10.6%), control 7 of 105 (6.7%).</p>
	<p>risk of case, 23.4% higher, RR 1.23, $p = 0.65$, treatment 11 of 104 (10.6%), control 9 of 105 (8.6%).</p>
<p><i>Pasinato</i>, 2/15/2024, Randomized Controlled Trial, Italy, peer-reviewed, mean age 4.2, 5 authors.</p>	<p>risk of case, 50.0% lower, RR 0.50, $p = 1.00$, treatment 1 of 25 (4.0%), control 2 of 25 (8.0%), NNT 25.</p>

Supplementary Data

Supplementary Data

Footnotes

- a. Viral infection and replication involves attachment, entry, uncoating and release, genome replication and transcription, translation and protein processing, assembly and budding, and release. Each step can be disrupted by therapeutics.

References

1. **Algahtani** et al., *The Prospect of Lactoferrin Use as Adjunctive Agent in Management of SARS-CoV-2 Patients: A Randomized Pilot Study*, *Medicina*, doi:10.3390/medicina57080842.
2. **Als-Nielsen** et al., *Association of Funding and Conclusions in Randomized Drug Trials*, *JAMA*, doi:10.1001/jama.290.7.921.
3. **Alsaidi** et al., *Griffithsin and Carrageenan Combination Results in Antiviral Synergy against SARS-CoV-1 and 2 in a Pseudoviral Model*, *Marine Drugs*, doi:10.3390/md19080418.
4. **Altman**, D., *How to obtain the P value from a confidence interval*, *BMJ*, doi:10.1136/bmj.d2304.
5. **Altman (B)** et al., *How to obtain the confidence interval from a P value*, *BMJ*, doi:10.1136/bmj.d2090.
6. **Alves** et al., *Inhibition of SARS-CoV-2 Infection in Vero Cells by Bovine Lactoferrin under Different Iron-Saturation States*, *Pharmaceuticals*, doi:10.3390/ph16101352.
7. **Andreani** et al., *In vitro testing of combined hydroxychloroquine and azithromycin on SARS-CoV-2 shows synergistic effect*, *Microbial Pathogenesis*, doi:10.1016/j.micpath.2020.104228.
8. **Andreu** et al., *Liposomal Lactoferrin Exerts Antiviral Activity against HCoV-229E and SARS-CoV-2 Pseudoviruses In Vitro*, *Viruses*, doi:10.3390/v15040972.
9. **Anglemyer** et al., *Healthcare outcomes assessed with observational study designs compared with those assessed in randomized trials*, *Cochrane Database of Systematic Reviews* 2014, Issue 4, doi:10.1002/14651858.MR000034.pub2.
10. **Berkowitz** et al., *Sigma Receptor Ligands Prevent COVID Mortality In Vivo: Implications for Future Therapeutics*, *International Journal of Molecular Sciences*, doi:10.3390/ijms242115718.
11. **Boulware**, D., *Comments regarding paper rejection*, twitter.com/boulware_dr/status/1311331372884205570.
12. **c19early.org**, c19early.org/timeline.html.
13. **c19early.org (B)**, c19early.org/treatments.html.
14. **Campione** et al., *Lactoferrin as Protective Natural Barrier of Respiratory and Intestinal Mucosa against Coronavirus Infection and Inflammation*, *International Journal of Molecular Sciences*, doi:10.3390/ijms21144903.
15. **Campione (B)** et al., *Lactoferrin as Antiviral Treatment in COVID-19 Management: Preliminary Evidence*, *International Journal of Environmental Research and Public Health*, doi:10.3390/ijerph182010985.
16. **Chang** et al., *Lactoferrin as potential preventative and adjunct treatment for COVID-19*, *International Journal of Antimicrobial Agents*, doi:10.1016/j.ijantimicag.2020.106118.
17. **Concato** et al., *NEJM*, 342:1887-1892, doi:10.1056/NEJM200006223422507.
18. **Crawford** et al., *Analysis of Select Dietary Supplement Products Marketed to Support or Boost the Immune System*, *JAMA Network Open*, doi:10.1001/jamanetworkopen.2022.26040.

19. **Crighton** et al., *Toxicological screening and DNA sequencing detects contamination and adulteration in regulated herbal medicines and supplements for diet, weight loss and cardiovascular health*, Journal of Pharmaceutical and Biomedical Analysis, doi:10.1016/j.jpba.2019.112834.
20. **Cutone** et al., *Lactoferrin binding to Sars-CoV-2 Spike glycoprotein protects host from infection, inflammation and iron dysregulation.*, Research Square, doi:10.21203/rs.3.rs-1605740/v1.
21. **De Forni** et al., *Synergistic drug combinations designed to fully suppress SARS-CoV-2 in the lung of COVID-19 patients*, PLoS ONE, doi:10.1371/journal.pone.0276751.
22. **Deaton** et al., *Understanding and misunderstanding randomized controlled trials*, Social Science & Medicine, 210, doi:10.1016/j.socscimed.2017.12.005.
23. **Deng**, H., *PyMeta*, Python module for meta-analysis, www.pymeta.com/.
24. **Eberhardt** et al., *SARS-CoV-2 infection triggers pro-atherogenic inflammatory responses in human coronary vessels*, Nature Cardiovascular Research, doi:10.1038/s44161-023-00336-5.
25. **Egger** et al., *Bias in meta-analysis detected by a simple, graphical test*, BMJ, doi:10.1136/bmj.315.7109.629.
26. **Faria** et al., *Genomics and epidemiology of the P.1 SARS-CoV-2 lineage in Manaus, Brazil*, Science, doi:10.1126/science.abh2644.
27. **Fiaschi** et al., *In Vitro Combinatorial Activity of Direct Acting Antivirals and Monoclonal Antibodies against the Ancestral B.1 and BQ.1.1 SARS-CoV-2 Viral Variants*, Viruses, doi:10.3390/v16020168.
28. **Gøtzsche**, P., *Bias in double-blind trials*, Doctoral Thesis, University of Copenhagen, www.scientificfreedom.dk/2023/05/16/bias-in-double-blind-trials-doctoral-thesis/.
29. **Harbord** et al., *A modified test for small-study effects in meta-analyses of controlled trials with binary endpoints*, Statistics in Medicine, doi:10.1002/sim.2380.
30. **Hayden** et al., *Baloxavir Marboxil for Uncomplicated Influenza in Adults and Adolescents*, New England Journal of Medicine, doi:10.1056/NEJMoa1716197.
31. **Ikematsu** et al., *Baloxavir Marboxil for Prophylaxis against Influenza in Household Contacts*, New England Journal of Medicine, doi:10.1056/NEJMoa1915341.
32. **Jadad** et al., *Randomized Controlled Trials: Questions, Answers, and Musings, Second Edition*, doi:10.1002/9780470691922.
33. **Jeffreys** et al., *Remdesivir-ivermectin combination displays synergistic interaction with improved in vitro activity against SARS-CoV-2*, International Journal of Antimicrobial Agents, doi:10.1016/j.ijantimicag.2022.106542.
34. **Jitobaom** et al., *Favipiravir and Ivermectin Showed in Vitro Synergistic Antiviral Activity against SARS-CoV-2*, Research Square, doi:10.21203/rs.3.rs-941811/v1.
35. **Jitobaom (B)** et al., *Synergistic anti-SARS-CoV-2 activity of repurposed anti-parasitic drug combinations*, BMC Pharmacology and Toxicology, doi:10.1186/s40360-022-00580-8.
36. **Karita** et al., *Trajectory of viral load in a prospective population-based cohort with incident SARS-CoV-2 G614 infection*, medRxiv, doi:10.1101/2021.08.27.21262754.
37. **Kobayashi-Sakamoto** et al., *Bovine lactoferrin suppresses the cathepsin-dependent pathway of SARS-CoV-2 entry in vitro*, International Dairy Journal, doi:10.1016/j.idairyj.2023.105805.
38. **Kumar** et al., *Combining baloxavir marboxil with standard-of-care neuraminidase inhibitor in patients hospitalised with severe influenza (FLAGSTONE): a randomised, parallel-group, double-blind, placebo-controlled, superiority trial*, The Lancet Infectious Diseases, doi:10.1016/S1473-3099(21)00469-2.
39. **Lee** et al., *Analysis of Overall Level of Evidence Behind Infectious Diseases Society of America Practice Guidelines*, Arch Intern Med., 2011, 171:1, 18-22, doi:10.1001/archinternmed.2010.482.
40. **López-Medina** et al., *Effect of Ivermectin on Time to Resolution of Symptoms Among Adults With Mild COVID-19: A Randomized Clinical Trial*, JAMA, doi:10.1001/jama.2021.3071.

41. **Lui** et al., *Nsp1 facilitates SARS-CoV-2 replication through calcineurin-NFAT signaling*, *Virology*, doi:10.1128/mbio.00392-24.
42. **Lv** et al., *Host proviral and antiviral factors for SARS-CoV-2*, *Virus Genes*, doi:10.1007/s11262-021-01869-2.
43. **Macaskill** et al., *A comparison of methods to detect publication bias in meta-analysis*, *Statistics in Medicine*, doi:10.1002/sim.698.
44. **Malone** et al., *Structures and functions of coronavirus replication–transcription complexes and their relevance for SARS-CoV-2 drug design*, *Nature Reviews Molecular Cell Biology*, doi:10.1038/s41580-021-00432-z.
45. **Mann** et al., *Hen egg white bovine colostrum supplement reduces symptoms of mild/moderate COVID-19: a randomized control trial*, *Future Science OA*, doi:10.2144/fsoa-2023-0024.
46. **Matino** et al., *Effect of Lactoferrin on Clinical Outcomes of Hospitalized Patients with COVID-19: The LAC Randomized Clinical Trial*, *Nutrients*, doi:10.3390/nu15051285.
47. **Mattar** et al., *Natural resources to control COVID-19: could lactoferrin amend SARS-CoV-2 infectivity?*, *PeerJ*, doi:10.7717/peerj.11303.
48. **McLean** et al., *Impact of Late Oseltamivir Treatment on Influenza Symptoms in the Outpatient Setting: Results of a Randomized Trial*, *Open Forum Infect. Dis.* September 2015, 2:3, doi:10.1093/ofid/ofv100.
49. **Meeus**, G., *Online Comment*, twitter.com/gertmeeus_MD/status/1386636373889781761.
50. **Meneguesso**, A., *Médica defende tratamento precoce da Covid-19*, www.youtube.com/watch?v=X5FCrIm_19U.
51. **Miotto** et al., *Molecular Mechanisms Behind Anti SARS-CoV-2 Action of Lactoferrin*, *Frontiers in Molecular Biosciences*, doi:10.3389/fmolb.2021.607443.
52. **Mirabelli** et al., *Morphological cell profiling of SARS-CoV-2 infection identifies drug repurposing candidates for COVID-19*, *Proceedings of the National Academy of Sciences*, doi:10.1073/pnas.2105815118.
53. **Moreno** et al., *Assessment of regression-based methods to adjust for publication bias through a comprehensive simulation study*, *BMC Medical Research Methodology*, doi:10.1186/1471-2288-9-2.
54. **Murigneux** et al., *Proteomic analysis of SARS-CoV-2 particles unveils a key role of G3BP proteins in viral assembly*, *Nature Communications*, doi:10.1038/s41467-024-44958-0.
55. **Navarro** et al., *Bovine lactoferrin for the prevention of COVID-19 infection in health care personnel: a double-blinded randomized clinical trial (LF-COVID)*, *BioMetals*, doi:10.1007/s10534-022-00477-3.
56. **Nichol** et al., *Challenging issues in randomised controlled trials*, *Injury*, 2010, doi: 10.1016/j.injury.2010.03.033, [www.injuryjournal.com/article/S0020-1383\(10\)00233-0/fulltext](http://www.injuryjournal.com/article/S0020-1383(10)00233-0/fulltext).
57. **Nonaka** et al., *SARS-CoV-2 variant of concern P.1 (Gamma) infection in young and middle-aged patients admitted to the intensive care units of a single hospital in Salvador, Northeast Brazil, February 2021*, *International Journal of Infectious Diseases*, doi:10.1016/j.ijid.2021.08.003.
58. **Ostrov** et al., *Highly Specific Sigma Receptor Ligands Exhibit Anti-Viral Properties in SARS-CoV-2 Infected Cells*, *Pathogens*, doi:10.3390/pathogens10111514.
59. **Pasinato** et al., *Lactoferrin in the Prevention of Recurrent Respiratory Infections in Preschool Children: A Prospective Randomized Study*, *Children*, doi:10.3390/children11020249.
60. **Peacock** et al., *The SARS-CoV-2 variant, Omicron, shows rapid replication in human primary nasal epithelial cultures and efficiently uses the endosomal route of entry*, *bioRxiv*, doi:10.1101/2021.12.31.474653.
61. **Peters**, J., *Comparison of Two Methods to Detect Publication Bias in Meta-analysis*, *JAMA*, doi:10.1001/jama.295.6.676.
62. **Piacentini** et al., *Lactoferrin Inhibition of the Complex Formation between ACE2 Receptor and SARS CoV-2 Recognition Binding Domain*, *International Journal of Molecular Sciences*, doi:10.3390/ijms23105436.
63. **Rosa** et al., *Ambulatory COVID-19 Patients Treated with Lactoferrin as a Supplementary Antiviral Agent: A Preliminary Study*, *Journal of Clinical Medicine*, doi:10.3390/jcm10184276.

64. **Rosa (B)** et al., *An overview on in vitro and in vivo antiviral activity of lactoferrin: its efficacy against SARS-CoV-2 infection*, BioMetals, doi:10.1007/s10534-022-00427-z.
65. **Rothstein**, H., *Publication Bias in Meta-Analysis: Prevention, Assessment and Adjustments*, www.wiley.com/en-ae/Publication+Bias+in+Meta+Analysis:+Prevention,+Assessment+and+Adjustments-p-9780470870143.
66. **Rücker** et al., *Arcsine test for publication bias in meta-analyses with binary outcomes*, Statistics in Medicine, doi:10.1002/sim.2971.
67. **Said** et al., *The effect of Nigella sativa and vitamin D3 supplementation on the clinical outcome in COVID-19 patients: A randomized controlled clinical trial*, Frontiers in Pharmacology, doi:10.3389/fphar.2022.1011522.
68. **Salaris** et al., *Protective Effects of Lactoferrin against SARS-CoV-2 Infection In Vitro*, Nutrients, doi:10.3390/nu13020328.
69. **Scardua-Silva** et al., *Microstructural brain abnormalities, fatigue, and cognitive dysfunction after mild COVID-19*, Scientific Reports, doi:10.1038/s41598-024-52005-7.
70. **Shousha** et al., *Hepatic and gastrointestinal disturbances in Egyptian patients infected with coronavirus disease 2019: A multicentre cohort study*, World Journal of Gastroenterology, doi:10.3748/wjg.v27.i40.6951.
71. **Stanley** et al., *Meta-regression approximations to reduce publication selection bias*, Research Synthesis Methods, doi:10.1002/jrsm.1095.
72. **Sweeting** et al., *What to add to nothing? Use and avoidance of continuity corrections in meta-analysis of sparse data*, Statistics in Medicine, doi:10.1002/sim.1761.
73. **Thairu** et al., *A Comparison of Ivermectin and Non Ivermectin Based Regimen for COVID-19 in Abuja: Effects on Virus Clearance, Days-to-discharge and Mortality*, Journal of Pharmaceutical Research International, doi:10.9734/jpri/2022/v34i44A36328.
74. **Treanor** et al., *Efficacy and Safety of the Oral Neuraminidase Inhibitor Oseltamivir in Treating Acute Influenza: A Randomized Controlled Trial*, JAMA, 2000, 283:8, 1016-1024, doi:10.1001/jama.283.8.1016.
75. **Wan** et al., *Synergistic inhibition effects of andrographolide and baicalin on coronavirus mechanisms by downregulation of ACE2 protein level*, Scientific Reports, doi:10.1038/s41598-024-54722-5.
76. **Willett** et al., *The hyper-transmissible SARS-CoV-2 Omicron variant exhibits significant antigenic change, vaccine escape and a switch in cell entry mechanism*, medRxiv, doi:10.1101/2022.01.03.21268111.
77. **Williams**, T., *Not All Ivermectin Is Created Equal: Comparing The Quality of 11 Different Ivermectin Sources*, Do Your Own Research, doyourownresearch.substack.com/p/not-all-ivermectin-is-created-equal.
78. **Xu** et al., *A study of impurities in the repurposed COVID-19 drug hydroxychloroquine sulfate by UHPLC-Q/TOF-MS and LC-SPE-NMR*, Rapid Communications in Mass Spectrometry, doi:10.1002/rcm.9358.
79. **Yang** et al., *SARS-CoV-2 infection causes dopaminergic neuron senescence*, Cell Stem Cell, doi:10.1016/j.stem.2023.12.012.
80. **Yazawa** et al., *Evaluation of SARS-CoV-2 isolation in cell culture from nasal/nasopharyngeal swabs or saliva specimens of patients with COVID-19*, Research Square, doi:10.21203/rs.3.rs-2676422/v1.
81. **Zavascki** et al., *Advanced ventilatory support and mortality in hospitalized patients with COVID-19 caused by Gamma (P.1) variant of concern compared to other lineages: cohort study at a reference center in Brazil*, Research Square, doi:10.21203/rs.3.rs-910467/v1.
82. **Zhang** et al., *What's the relative risk? A method of correcting the odds ratio in cohort studies of common outcomes*, JAMA, 80:19, 1690, doi:10.1001/jama.280.19.1690.